

TELE-HEALTH SOLUTIONS IN PRACTICE: A FEASIBILITY STUDY OF THEIR
ADVANTAGES, LIMITATIONS, AND ADOPTION POTENTIAL

by

Rohit Jain

DISSERTATION

Presented to the Swiss School of Business and Management Geneva

In Partial Fulfillment

Of the Requirements

For the Degree

DOCTOR OF BUSINESS ADMINISTRATION

SWISS SCHOOL OF BUSINESS AND MANAGEMENT GENEVA

MARCH, 2026

TELE-HEALTH SOLUTIONS IN PRACTICE: A FEASIBILITY STUDY OF THEIR
ADVANTAGES, LIMITATIONS, AND ADOPTION POTENTIAL


by

Rohit Jain

Supervised by

Dr. Atul Pati Tripathi

APPROVED BY



Dissertation chair

RECEIVED/APPROVED BY:

Renee Goldstein Osmic

Admissions Director

Dedication

I respectfully dedicate this thesis to my parents and wife, whose unwavering support, encouragement, and sacrifices have been the cornerstone of my academic and personal pursuits. Their constant faith in my abilities has been a source of strength and inspiration throughout this journey.

I also dedicate this work to my esteemed teachers and mentors, whose invaluable guidance, wisdom, and encouragement have significantly contributed to my intellectual growth.

Acknowledgements

I would like to express my profound gratitude to my supervisor and guide, **Dr. Atul Pati Tripathi**, for their continuous guidance, insightful suggestions, and unwavering support throughout the course of this research. Their expertise and mentorship have been instrumental in shaping this work.

I am sincerely thankful to **SSBM** for providing the necessary academic environment, resources, and facilities required to complete this study.

I would also like to acknowledge the support and cooperation of my faculty members, and colleagues, whose constructive feedback and encouragement have greatly contributed to this research.

I extend my heartfelt appreciation to my family for their constant encouragement, patience, and moral support during the course of this work.

Finally, I wish to thank all those who, directly or indirectly, along with upGrad Buddy, contributed to the successful completion of this thesis.

ABSTRACT

TELE-HEALTH SOLUTIONS IN PRACTICE: A FEASIBILITY STUDY OF THEIR ADVANTAGES, LIMITATIONS, AND ADOPTION POTENTIAL

Rohit Jain

2026

Dissertation Chair: <Chair's Name>

Co-Chair: <If applicable. Co-Chair's Name>

The fast-paced innovation of digital technology was a game-changer in the field of healthcare delivery, and it made “telehealth” a sustainable and significant concept, which contributed to the improvement of the three pillars of Healthcare (HC): accessibility, affordability, and quality. The researcher was interested in finding out the practicality, acceptability, and “telehealth” services by looking at the users' viewpoint, their experiences, and the factors that determine their willingness to try and keep using it. A mixed-method research design was adopted; the first part of the design was the gathering of quantitative data through a Google Forms survey consisting of a fixed number of questions and a convenience sample size of 150 participants; the second part was collecting qualitative data from research study, reports, and journals that had been published before. Among the different methods of statistics for both inferential and descriptive use were

frequency analysis, descriptive statistics, chi-square tests, correlation tests, and Mann-Whitney U tests. Content analysis was the technique used for the qualitative part of the data analysis. The results indicated that “telehealth” has not only become a temporary option but also been integrated into the modern HC system. The respondents were extremely satisfied with the convenience, time-saving, and access to services the new system provided. However, other problems like technical constraints, limited physical examinations, and data security still exist. Altogether, these results indicate that “telehealth” can be very promising in improving HC delivery and equity, though stable policies, stable infrastructure, and thorough education of both users and providers are required to make “telehealth” a regularly incorporated part of the global HC systems.

Keywords: “Telehealth”, Digital Healthcare, Technology Adoption, Healthcare Accessibility, Feasibility Study, Healthcare Innovation.

TABLE OF CONTENTS

List of Tables	ix
List of Figures	xi
List of Abbreviations	xii
CHAPTER I: INTRODUCTION.....	1
1.1 Introduction.....	1
1.2 Research Problem	9
1.3 Purpose of Research.....	15
1.4 Significance of the Study	16
1.5 Research Purpose and Questions	17
1.6 Business & Strategic Context of Telehealth	19
CHAPTER II: REVIEW OF LITERATURE	25
2.1 Introduction.....	25
2.2 Theoretical Framework.....	26
2.3 Conceptual Framework and Definitions of “Telehealth”	33
2.4 Global and National Trends in “telehealth” Adoption.....	41
2.5 Advantages and Opportunities of “telehealth.”	46
2.6 Challenges and Limitations of “telehealth” Implementation.....	55
2.7 Future Prospects and Emerging Directions in “telehealth” ..	62
2.8 Synthesis of Literature and Research Gaps	69
2.9 Telehealth Business Models and Strategic Management.....	93
CHAPTER III: METHODOLOGY	105
3.1 Overview of the Research Problem	105
3.2 Operationalisation of Theoretical Constructs	107
3.3 Research Purpose and Questions	109
3.4 Research Design.....	110
3.5 Population and Sample	111
3.6 Participant Selection	112
3.7 Instrumentation	114
3.8 Data Collection Procedures.....	116
3.9 Data Analysis	117
3.10 Research Design Limitations	120
3.11 Conclusion	122
CHAPTER IV: RESULTS.....	124

4.1	Frequency Analysis.....	124
4.2	Descriptives.....	132
4.3	Hypothesis Testing.....	133
4.4	Summary of Findings.....	143
CHAPTER V: DISCUSSION.....		149
5.1	Discussion of Results.....	149
5.2	Discussion of Research Question One	151
5.3	Discussion of Research Question Two	153
5.4	Discussion of Research Question Three	157
5.5	Discussion of Research Question Four	159
5.6	Discussion of Research Question Five	160
5.7	Strategic Analysis Using Business Strategy Frameworks ..	163
5.8	Feasibility Evaluation Using TELOS Models	166
5.9	Financial Feasibility Analysis of Telehealth Implementation	171
5.10	Telehealth Readiness Assessment Tool (TRAT): A Practical Framework for Healthcare Organisations	175
5.11	Strategic Implications for Healthcare Organisations	178
5.12	Business Model Innovation Opportunities.....	180
5.13	Change Management Framework for Telehealth Implementation	181
CHAPTER VI: SUMMARY, IMPLICATIONS, AND RECOMMENDATIONS...		184
6.1	Summary	184
6.2	Telehealth Performance Measurement Framework	185
6.3	Implications.....	187
6.4	Strategy Recommendations-Stakeholders.	189
6.5	Conclusion	210
REFERENCES		214
APPENDIX A:.....		239
APPENDIX B: DATASET		241

LIST OF TABLES

Table 3.1: Research Variables or Constructs	109
Table 3.2: Inclusion and Exclusion Criteria for Participant Selection.....	113
Table 4.1: Have you ever used “telehealth” services before?.....	124
Table 4.2: If you have used “telehealth” services, how satisfied were you with the experience?	125
Table 4.3: What is the main reason you use “telehealth” services?.....	126
Table 4.4: How do you feel about the quality of care received through “telehealth” services compared to in-person care?	127
Table 4.5: Have you ever experienced technical difficulties during a “telehealth” appointment?.....	128
Table 4.6: In your opinion, what are the advantages of “telehealth” services? ...	129
Table 4.7: In your opinion, what are the disadvantages of “telehealth” services?	130
Table 4.8: Would you be willing to use “telehealth” services again in the future?	131
Table 4.9: Descriptive Statistics	132
Table 4.10: Hypothesis Test Summary	133
Table 4.11: One-Sample Chi-Square Test Summary.....	134
Table 4.12: One-Sample Chi-Square Test Summary.....	135
Table 4.13: Correlations.....	137
Table 4.14: Ranks	138
Table 4.15: Test Statistics	139
Table 4.16: Correlations.....	141
Table 4.17: Correlations.....	142
Table 5.1: SWOT Analysis	163
Table 5.2: TELOS Feasibility Evaluation.....	166
Table 5.3: TRAT: Organisational Dimensions (7 Dimensions)	175
Table 5.4: Telehealth Implementation Success Factors Model (TISFM) Factors.....	177
Table 5.5: Summary of Feasibility & Strategy Findings	183
Table 6.1: Telehealth Performance Dashboard KPIs.....	186

LIST OF FIGURES

Figure 1.1: “Telemedicine” Workflow	1
Source: (Self-Created)	1
Figure 1.2: “Telemedicine” Market Size 2024 to 2034 (USD Billion)	2
Figure 2.1: The “technology acceptance model”	27
Source: (Davis, 1989)	27
Figure 2.2: The “unified technology acceptance and use of technology theory” (UTAUT)	33
Source: (Venkatesh et al., 2003)	33
Figure 3.1: Conceptual Model of the study	115
Source: (Self-Created)	115
Figure 4.1: Have you ever used “telehealth” services before?	124
Figure 4.2: If you have used “telehealth” services, how satisfied were you with the experience?	125
Figure 4.3: What is the main reason you use “telehealth” services?	126
Figure 4.4: How do you feel about the quality of care received through “telehealth” services compared to in-person care?	127
Figure 4.5: Have you ever experienced technical difficulties during a “telehealth” appointment?	128
Figure 4.6: In your opinion, what are the advantages of “telehealth” services? ..	129
Figure 4.7: In your opinion, what are the disadvantages of “telehealth” services?	130
Figure 4.8: Would you be willing to use “telehealth” services again in the future?	131
Figure 4.9: Have you ever used “telehealth” services before?	135
Figure 4.10: Would you be willing to use “telehealth” services again in the future?	136
Figure 5.1: Telehealth Adoption and Satisfaction Model (TASM)	150
Source: (Self-Created)	150
Figure 5.2: Telehealth Implementation Success Factors Model (TISFM)	178
Source: (Self-Created)	178
Figure 6.1: Telehealth performance dashboard	187

LIST OF ABBREVIATIONS

Abbreviations	Full Form
HC	Healthcare
IoMT	Internet of Medical Things
CAGR	Compound Annual Growth Rate
TAM	Technology Acceptance Model
AI	Artificial Intelligence
DOI	Diffusion of Innovations Theory
UTAUT	Unified Theory of Acceptance and Use of Technology
EDMS	Electronic Document Management Systems
UIMS	University Information Management System
CVDs	Cardiovascular Diseases
HIT	Health Information Technology
GPs	General Practitioners
RPM	Remote Patient Monitoring
VR/AR	Virtual and Augmented Reality
GDPR	General Data Protection Regulation
CNN	Convolutional Neural Network
ASD	Autism Spectrum Disorder
EHRs	Electronic Health Records
MCQs	Multiple-Choice Questions

CHAPTER I: INTRODUCTION

1.1 Introduction

Telehealth uses digital communication technologies to deliver healthcare services to patients who need remote medical care from their providers. The initial purpose of telehealth is to increase healthcare accessibility rather than focusing on its technical organisation, but it has evolved into a vital operational model that current healthcare systems use to deliver services (Haleem et al., 2021). Over 60% of US Healthcare (HC) facilities already use “telehealth” in some form, demonstrating its tremendous traction in HC systems globally (Tuckson et al., 2017). A growing number of prominent HC organisations, including Intermountain HC, Geisinger Health System, Partners HC, and the Veterans Health Administration, have adopted “telehealth” interventions.




Figure 1.1: “Telemedicine” Workflow

Source: (Self-Created)

Figure 1.1 shows a typical “telemedicine” workflow, with the electronic transmission of medical data and the stages needed in remote consultations, to help explain the operational aspects of “telemedicine”.




Figure 1.2: “Telemedicine” Market Size 2024 to 2034 (USD Billion)

Source: (Research, 2025)

As shown in Figure 1.2, the global “telemedicine” market is projected to expand from 160.13 billion USD in 2025 to more than 709.69 billion USD in 2034, representing a commendable CAGR of 17.99%. Estimates place the North American “telemedicine” market size at USD 46.14 billion in 2024, with a predicted CAGR of 18.16% for the projected year. The market size and estimates are based on revenue in millions or billions of US dollars, with 2024 serving as the base year.

Although the telehealth implementation can have great benefits, it is associated with serious organisational challenges. Healthcare facilities need to consider telehealth as a promising investment that influences the structure of operations, cost management, and positioning. The telehealth initiatives may not be effectively utilised without a clear knowledge of the user acceptance dynamics, even though the technology is ready.

The purpose of this study is not a large-scale descriptive report on telehealth technologies around the globe. Rather, it aims at detecting the behavioural determinants, which determine telehealth adoption, especially the perceived usefulness and perceived ease of use as moderators of behavioural intention. Using the Technology Acceptance Model (TAM), the study analyses the issue of telehealth adoption as a decoding problem in the context of healthcare managers and policymakers.

The research will examine the evolution of “telehealth” from both a national and an international perspective while evaluating its merits and demerits in the subsequent sections. In an effort to assist decision makers and health care investors in capturing the full potential of “telehealth” as a remedy for today's health care problems, the study gives the health care sectors advice on how to smoothly incorporate “telemedicine” and eHealth. This study can inform managerial decision-making by providing evidence-based understanding of the conditions that condition the sustainable deployment of telehealth through the analysis of the perceptual factors that determine its adoption. The discoveries are expected to help healthcare leaders make the most of telehealth investment according to organisational strategy and long-term performance goals.

Introduction to “telehealth” Solutions

“Telehealth” solutions intend to make HC services more accessible, efficient, and of better quality by providing a variety of HC services through digital technology at a distance (Haleem et al., 2021). Among the methods that fall within this category are live video conferencing and other synchronous modalities, as well as asynchronous techniques such as store-and-forward systems, remote patient monitoring via wearable technology, health applications for mobile devices, diagnostic tools using artificial intelligence, virtual reality treatments, and blockchain-based secure data management for HC transactions. Telehealth provides better access to medical services while enabling healthcare systems to operate more flexibly. The system needs both patients and providers to continue using it for its success. The complete success of telehealth depends on three factors, which include its technological capabilities, how users view it, how prepared organisations are to implement it, and how their strategic plans match with it.

“Telehealth” was quickly adopted due to the outbreak (Chu et al., 2021; Demeke et al., 2021). Concerns about long-term viability, equal access, and “telehealth's place in the

HC system emerge as many groups' reliance on emergency medical services during the pandemic winds down (Shachar et al., 2020). The COVID-19 pandemic increased the use of telehealth around the world and made it a necessity to introduce it quickly and as a part of regular healthcare. Nevertheless, expediency-based growth is not automatically sustainable in the long term. With the pressures caused by the pandemic under control, the healthcare organisations are now faced with the challenge of understanding whether or not telehealth can be strategically integrated into the regular care models instead of being a crisis-response tool (Shaver, 2022). It is possible that the quick acceptance of “telemedicine” during the pandemic emergency phase is to blame for the large variation in practices, definitions of “telehealth” and “telemedicine”, and state regulations that have resulted from the emergency use of “telehealth” (Omboni et al., 2022; Roy et al., 2021). These new and enlarged adoptions have occasionally involved waivers from the standard regulatory procedures; these waivers are currently under review. At the conclusion of the emergency adoption period, there is a chance to reassess the criteria and components of “telehealth”. This may uncover the less obvious danger of maintaining less-than-ideal practices that were implemented during the pandemic's emergency phase.

Some publications differentiate between “telehealth” and “telemedicine” when defining “telemedicine”, while others use the words synonymously (Ansarian & Baharlouei, 2023; Fatehi & Wootton, 2012). “Telemedicine” is a method of diagnosing and treating patients remotely over telecommunications networks, whereas “telehealth” is commonly considered a subset of “telemedicine” and encompasses all services that are inherently related to this (Amjad et al., 2023; Mathews et al., 2023). In this study, the digital-based remote healthcare services are broadly described as telehealth or telemedicine. It is not about the terminological differences but rather about the behavioural

and organisational variables that have an impact on the adoption and continued use of such services.

In order to better organise content and promote more accurate communication within the field, standardised terminology might be useful in clinical disciplines, just like in other fields (Andrews et al., 2016; R. Bashshur et al., 2011). Even before the pandemic, “telemedicine” was underutilised in both rural and urban settings, with particular gaps in coverage in the fields of neurology and critical care (Chu et al., 2021; Herasevich & Subramanian, 2019). When patients have improved interactions and their expectations are satisfied by their doctors, they are more likely to be active participants in their care. This could be an unintended result of defining language in the “telehealth” sector (Dickinson et al., 2017; Flickinger et al., 2013). As the field of “telehealth” advances, better communication between researchers, legislators, patients, and providers will be attainable with a common definition of the term.

Recent years have seen tremendous change in the worldwide “telehealth” scene as a result of the fast development of digital technology, the broad usage of smartphones, and the increasing need for remote HC access, which has been worsened by the COVID-19 epidemic. The rise of telehealth globally is indicative of a wider shift in the model of healthcare delivery, with digital platforms becoming more and more useful in complementing the face-to-face services (Ezeamii et al., 2024). Nevertheless, the forecasts of market size growth do not always guarantee the effective exploitation and the organization's success. Strategic decision-makers should then go further to consider telehealth not just on a growth basis but also on the behavioral considerations of whether the investments will create long-term value. The developments highlight the growing global presence of “telehealth” and its essential potential to change HC systems in the direction of a more inclusive and resilient system.

Even with such a great growth trend, the use of “telehealth” and sustainability has remained a challenge in various ways. People who use telehealth services after the initial surge show a primary management problem because their acceptance of telehealth services, their ability to use telehealth services, and their planned use of telehealth services need proper management. Organisations need to develop their technological capabilities to manage telehealth services because they need to combine their strategic plans with their current user demand and organisational resources (Su et al., 2024). Organisations face ongoing difficulties because people lack digital skills, and they worry about their privacy, and they see digital tools as having limited benefits. The behavioural and perceptual elements show that researchers need to study telehealth systems by using a complete theoretical framework, which will help them understand how users adopt these systems. These changes point to the fact that although “telehealth” is already part of the standard HC delivery system, the processes for the sustainability of “telehealth” implementations will demand careful handling of the technology, regulations, and the practice, that is, the patients and the providers. Furthermore, hurdles like a lack of digital skills, the need for securing cyberspace, ensuring privacy of data, and the disparity of internet access, especially among the low-income and rural population, are still some of the major hindrances to the “telehealth” equity issue. Nonetheless, the telehealth's contribution to the future of HC delivery seems to be a solid one, together with the growing demand for the hybrid care models that combine the digital and the traditional in-person care. The global HC system will benefit from the increased access, efficiency, and quality of “telehealth” solutions.

This study, therefore, approaches telehealth not as a technological phenomenon but as a strategic adoption decision shaped by user perceptions and organizational considerations. The research shows how telehealth initiatives achieve sustainable

implementation through their examination of perceived usefulness and perceived ease of use, which serve as the main factors driving people to use telehealth services.

Importance of Feasibility Analysis in “telehealth”

The feasibility analysis is a key step in the successful deployment of “telehealth” solutions because it delivers a systematic study of the viability, the practicability, and the impact of the proposed interventions (Gregory et al., 2023). Feasibility evaluations in “telehealth” consider a large number of factors, including but not limited to technical capability, financial viability, regulatory compliance, organisational readiness, and patient willingness. Such studies allow HC providers, policymakers, and technology developers to foresee potential issues, remove the hazards, and make the right choices based on their knowledge prior to large-scale implementation (Sittig et al., 2020). To illustrate, getting the proper technical infrastructure that guarantees a trustworthy internet link, secure data transfer, and communication with the current HC systems may result in significant financial savings, which would be otherwise wasted in fixing a failed implementation. On the same note, financial feasibility studies assist in establishing whether “telehealth” programs are economical, expandable, and can deliver a favourable payback on the investment by HC organisations (Snoswell et al., 2020). Also, assessing patient readiness and acceptance, which implies the assessment of the level of digital literacy, trust in technology, and willingness to use virtual care, will make “telehealth” solutions easy to use and adoptable by many. Finally, extensive feasibility analysis will offer a blueprint to implement “telehealth” programs that are feasible, sustainable, and have the potential to enhance HC access, performance, and quality among various groups of people in various environments.

Increased reliance on “telemedicine” and other forms of online health care is one way in which the COVID-19 pandemic altered the delivery of patient care (Bitar &

Alismail, 2021; Garfan et al., 2021). In order to assist in preventing the spread of the SARS-CoV-2 infection, the virtual modality decreased face-to-face patient encounters. There is also uncertainty about the future of “telemedicine” as a vital patient care modality after COVID-19 becomes endemic, despite its widespread use in many medical specialties at present (Seifert et al., 2020; Shirke et al., 2020). Knowing what helps and what hinders the successful adoption of “telemedicine” is an important consideration (Anthony et al., 2018; Şahin et al., 2024) when “telemedicine” is most effective, how both patients and providers feel about it, and how widespread its use is (Mudiyanselage et al., 2019; Seifert et al., 2020; Shirke et al., 2020; Wootton, 2012).

The most effective aspect of carrying out a thorough feasibility analysis is that it allows one to not only comprehend how successful the “telehealth” programs can be but also to predict and overcome challenges prior to implementation (Gregory et al., 2023). A study conducted in the past has revealed that looking at possibilities first not only increases the adoption and continuation of the program, but also, in a resource-limited environment, it plays an important role. For instance, in doing a feasibility assessment, the criteria for the physical setup generally entail having reliable internet access, the necessary hardware, and interoperability with the current electronic health records, which cannot be avoided in “telehealth” (C. S. Kruse et al., 2017). In addition, the financial feasibility studies support the organisations in recognising the areas of cost-effectiveness, budgeting, and sustainability, which together make it possible for the “telehealth” programs to be of long-term value to both the providers and the patients. When the stakeholders focus on these areas during the early stages of the planning process, they are able to come up with evidence-backed strategies that decrease the risks involved in the operations and, at the same time, increase the effectiveness of the programs.

Feasibility analysis, in addition, contributes significantly to the understanding of patient acceptance and participation, which is one of the factors that determine the desired HC outcomes. The recognition of patient attitudes, for instance, digital literacy, willingness to use “telehealth” services, and perceived mistrust of virtual care, will help the HC organisations to come up with user-friendly solutions that meet community needs (Bober et al., 2024). A few real-world examples illustrate this point: it was shown that in a feasibility study in the rural areas of Malaysia, the evaluation of patient readiness and the assessment of the need for technical support were among the most reinforcing factors leading to the increased adoption rates and the corresponding satisfaction associated with “telehealth” services (Nazar et al., 2024). Similarly, the viability assessments related to the COVID-19 pandemic in the United States indicated that the prompt recognition of logistical and regulatory barriers facilitated the program implementation and widened the use of “telehealth” services (Bestsenny et al., 2021). The findings underline that the feasibility study is not a one-time activity but rather a strategic tool that guarantees “telehealth” solutions are admissible, sustainable, and bring about the desired level of health care improvements among various populations.

1.2 Research Problem

As indicated in previous studies, using technology—in this example, “telemedicine”—to alleviate health issues requires considering a broad range of interactions. Since the 1990s, sociological research on technological innovation has supported the idea that the understanding of innovativeness must be approached from a holistic point of view, taking interactions among all of its elements into consideration. In actuality, technology adoption models such as the Technology Acceptance Model (TAM) have shown that technological innovation is much more than a planned and linear process.

Technology innovation is a complicated system that reacts to trial-and-error procedures, operates disruptively, and rarely follows a set timeline.

Three related developments are presently influencing “telehealth”. The first is how “telehealth” has evolved from being used to make medical care more accessible to being more convenient and eventually less expensive (Lin et al., 2018). The second is the expansion of “telehealth” to include episodic and chronic disorders in addition to acute ones (Lin et al., 2018). The third is the expansion of “telehealth” from hospitals and remote clinics to home and portable electronics (Lin et al., 2018). From a patient's point of view, Telehealth’s primary goal is to improve access to treatment, and as such, it has traditionally improved access to health care for ailments and populations for which care was otherwise unavailable. Programs to give treatment to people in the military, jails, and rural areas have been among the early and persistent applications of “telehealth” (Lin et al., 2018).

The Internet is facilitating the easy delivery of health care in addition to expanding access, much as it has done for travel, shopping, and banking. For the "most prevalent, most annoying, most inconvenient" ailments, several institutions, from academic health facilities to start-ups, now provide affordable virtual visits (less than \$50 each visit), round-the-clock. In comparison, it takes 20 days on average to get a doctor's appointment, which, when combined with travel time and waiting time, takes 2 hours. Given the increased need to bend the cost curve, “telemedicine” may give intense treatments to the 20% of people who are responsible for 80% of HC costs. The United States Senate Committee on Finance put it this way: "Traditionally, 'telehealth' has been seen as a tool to enhance access to treatments, but attention is developing to determine if 'telehealth' has the potential to cut health care costs" (Y. T. Yang, 2016).

More lately, “telemedicine” has grown to cover care for episodic diseases like sinusitis, using a variety of care models including medical assistant visits to schools, video

calls, phone calls, and online algorithms (R. L. Bashshur & Shannon, 2009). “telehealth” applications for chronic diseases have previously been restricted exclusively to usually adaptive surveillance (e.g., text messages) or telephone help, with the significant exception of mental health. In 141 randomised, controlled studies of “telehealth” therapies for chronic illnesses, for instance, just 10 included videoconferencing with a doctor. Despite this scant data foundation, “telehealth” is becoming increasingly popular for treating various chronic illnesses, which impact 140 million people in the US and are responsible for 80% of all medical costs (R. L. Bashshur & Shannon, 2009). Future models will expand on the largely colloquial variant of today into one that incorporates rich data transmission from remote observing (through the usage of wearable detectors and mobile diagnostic tools, including such electrocardiograms), skills training of patients, and regular virtual and in-person visits from doctors, nursing staff, practitioners, and counsellors (R. L. Bashshur & Shannon, 2009).

The third “telehealth” development is the shift in treatment away from hospitals and clinics. Initial “telehealth” applications provided patient treatment in settings like hospitals and satellite clinics, typically requiring pricey technology gear and on-site clinical or technical assistance. “Telehealth” is quickly relocating to the home because of widely accessible broadband and portable diagnostic technology (Lin et al., 2018). The patient-centred medical home will increasingly serve as the patient's residence for those with chronic diseases, including the 2 million elderly people who are effectively housebound. Even care for acute disorders like pneumonia and stroke²⁴ is shifting from the emergency department to the doorstep or bedroom, thanks to the use of video visits in the ambulance or at home. The trend in banking, where services were relocated from the bank lobby to mobile devices through automated teller machines and the Internet, may be seen in the

provision of health care to people in retail clinics or homes and over the phone (Lin et al., 2018).

By facilitating physical separation, monitoring symptoms and outbreaks, and assisting policymakers in predicting requirements and choosing suitable and timely measures, “telehealth” plays both direct and indirect roles in limiting the spread of illnesses (Lin et al., 2018).

“Telehealth” administration, though, demands modifications to working procedures and data curation; it is not only about technology. Changing work procedures has been one of the previous challenges to “telehealth”. To facilitate functioning without physicality in the workspace in order to safeguard patients and HC personnel, “telemedicine” technologies have been widely implemented throughout the pandemic (Dorsey & Topol, 2016).

To preserve patient privacy, the most recent data security and encryption methods must be implemented. To protect data security and prevent the release of sensitive personal information, patients must be informed of the privacy decisions they are making (Fagherazzi et al., 2020). The interests of the individual must be weighed against the interests of the public when deciding on data-sharing policies, keeping in mind that various cultures often place different values on these interests. “Telehealth” solutions must be simple to use and maintain, which includes having reliable internet connectivity, to maximise digital inclusion. Organisations must take into account both the requirements and customs of immigrants and those of permanent residents. In order to be inclusive, HC organisations and individuals must be able to pay it, which may call for strategic finance model modifications (Fagherazzi et al., 2020).

Remote areas with limited access to traditional HC practitioners have historically been the early adopters of “telehealth”. Results and experiences with “telehealth” rely on

specific design considerations and elements, including health literacy, digital literacy, and the degree of clinical care pathway integration. Wharton and colleagues highlighted that there is "scarce evidence of their influence on clinical outcomes and quality of care" when discussing virtual primary care services (Topol et al., 2015). Improvements in clinical outcomes may not be Telehealth's primary benefit, though. Instead, those advancements may result from better systems that provide more affordable HC. To reach true economies of scale, "telehealth" solutions should be used worldwide. Organisations must cooperate and understand what, when, where, why, and how properly in order to maximise the long-term advantages of "telehealth". Governments must assist the "telehealth" sector in creating and evaluating unique, safe, and quick "telehealth" solutions. To achieve digital inclusion, data security, and solutions that are clear, adaptable, and suited to user demands, the industry must collaborate with experts and patients (Summerfelt et al., 2015).

It will be easier to change ingrained work routines with this customisation. However, new skills and information may be empowering for physicians, patients, and caregivers alike. It is important to take into account the financial burden of care placed on patients and their families. "Telehealth" can speed up health literacy in the same way COVID-19 has sped up digital literacy (Snoswell et al., 2020).

Remembering that care is a human and social activity while imagining a "telehealth"-enabled future is important since many clinical problems need physical exams or actions. In order to properly utilise these technologies, HC systems and lifestyles must be modified, and inadequate investment has been made in their development. The COVID-19 pandemic offers a significant opportunity and impetus for the creation of cutting-edge "telehealth" technologies that have the potential to revolutionise local and global health care and people's lives (Snoswell et al., 2020).

The expanded use of “telehealth” during COVID-19 has shown critical limitations in the capacity of health providers to regularly provide “telehealth”. Inadequate “telehealth” and system interoperability, knowledge and capacity shortages in the present health staff, unsustainable funding structures, and difficulties with data sharing are a few of them. To decrease these problems and allow HC providers to properly develop their services, solutions are needed. Additionally, guidelines on how “telehealth” should be supported and regulated in the future are needed for governments and other funding organisations (Zhou et al., 2020).

Despite the potential of tele-health to offer convenience and time saving to the patient, some of the operational and technical challenges remain unresolved and limit its optimum use. Lack of consistent user experience can be attributed to such issues as inconsistent connectivity and usability problems on the platform, inability to conduct physical examination, uncertainty about how to diagnose complex conditions, and irregular user experience. Such issues cast doubts on the reliability, safety, and clinical adequacy of tele-health as a primary healthcare modality.

Moreover, the swift growth of tele-health has not been accompanied by the corresponding increase in digital literacy, system interoperability, and workflow alignment between conventional and virtual care. Consequently, the users can experience challenges with navigation through the tele-health platforms, interpretation of remote directions, or developing faith in online consultation. Such hindrances are especially significant in environments where tele-health is adopted due to the need and not the willingness.

Meanwhile, tele-health is steadily becoming accepted by patients who appreciate its assumed benefits. In any case, it is not clear how much these benefits will supersede the real challenges. Empirical data about the ways in which users assess tele-health regarding convenience, satisfaction, perceived quality of care, and the desire to continue using this

type of service is lacking. Besides, the extent of benefit and limitation of tele-health adoption differs among groups of users, and it is influenced by their previous experience, trust in any digital technology, medical conditions, and the use setting.

Thus, an obvious research problem arises: although there is an increasing number of tele-health solutions available and promising, it has not been studied how users perceive their benefits, encounter their challenges, and create their evaluations regarding their overall viability and further utilisation. This gap should be addressed in an attempt to come up with tele-health systems that are clinically reliable, user-friendly, and sustainable in everyday healthcare delivery.

1.3 Purpose of Research

This study aims to assess the viability and perceived feasibility of telehealth services through assessing perceived benefits, perceived challenges, satisfaction level, perceived quality of care relative to face-to-face services, and persistence of use of telehealth. It presents a theoretical framework that tackles the challenges to integrating and standardising the use of “telemedicine” in HC facilities following a literature assessment that highlights significant theories, models, and frameworks utilised in the science of execution.

Objectives:

- To identify the advantages of “telehealth”, examining its impact on healthcare affordability, quality improvement, and accessibility for diverse populations.
- To explore the drawbacks and constraints associated with “telehealth”, such as funding and regulatory challenges, technical complexities, and patient acceptance issues.
- To assess the association between prior telehealth use and willingness to use telehealth again.

- To compare perceived telehealth care quality with in-person care and examine its statistical relationship with overall user satisfaction.
- To assess the future prospects and challenges for the growth of telehealth, evaluating its implications for healthcare systems, providers, and patients.

This research aims primarily to steer the research process, to perform a comprehensive analysis of “telehealth”'s most notable characteristics, and to generate knowledge that is usable in the implementation of beneficial and necessary changes in HC delivery.

1.4 Significance of the Study

The primary reason for the importance of this study is its ability to obtain a comprehensive insight about the “telehealth” solutions and their potential application in different types of HC settings, not only in the U.S.A but also globally. “telehealth” has turned out to be exceptionally beneficial for the health services sector by eliminating the distance barrier between patients and doctors and allowing the latter to provide their services without having to meet in person which thus creates a very good chance for people especially the marginalized, the poor, or those living in remote areas to access health services. With its systematic review of the pros and cons along with the “telehealth” use, this research will be an important source of information for finding out how digital health technology can improve efficiency, quality, and affordability of medical services. The research examines factors such as organisational, financial, regulatory, and technological ones influencing successful “telehealth” adoption, among others, to help the stakeholders in identifying the opportunities and predicting the possible difficulties. In this way, it backs the Evidence-based Decision Making and broadens the literature on digital HC innovations while supporting the future research and initiatives about “telehealth” integration into the regular HC process that would be able to reap the study's benefits.

In addition, the results of the study are of practical significance to a range of stakeholders such as HC providers, policymakers, technology developers, and patients. The HC system could benefit from the research output by devising strategic models that would contribute to the effective utilisation of “telehealth”, patient involvement, and the continuous provision of services. The evidence generated from the study will also be a strong argument for the lawmakers to pass the necessary legal, financial, and infrastructural measures to guarantee equitable access to “telehealth” services based on scientific data. Furthermore, the study identifies and explicitly mentions the obstacles, which are, among other things, patient digital literacy, regulatory restrictions, funding problems, and technical challenges, and indicates which parts would require targeted actions in order to make “telehealth” the most efficient method. The information gained is, therefore, going to be an important source of knowledge for all the global and local initiatives that are going to be set up for the purpose of extending “telehealth” utilization, enhancing accessibility and quality of HC, introducing digital health delivery innovations, and at the same time, ensuring a HC system that is both resilient and able to cope with both ordinary health services and public health emergencies.

1.5 Research Purpose and Questions

The importance of this study is that it offers an empirical review of the feasibility of telehealth on the user side, not through policy discourse or market growth rates alone. The study improves the more organised knowledge on telehealth sustainability in real-life health care environments by analysing perceived benefits, challenges, satisfaction, perceived quality, and reuse intention.

Conceptually, the study will build on the current technology adoption literature by incorporating variables of feasibility assessment and user perception, which enhances the

conceptual connections between the acceptance models and the science of healthcare implementation.

Practically, the results will be useful in helping healthcare providers to determine obstacles that restrict the use of telehealth and to model user-centred methods of improvement. Knowledge of the effect of perceived quality and satisfaction on further use can be used in service redesign, training programs, and digital infrastructure planning.

In terms of policy, the research could offer evidence that could inform regulatory policies, the distribution of funding, and the approach to investing in digital health with the goal of achieving equitable, reliable, and sustainable telehealth systems. The research will also include a literature review, trend definition, and evaluation of the opportunities and challenges to generate evidence-based research that is useful for policymakers, HC providers, and even tech developers in the process of developing, implementing, and scaling effective “telehealth” solutions. Furthermore, the entire research is focused on the “telehealth” discussion from the perspective of access, quality, and cost, with the aim of providing roads for its successful implementation and effective usage in different HC settings.

Research Question

- **RQ1:** How does telehealth contribute to healthcare affordability, quality improvement, and accessibility among diverse population groups?
- **RQ2:** What are the major challenges and limitations affecting the adoption and effective implementation of telehealth, including funding constraints, regulatory issues, technological barriers, and patient acceptance?
- **RQ3:** What is the relationship between users’ prior experience with telehealth services and their willingness to use telehealth in the future?

- **RQ4:** How do users perceive the quality of telehealth services compared to traditional in-person healthcare, and what impact does this perception have on their overall satisfaction?
- **RQ5:** What are the future prospects and challenges for the growth of telehealth, and how might these developments impact healthcare systems, providers, and patients?

1.6 Business & Strategic Context of Telehealth

Telehealth adoption has continued to transition to being perceived as a clinical innovation, rather than as a strategic business investment in healthcare organisations. Although the initial discourse of telehealth focused on enhancing access to care and clinical efficiency, in modern healthcare, telehealth programs must be assessed from the perspectives of organisational strategy, financial sustainability, and value creation in the long term. Telehealth is no longer a technological option but a strategic choice that entails significant capital and operating restructuring and risk management for healthcare executives and decision-makers.

The environment in which healthcare organisation's function has been typified by increased costs, competition, complexity in regulations, and increased patient expectations. In this situation, telehealth is an opportunity, as well as a strategic challenge. The adoption of telehealth should be rational not only in service delivery improvements but also on the basis of a return on investment (ROI), scalability, and institutional goals. In turn, it is important to interpret telehealth as a business and strategic choice in order to take informed managerial action.

Telehealth as a Strategic Decision to Invest

Strategically, the implementation of telehealth is a capital investment choice that entails the use of financial, human, and organisational capital. The investments in telehealth

are usually accompanied by the need to spend money on digital platforms, licensing, integration of systems, personnel training, and maintenance. In contrast to the conventional infrastructure healthcare investments, telehealth projects have intangible resources, in terms of digital capabilities, organisational learning, and service innovation, that are difficult to quantify and assess.

The healthcare executives must thus evaluate the telehealth initiatives based on decision criteria that go beyond the immediate cost reduction. In this respect, strategic investment appraisal means determining whether telehealth can lead to competitive advantage, service differentiation, retention of patients, and market growth. In most scenarios, telehealth investment is made under the uncertainty of demand trends, reimbursement models, and regulatory stability, which makes it difficult in traditional forms of capital budgeting (Drummond et al., 2015).

Telehealth investment is also long-term in nature, which enhances the strategic aspect of telehealth. Telehealth systems usually transform the organisational processes, models of patient interaction, and the priorities of resource allocation once implemented. Accordingly, the telehealth adoption has path-dependent consequences, and initial strategic decisions are important in long-term organisational performance.

Uncertainty in the market and Strategy risk.

Market uncertainty and strategic risk are two of the most significant issues in telehealth adoption. Several external drivers, such as patient acceptance, regulatory policies, reimbursement structures, and competitive dynamics, affect demand for telehealth services. They are not very stable and differ in regions and healthcare systems, and therefore, managers are not able to predict the level of utilisation and financial returns (Sarrami-Foroushani et al., 2014).

The strategic risks associated with telehealth adoption are the fact that investments can fail to achieve the expected value because of low user adoption, policy reversals, or operational inefficiencies. With their high reliance on behaviour and perception, unlike traditional healthcare services, the uptake of telehealth presents behavioural dialecticism in the strategic planning process. The healthcare managers, therefore, need to keep in mind that technological feasibility is not the only source of strategic risk, but also user acceptance and continued engagement.

In addition, this uncertainty is exacerbated by competitive pressures. With the increased commoditization of telehealth, healthcare organisations need to be aware of the commoditization risks in the digital services, where services have limited differentiation and margins are compressed. Strategic positioning thus becomes very vital in determining whether telehealth improves organisational performance or raises the level of complication in its operations without corresponding returns.

Telehealth Implementation Cost Structures.

Realising the cost structure of telehealth is necessary to make sound strategic decisions. Costs associated with telehealth may be grouped into fixed costs, variable costs, and indirect costs. The fixed costs usually constitute the platform acquisition or development, licensing, the provision of cybersecurity, and system integration costs. These expenses are made regardless of the volume of service and hence form a major entry barrier to the smaller healthcare entities.

Variable costs are related to service delivery and comprise staffing expenses, consultation time, technical support, and incremental infrastructure use. The telehealth approach changes the dynamics of costs in contrast to in-person services, as it allows setting more flexibility in staffing and the use of resources. These efficiencies are, however,

dependent on having sufficient scale, without which variable costs can supersede benefits (Whitten & Love, 2005).

Telehealth planning often underestimates indirect costs like training, change management, and adaptation of the organisation. Staff resistance towards change, alterations of workflow, and cultural adaptation add extra costs that cannot be readily reflected by financial statements but which greatly influence the success of the project. Strategically, the costs may not be taken into consideration, which will result in over-optimistic ROI estimates and future managerial disappointment.

Telehealth Revenue Models.

Revenue models that would affect financial sustainability also need to be carefully considered when adopting telehealth. Telehealth revenue models are common, and these are subscription-based services, fee-based services, and bundled care services. The models are characterised by different strategic implications and risk profiles.

Models that are subscription-based offer a steady revenue flow and are commonly used in cases of chronic care management and patient engagement. They must be able to sustain a consistent number of patients and value, though they must feel the value to warrant a recurring payment. Instead, fee-for-service models are more consistent with traditional healthcare reimbursement models and face the risk of being sensitive to volume changes and regulatory limitations (Dorsey & Topol, 2016).

Bundled care models combine telehealth with other types of care packages, which focus on value-based achievements and not on the service transactions. Although a reasonably appealing strategy, bundled models require advanced coordination and outcome measurement applications, which can be a heavy burden on managerial and analytical resources. The choice of a suitable revenue model is thus a tactical move that should consider the organisational potential, market factors, and regulatory measures.

The telehealth adoption has managerial challenges.

Regardless of the strategic value of telehealth, healthcare managers continue to face challenges in returns on investment, sustainable scaling of services, and aligning telehealth programs with business strategy. Many telehealth benefits turn out to be difficult to justify using ROI because of their long-term and intangible nature, including patient satisfaction, brand reputation, and company education.

Another important issue is the scaling of telehealth on a sustainable basis. Although pilot projects might prove operationally viable, scaling will necessitate effective governance frameworks, standardised processes, and uniform service quality. Lack of strategic alignment means that telehealth initiatives will still be seen as fragmented add-ons instead of part and parcel of organisational strategy (Greenhalgh et al., 2017).

Corporate strategy fit is essential to prevent strategic drift. Either opportunistic or reactive telehealth programs might not support the organisation's overall objectives, such as market positioning, service differentiation, or financial sustainability. Strategic coherence is necessary since investments in telehealth support do not water down organisational priorities.

Placing the Present Study.

The present study fits within this context as a decision-support model of healthcare leaders as opposed to a technical or clinical analysis of telehealth systems. The study offers a reflection on the idea of behavioural factors that can have a direct impact on the adoption of telehealth and, consequently, the attainment of the strategic and financial value, given that the research concentrates on the perceptions of usefulness and ease of use as the factors in the perceptions of users.

By knowing these behavioural determinants, managers are then able to make better decisions as to investment priorities, system design decisions, and implementation

strategies. Instead of focusing on the technological features, the study will be useful in the strategic decision-making process by determining the circumstances in which the investments in telehealth are more prone to succeed in business terms.

By so doing, the research will fill the gap between the technology adoption theory and strategic healthcare management. It provides empirical evidence on the managerial choices involving resource allocation, risk mitigation, and long-term planning and raises the strategic value of telehealth adoption in healthcare organisations.

CHAPTER II: REVIEW OF LITERATURE

2.1 Introduction

One more technical enhancement in the present-day HC system is “telehealth”, which, thanks to the merger of digital technology, can smoothly get rid of the barriers related to access, cost, and location. The concept of “telehealth” is already established due to the rapid progress in communication technology, "Artificial Intelligence" (AI), and mobile health apps; in fact, they are now indispensable components of the health systems globally. The chapter provides a critical perspective on the definition, concepts, and theoretical foundations of “telehealth” and functions as a concluding summary of the literature on the topic. The chapter brings forth the history of prior research, aligning it into a single entity in order to create a theoretical and applied framework for evaluating “telehealth” solutions.

The assessment also discusses acceptance and use patterns, the technologies currently being used to speed up the “telehealth” industry, and the national and worldwide levels. It includes a critical assessment of the benefits of “telehealth” in terms of enhancing accessibility, affordability, and quality, as well as the problems, which include acceptance, technical, and regulatory challenges. In doing so, the chapter not only summarises the body of information already in existence but also identifies research gaps that will be necessary for future assessments of the potential of “telehealth” solutions. This systematic review would not only suit the objective of the work, but also furnish the foundation of evidence-based suggestions that would be used to enhance the “telehealth” implementation and adoption at the international level.

2.2 Theoretical Framework

Technology Acceptance Model (TAM)

This model was created based on the Theory of Reasoned Action (TRA), and a particular model was created to describe how people develop their attitudes and intentions to use information systems. TAM also focuses on the way the cognitive assessments of users influence the decisions made towards using technology. It is especially useful when the technological systems are implemented to enhance efficiency, effectiveness, and service provision. TAM has been a fundamental model in analysing technology adoption practices in organisational and personal settings due to its flexibility and descriptive capability.

The perceived Usefulness shows how much the user believes that the utilization of a specific system will result in better performance, or will bring about better results, whereas the Perceived Ease of Use shows how much the user thinks that using the system will be effort- and complexity-free. The beliefs are known to alter the attitude of the users of the technology, and this attribute affects their desire and intention to use the technology. Regarding telehealth, PU can be represented by the perceived enhancement of healthcare access, efficiency, and quality of care, whereas PEOU is associated with the ease, functionality, and design of telehealth systems. These constructs collectively offer an objective prism through which telehealth technology acceptance and resistance by users may be considered, thus rendering TAM an appropriate and powerful framework to examine the potential of telehealth adoption in modern healthcare systems.




Figure 2.1: The “technology acceptance model”

Source: (Davis, 1989)

The Technology Acceptance Model (TAM) describes how users arrive at the acceptance and utilisation of a new technology by demonstrating the associations among the important psychological and behavioural variables. The model demonstrates that there are external factors like system design, user training, and organisational support that affect the Perceived Usefulness (PU) and Perceived Ease of Use (PEOU) of users. Perceived Usefulness describes how much the users feel the technology would improve their performance or results, whereas Perceived Ease of Use describes how easy the system is viewed to be. The model also suggests that more accessible technologies are the ones that are more perceived to be helpful. Both PU and PEOU influence the attitude of the users concerning the use of the technology, an element that determines the intention of users to use the technology. One of the determinants of the actual system use is behavioural intention, which is the actual adoption and sustained use of the technology. Applying this model to the context of telehealth, this model allows clarifying the effects of the perception of usability and effectiveness on the desire of users to adopt and use digital healthcare services regularly.

Core Constructs of the Technology Acceptance Model

Perceived Usefulness (PU)

Perceived Usefulness can be defined as the level at which one feels that adoption of a given system will support his or her job performance or effectiveness (Davis, 1989). That is, a user will become more inclined to use a technology when they see obvious advantages in it, i.e., increased efficiency, better results, time, or improved service quality. A study conducted by (Holden & Karsh, 2010) indicated that healthcare professionals tend to adopt the health information technologies when they are convinced that the systems would have beneficial effects on the clinical performance and patient outcomes. Equally, (Chau & Hu, 2002) discovered that the uptake of the telemedicine technologies by the physicians was mainly influenced by the perceived utility of the technology and not necessarily due to pressure or organisational requirements.

Perceived Ease of Use (PEOU)

Perceived Ease of Use is the degree to which an individual holds the belief that s/he will not be required to undertake hectic efforts in an effort to utilise a specific technology (Davis, 1989). The lower acceptance of technologies that seem complicated, time-consuming, or hard to use is observed, even in the case of their potential gains. Ease of use in telehealth applications involves simple navigation, intuitive system design, limited technical complexity, reliable connection, and user-friendly interfaces to both the patients and the healthcare providers. PEOU is of particular importance to populations that lack digital literacy, such as elderly patients or users in rural areas, where adoption willingness depends particularly on its effects. The literature is always in favour of the contribution of PEOU in technology acceptance in healthcare. Research by Venkatesh & Bala (2008) indicates that ease of use not only has a direct relationship with behavioural intention, but also an indirect connection with the perceptions of usefulness since systems that are easier

to use are perceived to be more useful. With telehealth, when users perceive a digital consultation platform as easy to use, they are more likely to appreciate their functionality benefits and maintain their use.

TAM and Behavioural Intention to Use Telehealth

TAM indicates that both perceived usefulness and ease of use affect the attitude of the users towards technology, which in turn affects their behavioural intention to use and actual adoption behaviour. Behavioural intention is a notion that is used in telehealth adoption as it determines how people and medical staff would like to participate in remote consultations, digital diagnostics, and other systems of healthcare delivery. Other studies have used TAM to investigate the acceptance of telehealth among the various stakeholders. Indicatively, P. J. Hu et al. (1999) used TAM to examine the acceptance of telemedicine technologies by the physicians and discovered that PU was significant on intention to use compared to PEOU, which underscored the performance-oriented aspect of clinical professionals. On the other hand, the ease of use is frequently emphasised as no less significant in patient-focused studies, especially when dealing with non-technical users (Zhao et al., 2018). These results indicate that effective telehealth implementation should be based on the concept of balancing between the functional value and the usability of the systems so that they can be both efficient, convenient and helpful.

Extensions and Applications of TAM in Healthcare

TAM has been expanded over time and modified to more effectively reflect the complexity of a healthcare setting. Other variables that researchers have included in their effort to augment the explanatory power of TAM in health technology adoption scenarios include trust, perceived risk, social influence, facilitating conditions, and privacy concerns. Trust in digital platforms and data security has been found to be a key issue in telehealth research in relation to perceived usefulness and intention to use (Asua et al., 2012). The

issue of privacy and confidentiality is especially relevant to healthcare since the information about patients is rather sensitive. Incorporation of TAM with trust and perceived risk constructs has proven to predict telehealth adoption behaviours better. Nevertheless, the original TAM is the theoretical framework to learn about telehealth acceptance based on its conciseness, empirical evidence, and flexibility. The constructs in its core are still helpful to understand why users accept or reject digital healthcare technologies.

Relevance of TAM to the Present Study

The Technology Acceptance Model (TAM) is a suitable and effective theoretical framework that can be used to analyse the potential of adopting telehealth solutions by users in the framework of the current study. Through the two fundamental constructs of Perceived Usefulness (PU) and Perceived Ease of Use (PEOU), the study aims to comprehend the cognitive process of utilising telehealth services by the users based on their functional value, convenience level, and general ease of use. These assessments are especially essential in telehealth, as the readiness of the end-users to substitute or complement conventional face-to-face scheduled appointments with online ones is largely conditioned by their attitude towards the effectiveness, efficiency, and comfort of interaction. The anticipated results are that increasing perceived usefulness and ease of use will have a positive impact on the behavioural intention of the users to adopt telehealth solutions, which will consequently increase the potential of continued usage and acceptance in the long-term.

Applying TAM to the process of studying telehealth adoption, this study is in line with previous research in the field of technology acceptance in healthcare and is relevant to the increasing literature on the topic of digital transformation of healthcare delivery systems. TAM offers a systematic and empirically verified method of determining the

willingness of users towards the deployment of telehealth technologies, determining the main psychological and technological obstacles to its adoption, and comprehending the factors that facilitate or could impede user adoption. Moreover, the use of TAM in the present study provides useful information to the healthcare community, policymakers, and technology developers by identifying areas that require improvement in the design of the system, training strategies, and implementation strategies. Therefore, by providing a stronger theoretical base to the current research, TAM also increases the usefulness of the research in practical use, educating evidence-based interventions designed to increase the uptake and use of telehealth.

Diffusion of Innovation Theory (Everett Rogers, 1962)

A theoretical framework used to clarify and predict the course and determining aspects of a sequence of "innovations," such as new ideas, technologies, or products in society, is known as the "diffusion of innovations theory" (DOI). Innovative diffusion theory was first studied by American sociologist Everett M (Rogers, 2003). Rogers's remark regarding the spread of specific agricultural advances. He noticed that many years after they first appeared, many new techniques and technologies that were extremely beneficial to local farmers were progressively accepted and used in Carroll, Iowa, the United States, where he was born. Rogers participated in a hybrid maize dissemination study and spoke with 148 farmers regarding the utilisation of cutting-edge items, including herbicides (Davis, 1989). After gathering research on the spread of innovation across several domains, covering kindergartens, driving schools, and medical circles, it was discovered that while industries and fields vary, the results of innovation diffusion throughout various domains are comparable. For instance, the innovation adoption process exhibits an "S" curve; the pioneers of innovation adoption (the first to accept innovation) are typically knowledgeable and experienced individuals, and there are variations in the

information sources and channels at different stages of the decision-making process. Rogers made his proposal for innovation in the early 1950s, attempting to explain how innovation spreads across time and social systems, for the first time in his seminal book *The Diffusion of Innovation* (1962). This proposal was based on long-term observation and investigation research on new technologies (particularly in the agricultural field) and thought communication in society, as well as the compilation and analysis of related diffusion research across multiple fields and reference from multidisciplinary research results from sociology, communication, economics, and pedagogy. A universal invention diffusion model is developed in the interim.

Unified Theory of Acceptance and Use of Technology (UTAUT)

Nowadays, the majority of nations understand how crucial electronic government, or e-government, is to contemporary public administration. A number of state services can be accessed through Turkey's outstanding e-government platform, which has been in use since 2008. To various individuals, the term "electronic government" may have varying connotations. In a broad sense, it describes the efficiency with which the government makes use of information and communication technologies to deliver electronic services to individuals, companies, and public organisations (Nam, 2014).

The provision of public services to citizens through the use of information and communication technology is a crucial function of e-government (Berlilana et al., 2017). In Turkey, the process of transferring documents from state institutions to electronic media started in 2013. To satisfy business needs, electronic document management systems (EDMS) allow studies to be digitised through computer systems and technology. This makes it simple for many organisations that use an all-inclusive EDMS to handle all information generated both internally and externally. As a result, this system continues to operate at a higher level than institutions that use conventional approaches in terms of

productivity and effectiveness in service provision. Every institution must successfully deploy EDMS since these technologies expedite business processes and offer users ease. There are numerous difficulties when replacing an outdated system or technology with a new one. According to certain studies, the new technology or system is not adopted or embraced (Davis, 1989; Norzaidi et al., 2008; Song et al., 2015). The Ministry of Development of the Republic of Turkey's Izmir Katip Çelebi University supports “University Information Management System” (UIMS), which has modules that can be combined with the software used by Bartın University's electronic document management system. UIMS is an e-university project consisting of integrated modules covering all facets of academics and university administration. Because of the project's nature, this system is made to work in both public and private universities. It includes an archive and document management system that is safe, dependable, and appropriate for the institution's goals and structure.




Figure 2.2: The “unified technology acceptance and use of technology theory” (UTAUT)
Source: (Venkatesh et al., 2003)

2.3 Conceptual Framework and Definitions of “Telehealth”

A combination of "telemedicine," "e-health," and "m-health," the phrase "telehealth" describes the use of ICTs to provide medical treatment to patients in a distant location. Although “telemedicine” has long been associated with distant clinical services,

the term has recently expanded to cover a wider range of applications in the healthcare industry, such as wellness programs, illness prevention, remote monitoring, and even administrative tasks. Establishing a clear conceptual framework is essential to differentiate these overlapping terms and to understand the theoretical foundations that guide their adoption and implementation. The feasibility analysis will mean that the combination of technology and HC delivery will be characterised in the specification of “telehealth” to such an extent that the researchers and decision-makers will evaluate its importance as a mere tech breakthrough and also as a complete HC model that introduces access, quality, and efficiency into new health systems.

Okesiji et al. (2024) use a strong mixed-methods study design, which incorporates both qualitative and quantitative measures of performance in understanding the intricate connection between digital health operation efficiency, technological innovation, and economic performance. Even though the methodological triangulation underlies the internal validity of the results, the study mostly presupposes that the financial performance improvements in the digitally mature healthcare environments can be generalised to the various health systems. The assumption can also provide limited generalizability of the findings, especially in resource-restricted or digitally disparate situations when the level of infrastructural preparedness and user capabilities differ greatly. Even though the writers have made a solid case that the benefits in the long run (i.e., better patient satisfaction, lower operational expenses, and lower readmission rates) outweigh the investment costs, the research studies fail to adequately explain the transition risks, difficulty in workforce adaptation, and the financial strain incurred in the short run during the early stages of the implementation. The sustainability of the economy claims, therefore, although a positive outlook, is context-based and should be further empirically tested in the settings of heterogeneous healthcare. The literature review in Okesiji et al. (2024) successfully follows

the development of remote care, from the simplest form of teleconsultation to sophisticated, multi-purpose telehealth tools, and the use of telemedicine is connected to the overall economic theories of efficiency and value creation. Nonetheless, although it singles out financial sustainability as the key issue, the research mainly addresses the concept of return-on-investment (ROI) and market expansion, which provide little information on how regulatory inflexibility, reimbursement lag, and governance restrictions can erode estimated economic returns. The focus on formation of partnerships and scalable business models underscores key strategic facilitators, but these prescriptions are represented in much of the organisational profitability prism, and little is done to focus on equity-based outcomes or resiliency of the public health system. This leaves a void that is filled by the current research by assessing the feasibility of telehealth based on financial indicators and sustainability, access, and policy-based perspectives.

The authors Olivencia & Sasangohar (2023) change the focus to the issue of the system level of integration, stating that the productivity of telehealth is not complete without the analysis of such sociotechnical variables as are embedded in the clinical workflows. Their study offers an important critique of technology-focused adoption models because they show how telehealth use alters provider practice patterns, thus affecting efficiency, quality of care, and patient safety. However, although the study is right in its identification of the lack of a well-developed theoretical framework, in which telehealth integration has to be considered, its conceptual focus restricts the degree to which empirical performance outcomes can be contrasted in different settings. Consequently, the applicability of the framework has not been sufficiently tested in a large-scale or longitudinal healthcare setting and is supported to date by three case studies of telehealth, which lead to a sociotechnical framework based on the principles of systems engineering (Olivencia & Sasangohar, 2023). In spite of the fact that this interdisciplinary

approach has the advantage of improving conceptual rigour, the small sample size of case studies denies the model the ability to explain variation, especially when it comes to explaining the difference in healthcare systems with different governance structures and funding systems. This weakness underlines the necessity to conduct empirical research that should combine sociotechnical models with quantifiable outcomes in terms of cost efficiency, patient satisfaction, and service accessibility, which is the trend followed in the present study. On the same note, Kobeissi & Hickey (2023) elaborate on a telehealth model that is based on the systems theory guided by the Donabedian structure-process-outcome framework. Their model provides an all-encompassing analytical prism through the examination of the interplay between policy requirements, organisational attributes, professional skills, as well as patient-level factors of health. Although this multidimensional approach is a major strength, the research is largely normative as it provides insights into how the telehealth systems are supposed to work, but does not empirically show how such interactions affect the performance of the operations or financial aspects. Moreover, the model fails to pay enough attention to the impact that the differences in digital literacy and access to infrastructures can undermine the equity goals it aims to advance.

This study, Dhabarde et al. (2023), investigates the value of “telehealth” and the various tools used to integrate artificial intelligence into the HMS. “Telehealth” refers to the method of remotely assessing and caring for patients in the event of a pandemic. Some of the key benefits of “telehealth” are shorter wait times, lower travel expenses, and better treatment quality. In remote or underserved locations, where medical treatment is scarce, it might be quite useful. There is a pressing need for better care of chronic diseases through the use of “telehealth”, particularly in emergencies. Readers learn about approaches that can help people cope with the demands of HC while also self-managing their chronic

diseases; this is crucial since doctors were not available to treat patients throughout the outbreak. Directly and indirectly, “telemedicine” helps reduce the spread of viruses by maintaining a safe distance between patients and HC providers. “Telemedicine”, especially for diagnostic and therapeutic purposes over remote distances, has proven to be very advantageous in the case of COVID-19, diabetes, mental disorders, and cardiovascular diseases (CVDs). Therefore, if they want to move along with the current trends, then the HC sector and hospitals have to think out of the box and come up with new ways to digitally transform their practices.

This study, Carroll (2023) It investigates how innovations in “telehealth” are revolutionising the accessibility and delivery of HC. The HC landscape has changed since the COVID-19 pandemic began, giving patients and doctors more options. For those who are unable to attend in-person visits, “telehealth” improves accessibility by facilitating telephone and virtual consultations. The rates of missed visits and appointment rescheduling have decreased as a result of this change in patient adherence to planned care. “Telehealth” has many advantages for both patients and HC providers, but there are still many obstacles to overcome before it can be considered a regular practice. This study examines current innovations and empirical evidence to make that case. As a result of reading this, stakeholders and HC providers should have a better grasp of Telehealth’s benefits and drawbacks, allowing them to make educated decisions about the method’s potential future uses.

During the Zhang et al. (2023) COVID-19 With strong backing from numerous federal and state agencies, “telehealth” programs significantly grew throughout the epidemic and public health emergency. Innovative health service delivery models that rely on factors other than financial support have their future sustainability cast into doubt due to the haziness of upcoming “telehealth” payment standards. In order to assess clinical

effectiveness and financial viability, they have created a system for collecting relevant data at different phases of program execution. Due to the lack of studies focusing on the development of long-term sustainable "telehealth" programs, this paradigm is applicable in many contexts and gives health equity great importance. They acknowledge the difficulties of sustaining "telehealth" programs with a fee-for-service model alone and urge all parties involved, especially payers and lawmakers, to consider the programs' cost-effectiveness and back alternative payment models to make them last.

“Telehealth” is Vystropova & Bass (2024) a contemporary area of medicine that offers excellent, reasonably priced HC. Just as crucial as in-person bedside contact is the doctor-patient exchange that takes place during an online session. A qualified professional who can assist patients remotely via a computer and telecommunication software is known as a “telehealth” doctor. For patients who live in rural or isolated locations with little access to HC, “telehealth” is especially crucial. The essay explains the importance of communication in productive “telehealth” interactions. The authors looked at several online “telehealth” services, including the Magical Doctor multidisciplinary medical clinic and the Gostelemed “telehealth” discussion service offered by state medical organisations, as well as the mobile medical platform Capsa HC. The authors examined the fundamentals of successful “telehealth” communication using the techniques of data analysis, generalisation, and classification, in addition to a terminological analysis of dictionary definitions. During online sessions, they determined the tools and resources for efficient verbal and nonverbal communication. Particularly in times of crisis or epidemic, “telehealth” has been shown to be a useful instrument for raising the caliber and availability of medical care. Positive relationships between patients and specialists are fostered by effective “telehealth” communication.

In Sahoo et al. (2023) A systematic study of the intellectual architecture, as well as the trends and patterns of publishing research studies in telehealth, is conducted by a complex of bibliometric review methods based on the SPAR-4-SLR protocol and content analysis, performance analysis, and clustering of keywords and co-occurring patterns. Although the methodological soundness and the depth of the review give essential insights into the development of the telehealth scholarship in journals, institutions, and geographical locations on the macro-level, the literature is rather descriptive in character. The ten major research themes identified, including e-health service guidelines and telemonitoring of critical diseases to change management and health information technology, provide a convenient thematic categorisation; nevertheless, they lack in the assessment of the relative maturity, empirical strength, and practical efficacy of the mentioned themes. Consequently, the review summarises what has been researched instead of critically evaluating the level of support of existing evidence for the actual practice of telehealth, especially in under-resourced healthcare systems. Moreover, even though Sahoo et al. (2023) go further to address the new fields of inquiry, like the Internet of Things (IoT) in health and the relative paradoxes of virtual and in-person care, these arguments are not empirical and are conceptually disaggregated. The commercial nature of the framing of the review is both strategically valuable and creates a bias of innovation diffusion and productivity stories in which little focus is placed on patient-level outcomes, equity, or operational viability in the context of public healthcare. Therefore, the study does not go any further to make practical implications for future research directions; therefore, it is not true that it lacks any success in identifying the future research directions. The current research addresses this weakness by empirically analysing the feasibility and effectiveness of telehealth outside of patterns of publication, namely perceptions of users, quality of services, and sustained outcomes.

Rintyarna et al. (2023) Concentrate on the opportunities of AI-based telehealth systems to enhance access to healthcare in rural settings and use a descriptive qualitative research methodology, which assumes thematic analysis of the available literature. The inclusiveness and access in the study are one of the main gaps that need to be filled in telehealth research, especially in underserved populations that are geographically isolated. Nevertheless, the use of secondary data and conceptual synthesis restricts the capacity of the study to prove its arguments about enhanced health results and cost reduction. Although the authors rightly point out regulatory constraints, security of data, lack of sufficient infrastructures, and digital literacy among key barriers to the success of big data initiatives, these issues are presented separately without connecting them using an operational or economic framework. In addition, the study presupposes that telehealth solutions developed by AI can be successfully implemented in rural settings without sufficiently considering the preparedness of medical facilities or the socio-cultural adoption of new technologies. This brings concerns about the scalability, as well as sustainability, of AI-based telehealth models in low-resource scenarios. The current study has overcome these weaknesses by empirically measuring the adoption of telehealth in different populations of users and hence providing a more realistic assessment of the viability beyond what is theoretically possible.

The study by Butt et al. (2024) is a contribution to the literature as it provides insight into mobile health (mHealth) monitoring as one of the fast-moving fields in remote health care provision. Their study emphasises how mobile applications have become an increasingly important part of the process that makes it possible to monitor patients, conduct teleconsultation, respond to emergencies, and manage chronic diseases. Although the study is effective in highlighting the technological developments in mHealth systems, it also reveals one of the biggest gaps in the current research, that is, the peripheral

consideration of telemedicine-related implementation issues in mHealth systems. Most of the reviewed studies do not discuss the technical aspects of a system, including clinical decision-making processes, interactions between patients and physicians, and care coordination, and therefore offer high-level system overviews. Also, the fact that the studies on patient rescue procedures and social differences in the health care settings have not been broadly developed limits the use of mHealth solutions in emergency and high-risk conditions. Despite the fact that Butt et al. (2024) recognise these gaps, their research is rather theoretical, providing small empirical data to substantiate the arguments on system effectiveness and safety. The present study is based on this criticism and critically examines the benefits and limitations of telemedicine-enabled mHealth systems in terms of feasibility and user experience, thus providing empirically-based knowledge in the field, which is yet to be thoroughly investigated.

In conclusion, a well-defined conceptual framework and accurate “telehealth” definitions are two aspects of the same tool that assist both scholarly research and practical implementation. The researchers and practitioners will only be able to do the latter if they clarify the differences between the terms “telemedicine”, e-health, and m-health. This proactive and systematic approach of outlining the concepts by researchers will help not only to clarify but to characterise “telehealth” as a complex phenomenon. Moreover, it will facilitate debate on its adoption, benefits, challenges, and future directions based on the agreed-upon understanding of the concept and the ways it integrates with the overall goals of modern HC transformation.

2.4 Global and National Trends in “telehealth” Adoption

The global use of “telehealth” has seen a remarkable rise owing to the new technological advancements, the growing medical demands, and the elimination of location and infrastructure barriers. “telehealth” is gradually being incorporated into HC systems of

different countries worldwide as the countries are making efforts to ensure “telehealth” access that is more accessible, less costly, and more efficient in-service delivery. In most developed countries, the digital infrastructure and favourable policies are conducive to “telehealth” adoption, whereas in developing countries, the mobile penetration and new solutions based on low costs determine the growth. The deployment of “telehealth” at the national level showcases the differences in socio-economic status, policy framework, and HC priorities, and therefore, it is necessary to study the global trends and the experiences of individual countries. Understanding such trends can be very useful, as they can highlight the conditions under which “telehealth” solutions turned out to be effective or ineffective in different settings.

Nonda et al. (2025) note that sustainable growth in telemedicine can be achieved by strategic partnerships, scalable business models, and active financial planning, which market-based strategies are presented as the mechanisms of providing equal access to disadvantaged populations and increasing revenues. Although such a dual emphasis on inclusivity and profitability adds to the strategic significance of the study, the analysis is premised to a large extent on the compatibility between commercial growth and equitable healthcare provision. This supposition ignores structural inequalities, including digital access inequalities and geographical regulatory fragmentation, that can frustrate the efficacy of market-based strategies in reaching the marginalised communities. In addition, as much as the research points out the necessity of regulatory changes and adaptive reimbursement systems, it fails to empirically investigate the impact of current policy limitations on the slow pace of the introduction of telemedicine into state governments. The study on the assessment of ROI, the development of the market, and corporate development strategies in Nonda et al. (2025) helps to take a more comprehensive view of telemedicine financing, but the analysis is firm-focused. This study provides minimal

information about patient-level affordability, service sustainability, and resilience of systems in the long term due to its exclusive focus on organisational returns and scalability. As a result, the study can be of valuable strategic guidance to telemedicine providers; however, its generalizability to a public or hybrid healthcare system is limited. The current study responds to this limitation by considering the economic viability analysis with the consideration of the service quality, access, and sustainability.

Hong (2024) analyses the telehealth technologies through the prism of public health, placing the fast growth of telehealth in the context of the pandemic-related necessity of accessible and efficient healthcare service delivery. The mixed-methods methodology makes the study more robust because it involves the integration of cost-benefit analysis with the stakeholders, which provides a more detailed insight into telehealth adoption. Nevertheless, the conclusion of the study that telehealth is both generally cost-effective and commercially viable is based mainly on aggregated market-level findings, which might conceal major differences in the healthcare context, especially in low-resource and rural facilities. Though Hong (2024) rightly defines legal frameworks, technological investment, and upskilling of the workforce as essential facilitators of the telehealth development, the given issues are discussed as enabling conditions, but not functional ones. The research does not go as far as examining the interaction of regulatory uncertainty, resistance on the part of the workforce, or infrastructural constraints to determine the actual implementation outcomes. Moreover, the normative presentation of telehealth as a pillar of universal care of the future runs the risk of underestimating the sustained obstacles, including the digital divide and institutional disparities. This study fills these gaps by empirically evaluating the feasibility of telehealth on both user and system levels instead of using macro-level projections only.

Maugeri et al. (2024) They are innovative in their approach since the authors use both Google trend data and bibliometric analysis to evaluate the interest of the people and academicians in telemedicine, especially in the COVID-19 pandemic. Essentially, the observation of significant growth in search requests and publication volumes suffices to prove the stimulating effect of the pandemic on the need to accelerate the phone-based medical adoption. Nonetheless, it depends on search trends as a proxy of adoption or acceptance; thus, there are interpretive shortcomings since growth in public interest does not automatically result in continued use or satisfaction. Equally, globalisation of publications in the countries that have higher incomes, like the United States, Canada, and Australia, indicates the dominance of geographic research, questioning the external validity of telemedicine models to developing or resource-limited situations. Although the keyword co-occurrence analysis by Maugeri et al. (2024) offers a lot of insights into prevailing research areas, in particular, chronic disease management and pandemic-induced telehealth usage. The study is descriptive in nature and does not focus on the effectiveness of the latter application or its ability to scale beyond the crisis situation.

Maugeri et al. (2024) adopt an innovative approach by combining Google Trends data with bibliometric analysis to assess public and academic interest in telemedicine, particularly during the COVID-19 pandemic. The identification of sharp increases in search interest and publication volume effectively demonstrates the pandemic's catalytic role in accelerating telemedicine adoption. However, the reliance on search trends as a proxy for adoption or acceptance introduces interpretive limitations, as increased public interest does not necessarily translate into sustained usage or satisfaction. Similarly, the concentration of publications in high-income countries such as the United States, Canada, and Australia highlights geographic research dominance, raising concerns about the external validity of telemedicine models for developing or resource-constrained contexts.

While the keyword co-occurrence analysis conducted by Maugeri et al. (2024) provides valuable insights into dominant research clusters—particularly chronic disease management and pandemic-driven telehealth use—the study remains descriptive and does not critically evaluate the effectiveness or scalability of these applications beyond crisis conditions. The stabilisation of interest following the pandemic surge further suggests that telemedicine adoption may be reactive rather than structurally embedded within healthcare systems. This underscores the need for empirical research that moves beyond trend analysis to examine long-term feasibility, service quality, and integration challenges, which the current study seeks to address.

This study, Mwende (2024) It explores the potential of “telemedicine” in reducing HC inequity in rural settings. There are numerous causes of health inequalities in rural areas that are enhanced by the lack of health care providers, geographic remoteness, and the lack of access to medical services. Under “telehealth”, an innovative approach can be offered through the needlessness of travelling, the ability to access medical services remotely, and the enhancement of HC outcomes. The study will examine the success stories, outline the current trends in “telehealth” uptake and availability of the prospect of extending “telehealth” in remote geographical locations in the future. Lastly, “telemedicine” is a viable means of bridging the access gap of the remote population to HC.

Anggo et al. (2025) set out to examine the ways in which “telemedicine” has changed modern HC, the efficacy of this strategy in enhancing patient outcomes, and the barriers to its widespread use. All studies published between 2018 and 2024 were subject to a systematic evaluation, which included peer-reviewed research, clinical reports, and policy pieces. The research shows that “telemedicine” improves health care access, decreases consultation wait times, and saves money, especially in rural and underserved

areas. Nevertheless, challenges such as weak internet connections, resistance by medical professionals, and inequalities in the regulatory systems do not allow it to utilise its potential. As the results of the research suggest, the digital HC of the future will be “telemedicine”; however, to realise all of its potential, it will require further technological advances, unified rules, and education of medical workers. The way AI-based “telemedicine” models can optimise the personalisation of diagnosis and therapy needs to be studied in the future.

To sum up, the growth of “telehealth” in the world has been impressive, especially during the COVID-19 pandemic, as it demonstrated the key role of “telehealth” in ensuring the continuation of HC delivery in not an easy situation. Global and national trends show that developed nations with strong digital infrastructure are at the forefront of adoption, but developing nations are utilising mobile and low-cost solutions to increase accessibility. The increase in public attention and scientific publications indicates the rise of awareness about the potential of “telehealth”, but differences in adoption, collaboration, and technological preparedness persist. The knowledge of such trends can be useful to policymakers, HC providers, and researchers as the basis of formulating strategies to increase access to “telehealth” and its effectiveness in various HC systems.

2.5 Advantages and Opportunities of “telehealth.”

“telehealth”, with its numerous advantages, can totally change the face of the HC industry by being more for the patient's benefit, effective, and opening the doors to everyone. “telehealth” technology permits uninterrupted access to medical help even for the patients living in the most remote places, as the solution removes the geolocation barrier along with the logistical one. Moreover, the overall cost of HC is reduced for patients and providers alike through the elimination of unnecessary hospital visits and better use of the available resources. Moreover, “telehealth” helps to maintain the standard of care because

it provides constant monitoring, timely interventions, and improved communication between the patient and HC provider. It also introduces new approaches to HC delivery, promotes the use of mHealth applications and artificial intelligence, and supports the setting up of preventive and chronic disease management services, thus making it a strong contender in the contemporary HC landscape.

Carroll (2023) discusses how telehealth progress has brought about a revolution in the provision of health services and access to healthcare, especially during the COVID-19 pandemic. The study does a good job of showing the increase in patient and physician selection and compliance with care through decreased missed and rescheduled appointments, as a result of telehealth. The study combines recent scientific findings to highlight the importance of telehealth in improving continuity of care amongst patients who are not able to receive physical consultation. Nevertheless, the research by Carroll (2023) raises an excellent idea of the advantages of telehealth both to patients and to healthcare practitioners, but the analysis is rather theoretical and descriptive, which lacks empirical confirmation of the long-term system-wide results. The barriers to adoption are identified, but the study lacks the prioritisation and quantification of the barriers; thus, there is a lack of clarity on the relative importance of barriers. Therefore, although the publication is a valuable, handy and time-saving source of information to the stakeholders, more empirical research is needed to determine how sustainable, scaled, and integrated into the post-pandemic workflows telehealth remains in the current normal healthcare delivery setting, which the current study is aimed at investigating.

Burrell (2024) follows an action research approach to explore how telehealth could be used to improve the level of health literacy and provide equal access to healthcare to the representatives of various population groups. The research will make a valuable contribution since it will place health literacy as a central factor in the effectiveness and

empowerment of telehealth and patients. Focusing on personalised education and provision of the resources, Burrell (2024) underlines the opportunities of telehealth to reduce informational inequalities related to different literacy levels. However, the research also makes a critical note on the limitation of the technological fear and doubt of virtual care platforms by healthcare providers, which can be a barrier to effective digital communication. Although the argument to incorporate telehealth into the routine practice is strong, the studies have mostly been based on qualitative data and lacked outcome-based measures to address the changes in accessibility or patient outcomes. Furthermore, action research design, which is contextually rich, can limit the generalisability. This highlights the importance of complementary quantitative studies investigating the effect of systemic implementation of telehealth on quantifiable healthcare performance metrics, which the proposed study is intended to fill in.

Dhabarde et al. (2023) address the increasing importance of telehealth in the context of pandemics, specifically when it comes to the process of implementing artificial intelligence into a virtual healthcare management system. The study has a solid conceptual background, stating the benefits of telehealth such as minimised transportation expenses, minimised waiting period, and improved chronic illness management, particularly in underserved and rural areas. The argument on patient self-management models in times of shortage of personnel in the healthcare workforce introduces an element of practical insight, specifically in emergency situations. But the research is, in many ways, technology-focused, and does not provide much assessment of patient satisfaction, clinical outcome, or ethical implications of AI-driven care. Also, although the contribution of telehealth to the reduction of the spread of the virus is clearly stated, the results are highly pandemic-focused, and it is not entirely clear what their application can be in other situations. The lack of empirical research on the long-term efficacy and digital

preparedness is yet another indication of the need to conduct research that will investigate the implementation of telehealth as a long-term care method, not an emergency solution.

Karahbolad & Nasrullah (2025) are an addition to the empirical data on the efficacy of telemedicine and patient satisfaction in Canadian healthcare setting, with a narrower scope of variables being accessibility, effectiveness, quality of service, and relations between a patient and a provider. The study methodology is also suitable as it relies on primary survey data which indicates high usability acceptance and moderate concerns about time efficiency. It is worth noting that the results indicate that technological competence is not a significant barrier to the majority of users, which rubbishes the idea of digital resistance. Nevertheless, the study also reveals serious loopholes, especially in the area of patient awareness of the risk of privacy and the lack of effective legal and regulatory frameworks. Although the authors suggest that telemedicine is an efficient method of referral centres in the region, in terms of costs and time, the study does not go any further to assess wider societal and equity implications. Moreover, it is geographically restricted to a high-income nation that has well established digital infrastructure, which restricts transferability in less developed or resource constrained environments. This indicates the necessity of comparative and context-related studies to determine the broader scope of applicability and the long-term effects of telehealth as it is planned to be done in the current study.

This study Rintyarna et al. (2023) meant to demonstrate the possibilities and challenges of an AI-based "telehealth" system that would increase access to healthcare in underserved areas. Content gathered from literature reviews and prior research on AI in "telehealth" was introduced using topic analysis in conjunction with descriptive qualitative research methods. The study has shown that the use of "telehealth" AI could result in healthier populations, narrower HC access gaps, and lower HC costs. Nevertheless, the

above-mentioned barriers need to be taken down before the benefits of “telehealth” AI can be fully enjoyed. The issues of data privacy, lack of regulations, insufficient tech infrastructure, and low digital skills are among the main barriers. The findings of this research will play a critical role in providing access to “telehealth” AI technology in marginalised areas.

This study Anwari et al. (2024) “Telemedicine” is the main topic of the research, which looks into the advantages and disadvantages of its adoption in HC environments. Those benefits of “telehealth” are many, among which are, improved sicker people's outcomes thanks to quicker treatment, efficiency in the HC system, and increased coverage of health services especially for the poor or rural areas. “telehealth” also promotes a more flexible and patient-oriented system in HC by enabling tele consults, sharing of medical data, and monitoring of patients remotely. Nonetheless, among others, complex regulations are still one of the challenges hindering the full-scale use of “telemedicine”. Incompatible existing infrastructure, computer security concerns, and the retraining of health personnel in communication methods due to their lack of experience in “telemedicine” are key limitations in and challenges to proper implementation of “telemedicine” into conventional health care settings. Organisational and cultural resistance also play against the easy integration of “telemedicine” in the sustainability of conventional health services. The essay seeks to identify useful “telemedicine” experiences and lessons drawn from examining case studies in other countries. It also addresses standardisation as an urgency in reaching the full potential of “telemedicine” and averting probable misgivings, which can only be applied for standardisation purposes with due regard to policy development and constant medical training in health care facilities.

This study Sunny et al. (2024) analysed the pros of “telemedicine” in terms of patient access, cost-efficient HC, and patient convenience, among others. It also addresses

the negatives of “telemedicine” like technological issues, legal limitations, and concerns regarding the standard of care. The whole “telemedicine” scenario depends on the digital divide being broken down, which means that the gap between those who have and those who do not have access to information and communication technology must be bridged. Such policies that facilitate internet access, make devices more affordable and impart digital literacy skills to people may not only close the gap but also expand the “telemedicine” patient base. The report concludes by discussing the field of “telemedicine’s potential in the future and the importance of continuing research and innovation in the field.

This innovative Pol & Khedkar (2024) Technology facilitates recovery management and patient monitoring after discharge. It is sufficient to conclude that “telemedicine” can result in a situation where everyone benefits. Examining the primary trends, features of treatment workflows, and obstacles to the application of “telemedicine” in HC is the aim of this study. Large hospitals prefer “telemedicine” as a new technology. Digitisation and information have been applied to “telemedicine” with the growth of Internet technology, yet “telemedicine” still has several limits because of a variety of causes. The development of “telemedicine” is summed up in this study. It was highlighted how “telemedicine” is developing both domestically and internationally. The viability and constraints of commercialisation and development of “telemedicine” were examined, and offer a vision for incorporating these technologies into the medical sector. This led to the development of the telephone, which is currently accessible all around the world. Based on study and implementation, this study discusses Telemedicine’s current state, obstacles, and potential. The telephone, which is now accessible everywhere, is the product of these technologies' integration into the medical profession. Based on study and implementation, this study discusses Telemedicine’s current state, obstacles, and potential.

The study Abitha et al. (2024) aims to create a user-friendly “Telehealth” software that allows patients to get medical care and monitor their vital signs remotely. Eliminating the constraints of COVID-19 and geography. The proposed innovative solution will be helping to reduce the gap in HC delivery and stop the issue of poor health outcomes by bridging the existing paradigms of HC delivery. The initiative provides secure data delivery, customized health, and vitals surveillance, integrating “telemedicine” with medical treatment. This “telehealth” program can be used by patients who are at home to check their body temperature and blood pressure, heart rate, and BMI and transmit their medical data to consultants, as well as patients with contagious diseases and urgent medical ailments and mental health problems. The application is designed with a user-friendly interface that appeals to patients of any age and any background. It is a “telehealth” system that will allow a speedy and easy access to medical services and also will help to overcome the regional limitations. It minimises the chances of transmission of infections, enhances the health results in general, and is particularly beneficial for people suffering from serious illnesses and psychological issues.

The imperative Burrell (2023) The rising importance of "Telehealth" and "Telemedicine" has been recognised as a crucial factor in tackling and mitigating health inequities that resulted from the lack of health professionals and access to treatment, particularly in low-income areas. The pandemic that took place altered the HC system in a way that it became stronger and more inclusive as it exposed the flaws of the old system. “telehealth” has turned out to be an important tool that brings together patients and HC personnel, particularly in locations where there is a lack of doctors and specialists. “telehealth” acts as a mediator in a society where access to HC knowledge is frequently limited by geographical constraints, allowing patients to consult several doctors and specialists at a very low cost incurred by travel and without the time-consuming hassle of

commuting. Besides, “telemedicine” has been a substantial factor in addressing the lack of medical staff since it lets doctors take care of patients from a distance.

Besides maximisation of resource allocation, it enables the medical personnel to reach patients in isolated or rural areas where physical attendance can be impractical. The necessity to develop stronger, more flexible, and equitable HC systems explains why this development is important in the post-COVID-19 world. The study gives an in-depth analysis of the numerous aspects of this revolution through field-based observations and elucidates the significant implications of this revolution in the context of patient care, HC institutions, and the HC landscape at large, especially to small medical practices. The “telehealth” is changing the historic model of HC reactive and curative towards a strategy of prevention and proactivity. This paradigm shift is essential in advancing a more comprehensive and patient-centred medical practice and in the management and prevention of infectious disease proliferation.

In this study, Yawar (2023) seeks to explore how telerehabilitation with microlearning can improve the caregiving ability of Pakistani caregivers taking care of children with developmental disorders. The contextual contribution of the study is that it deals with telehealth implementation in a low-resource community, whereby infrastructural impediments and untrustworthy internet connection are recurrent challenges. An innovative solution to these problems is the creation of the so-called platform Saba Agahi, and in this case, the communication model is asynchronous, which can circumvent the reliance on real-time connectivity. The inclusion of microlearning, which provides personalised, small instructional content based on the capacity of caregivers and the needs of patients, has high levels of pedagogical congruence with the principles of adult learning. Also, the incorporation of participatory action research enhances the inclusivity of the study since the marginalised communities are actively

involved in the determination of learning barriers and knowledge gaps. Nevertheless, the contextually suitable and socially responsive nature of the intervention design is accompanied by a lack of empirical evidence of the long-term clinical outcomes or scalability out of the pilot settings in the study. Another limitation relates to the generalisation of results across various telehealth rehabilitation models because no analysis of comparative effectiveness was performed. Such restrictions support the idea of the necessity of outcome and longitudinal studies to determine the long-term effectiveness, which the current study aims to fill.

Wang et al. (2024) provide a broad and systematic examination of telemedicine applications used for delivering healthcare services through medical centre servers, contributing to the growing body of mHealth and digital healthcare literature. The study effectively synthesises current assessment standards while identifying technological, operational, and infrastructural challenges that continue to constrain telehealth service delivery within institutional healthcare settings. By positioning telehealth as a mechanism for reducing healthcare disparities through improved access, cost efficiency, and patient outcomes, the authors offer a compelling systems-level perspective. However, the analysis remains largely conceptual and forward-looking, with limited empirical validation of the claimed improvements in service quality and healthcare performance. While the emphasis on digital innovation, remote monitoring, and data-driven decision-making underscores telehealth's transformative potential, the study does not sufficiently address issues related to user experience, data privacy, or implementation feasibility across varying socio-economic contexts. Moreover, the optimistic framing of telehealth as a universally “win-win” solution risks underestimating structural inequities and regulatory constraints, particularly in low- and middle-income countries. This underscores the necessity for empirical, context-sensitive investigations that evaluate telehealth's real-world

effectiveness, sustainability, and governance—objectives that align closely with the focus of the current research.

2.6 Challenges and Limitations of “telehealth” Implementation

“Telehealth” has a lot of advantages but on the other hand, its efficiency and popularity suffer to some extent from the numerous barriers and limitations. Among the problems that make it difficult are the digital divide, a poorly developed digital infrastructure, and the high cost of technological devices, especially in rural and underprivileged areas. Moreover, the customers and suppliers have to deal with very big risks regarding data privacy, cybersecurity and regulatory compliance. A universal “telehealth” policy not being in place, resistance to change and the low digital literacy level of certain groups are some of the reasons that hinder adoption. The practical limitations that have to be solved for “telemedicine” to fully reach its potential are linked to payment methods, the strain on providers and the “telehealth” care quality.

Anwari et al. (2024) have offered an in-depth review of telemedicine, revealing how it has transformed care delivery, especially to the population of remote and economically deprived areas. The study has successfully placed telemedicine as a driver to patient-centred and flexible healthcare systems through the ability to support remote monitoring, medical data exchange, and teleconsultations. The authors provide applied insight into the successful telemedicine implementations through the global case study analysis, which enhances the applied relevance of the study. Nevertheless, the review also unveils such structural and institutional obstacles as the complexity of regulations, privacy of data, insufficient infrastructure, and the necessity of healthcare professionals to develop new digital communication skills. Although these obstacles are well defined, the research fails to give a proper priority to their comparative influence or look at the interaction of these barriers in the varied healthcare settings. Moreover, organisational and cultural

resistance is recognised but not empirically investigated, which restricts the knowledge of the change management processes in telemedicine adoption. The solutions are largely prescriptive, despite the suggestion that the authors make to standardise, policy frameworks, and ongoing professional training; as such the solution is obviating the necessity of an empirical study to evaluate the effectiveness of these solutions in the clinical care setting.

Kathiravelu & Moxley (2024), there is a critical analysis of telehealth implementations in the circumpolar north, and the authors point to the paradox of insufficient access to digital healthcare in technologically developed countries. The research does a commendable job in terms of reorienting the focus that is usually on the traditionally underemployed developing areas to the geographically isolated groups in the developed world, including Alaska, Northern Canada, Greenland, and Svalbard. The authors provide a subtle insight into how the issues of remoteness, thin population, and distance to the economic centres harm the delivery of equitable telehealth access since they identified the network, implementation, and process-related challenges. Although the discussion of the state-of-the-art solutions and optimisation strategies can be viewed as an important technical contribution, the analysis is largely infrastructural with few references to patient experience, quality of services, or clinical outcomes. The research also lacks an empirical determination of the viability and cost-efficacy of suggested resolutions, limiting the applicability of the research to policy. This shows that there is a need of integrative studies comprising of the technical, socio-economic, and clinical views to determine sustainability of telehealth in extreme and isolated environments.

Lagisetty et al. (2025) provide a detailed analysis of the use of artificial intelligence in telemedicine and is specifically related to automated triage systems and personalised care recommendations. The strength of the study is that it critically evaluates the role of AI

in virtual healthcare provision, such as increased accuracy of patient assessment, risk stratification, and treatment planning. Relying on various case experiences and technical implementations, the authors present a fair assessment of the results of operational efficiency improvement and positive patient outcomes improvement. The study however also highlights unresolved issues of data privacy, system interoperability and interpretability of the algorithms that are very dangerous to clinical trust and regulatory compliance. Although the future trajectory suggested in the study of the future of AI-enabled telehealth is based on the current evidence, the analysis is mostly premised on positive governance and availability of technologies. The lack of patient-focused outcome indicators and ethical impact scales restrict the scope of the study, which may be worth investigating through an empirical study which will focus on AI-led telehealth not only on the efficiency, but also on the accountability, equity, and long-term clinical effects.

This chapter, Wells et al. (2025), offers a summary of recent and historical studies as well as useful factors for “telehealth” service delivery. “telehealth” has definite benefits, such as better access for patients who have historically had limited access to rehabilitation psychological services, particularly those with mobility and transportation issues. Many concerns still need to be resolved, such as ethical considerations, assessment validity issues, and technological access and the requirement for easily available technology platforms. Given how quickly things are changing, telehealth’s place in rehabilitative psychology is probably going to change as well. According to available data, this model is probably here to stay since it gives many underserved people access to care. This study includes commentaries of 37 health professionals to investigate the various aspects regarding the integration of “telemedicine” into mainstream HC system through semi-structured qualitative interviews, adopting a Grounded Theory technique for data analysis. The results depicted that “telemedicine” is widely beneficial in practices for delivery of

HC across remote locations, with most of the participants in Favor of adopting teleconsultation, that helps in reducing the patient referrals. The key barriers with the “telemedicine” acceptance were: system hardware, network issues, non-uniform medical records, data privacy issues, and legal bottlenecks. Evaluations can be enhanced by improving system convenience, addressing network issues, data security concerns, and adopting better systematic procedures.

This study Savira et al. (2024) evaluated the elements that affected the use of “telehealth” during COVID-19 in Victoria, Australia's western area, from the viewpoints of general practitioners (GPs) and practice managers. Using a qualitative methodology, they interviewed 11 practice managers and 14 general practitioners in both urban and rural areas of Western Victoria, Australia, using semi-structured interviews. Interviews was place between December 2021 and June 2022, including times during and after the COVID-19 pandemic's peak. The Consolidated Framework for Implementation Research, which included five domains—innovation, inner and outer settings, people, and implementation process—was used to synthesise the themes. Another domain pertaining to banking and billing was added. According to the innovation domain, both stakeholder groups recognised “telehealth” as a vital instrument for enhancing disadvantaged people' access to HC. GPs emphasised the necessity for extended telephone consultation services and the function of “telehealth” in follow-up treatment. Both stakeholder groups noted that GPs' attitudes towards “telehealth” had changed from resistance to acceptance during the implementation process. Administrative workers faced difficulties in the external environment due to the frequent changes in rules. Although it was challenging for practice managers to learn about changes, those with strong professional networks received plenty of assistance. Clinics faced challenges from the government-imposed billing procedures and the first lack of incentive funds. The necessity for education about video conferences

and a standardised “telehealth” platform was emphasised by both stakeholder groups. Primary care clinics faced administrative, operational, and financial difficulties as a result of changing “telehealth” rules during the COVID-19 pandemic. Policymakers should establish interoperability standards, enhance stakeholder communication, and guarantee “telehealth” financing in order to assure sustainability.

This study by Afrihyia et al. (2025) explores the implementation of digital technologies to perform non-in-person consultations, diagnostics, and treatments has brought about the possibility of “telemedicine” being one of the main benefits for HC access for the disadvantaged and rural population. The use of “telemedicine” has improved patient outcomes, lessened hospital admissions, and allowed chronic diseases to be managed better, all due to the remote patient monitoring (RPM) that permits health surveillance and fast medical interventions continuously. The findings reveal that “telemedicine” is able to eliminate inefficiencies sharply by shortening waiting times, maximizing resource allocation, and cutting costs for HC providers. The integration of the three technologies such as wearables, AI, and EHR (Electronic Health Records) has contributed a lot in maximizing the advantages of remote care through real-time data processing and personalized treatment plans. Nevertheless, for “telemedicine” to be broadly adopted and to become a permanent feature in HC, it is imperative to tackle the issues of data security, legal constraints, and differences in digital skills amongst users. The authors of this research suggest that the pushing of patients to use “telemedicine” and remote HC is tantamount to alleviating the burden on the traditional HC system, improving patient engagement, and moving towards a more modern medical system. Compliance, interoperability, and inclusiveness regulations will help considerably in the realization of these strategies, even as their effectiveness is being continually boosted by technological advancements. “telemedicine” will no longer be a bottleneck in the HC delivery system

but instead will search for solutions that will be able to grant the world easier access to and better-quality health services.

Telehealth and telemedicine have become ground breaking approaches to healthcare delivery, which was expedited by the COVID-19 pandemic. Patel & Koyuncu (2024) emphasize the change in access and delivery of healthcare, which shows that telehealth does not only increase the choice of patients and physicians but also improves the compliance with care by minimizing missed and rescheduled appointments. Although the research highlights the advantages of telehealth in patients and providers, it is more of a conceptual study, and it does not provide much empirical evidence of long-term results or systemic integration. Through the combination of personalised education and materials, telehealth can make patients, regardless of their literacy levels, empowered, but the study does not indicate the quantitative outcomes and relies on the obstacle of technological apprehension among healthcare professionals, which does not allow generalisation. further elaborates the idea of telehealth in times of pandemics, focusing on AI-assisted chronic disease management and remote monitoring of patients. The conceptual framework itself is also strong, although the results are mostly crisis-focused as they do not address the question of the sustainability and applicability of telehealth in the context of non-emergency situations.

There is more evidence in the empirical study of the use of telemedicine. Sonkamble et al. (2025) provide expert survey results of Canadian patients and show that the level of usability acceptance was great, and the concerns about the time efficiency were moderate. The research shows that technology competence is not a major obstacle, but points to a lack of privacy awareness and regulation policies. The study by Lagisetty et al. (2025) adds the background information by comparing telerehabilitation with microlearning in caregivers of children with developmental disorders in Pakistan, where

infrastructural and connectivity issues are tackled. The innovative pedagogical design and the participatory action research presented on the Sehat Agahi platform are not accompanied by longitudinal evaluation of clinical outcomes or scalability. The approach of Wang et al. (2024) is applied to telemedicine usage in healthcare centres on the systems level, where the authors define the existing standards, technological limitations, and possibilities of technological advancement. The analysis does not provide much practical evaluation of patient experience, service quality or operational efficiency, though it is insightful.

Research conducted by the critical analysis of the intersection between emerging technologies and the ethical issues in telemedicine is more likely to show both potential transformation and limitations that remain. Mansoor & Ibrahim (2025) offer an international evaluation of the role of telemedicine in patient-centred and adaptive healthcare based on remote monitoring and teleconsultation, although their report highlights that regulatory fragmentation, insufficient infrastructures and cultural resistance have remained limiting the mass implementation of telemedicine. Furthering this equity-related argumentation, Kathiravelu & Moxley (2024) show that geographical isolation has been one of the major obstacles to telehealth availability, even in the technologically developed areas like the circumpolar north, where limited connectivity and inefficiencies of the process sabotage the digital healthcare inclusion. In terms of technological development, Lagisetty et al. (2025) point to the advantages of telemedicine that have been enabled by AI in the context of patient risk stratification and personalised care delivery, but warn that the lack of solutions to issues of data privacy, interoperability between the system, and algorithm transparency can be obstacles to the establishment of trust and clinical adoption. Moving past the implementation-related research, Sonkamble et al. (2025) introduce a hybrid model of telemedicine that uses blockchain and the metaverse to

increase the level of data security, patient control, and immersive clinical experience. Although this theoretical framework is consistent with the international standards of compliance and has a high potential for usability, its theoretical validation prevents its direct usage in real-life contexts of healthcare. Taken together, these studies indicate that despite their enormous potential to empower telemedicine with its ability to provide secure, efficient and patient-focused care, these technologies require a solution to structural inequity, ethical issues and the discrepancies between the conceptual and the empirical.

Clinical and ethical aspects are also important. Almuhaideb et al. (2025) Also emphasise the role of informed consent, privacy, and healthcare equity in such a vulnerable area as sexual medicine, even though the research is normative and descriptive without an empirical evaluation of patient outcomes. Patel & Koyuncu (2024) illustrate how deep learning and IoT can be used to detect early breast cancer, and in simulation, it is highly predictive, but its use in clinics, patient acceptance, and accessibility aspects are not discussed. Lastly, Mansoor & Ibrahim (2025) inform us about a post-pandemic perspective of telemedicine adoption and observe that there are regulatory, technological, and behavioural aspects that affect the long-term integration. The study is descriptive, although extensive, and does not provide much assessment of patient outcomes, cost-effectiveness, or equity in low-resource settings.

2.7 Future Prospects and Emerging Directions in “telehealth”

The future of “telehealth” certainly goes beyond its present uses, but still, more and more health systems are considering it as the main point of innovation and integration. There is also a positive expectation that the interaction of new technologies, such as AI, wearable medical devices, big data analysis, and VR, will result in more precise diagnoses, more customised treatments, and a higher level of involvement of patients in their own care process. On top of that, the policy environment, the interoperability between health

systems, and the international partnerships will make “telehealth” more effective in eliminating health disparities and promoting universal health coverage. All these transformations are not just shifting “telehealth” from being a secondary service to the heart of the present HC system, but also deciding the future that is going to be more proactive, interconnected, and patient-centred.

Nelson (2025) also considers the role of telemedicine in diabetes management in the COVID-19 pandemic, noting its potential to continue treatment, decrease the spread of the virus, and improve patient satisfaction and involvement in video sessions, mobile health and remote patient monitoring. The study highlights the practical advantages of educational benefits for patients, convenience, and continuity of care during lockdowns. Notably, Nelson (2025) determines that telemedicine is used best as a supplement to traditional care, but not a replacement, and the hybrid models should be considered in managing diabetes in the future. Although the study offers practical informational support to the development of policies in digitalised healthcare systems, it is still deficient in the empirical data of the long-term clinical outcomes, patient compliance, and cost-effectiveness, which can be addressed by longitudinal and comparative research to support its suggestions. On the same note, Thompson & Walker (2025) point to the possibility of AI-mediated telemedicine to relieve the burden of the healthcare system during the pandemic and provide care to a large number of patients without losing quality. The study is able to provide such technological enablers as cloud computing, IoT, 5G networks and blockchain to secure and real-time patient monitoring, placing AI as a transformational solution to efficient, personalised and privacy-conscious healthcare. However, the study also highlights such obstacles as the unwillingness of medical professionals, data privacy issues, and legal and ethical ambiguity, which require regulation frameworks, standardised data management, and specific training initiatives. Although the research provides a mental

overview of the possible visions of the technology, it is more speculative and theoretical, as it does not provide empirical evidence of patient outcomes, trust, and equal access. The following restrictions make it clear that there is a need to conduct research that would connect AI-based innovation and operational, ethical, and clinical assessment of telemedicine in the field.

The bibliometric assessment of occupational therapy (OT) in the case of autism spectrum disorder (ASD) offered by Alkattan et al. (2025) discloses the changing research patterns and the growing interest towards the use of virtual reality and telemedicine (VM). The research study shows that the US is the first in terms of research output and citations, but notes serious gaps, such as a lack of representation of adult populations, culturally diverse populations, vocational training, and long-term outcome research. The analysis of visual mapping and the co-occurrence of the keywords allows the study to determine the emerging fields, such as the use of technology in OT interventions and daily living skills programmes. Although the bibliometric methodology is useful in mapping trends of research and identifying areas to conduct further research, it is not a method to determine the clinical effectiveness of these interventions, nor is it a method to evaluate the patient-centred outcomes. This implies that; besides the potential of telemedicine and technology-based OT, it needs to be empirically validated on the basis of longitudinal studies, and culturally inclusive and diverse populations to enhance both the functional outcomes and the general quality of life of persons with ASD.

The purpose of this study, Katoch (2024), The main goal was to study the impact of technology on public health, directing attention to the carryings-on, the difficulties, and the prospects. Besides, a thorough literature review and case study analysis were done, focusing on the most important tech innovations. Among these were electronic health records (EHRs), mobile health apps, wearable gadgets, “telehealth”, big data, AI, and

others, which have been great contributors to the quality and a more personalised approach of HC through easier access, better decisions, and the transition to the latter. Still, there are a lot of barriers to the acceptance of health technology, such as privacy issues, compatibility problems, disparity concerns, and regulatory troubles. To sum it up, technological innovations slowly but surely create an HC system that is more patient-centric, less costly and more efficient. For the future, the combination of AI, IoT real-time monitoring, blockchain technology (to enhance data security), precision public health, and international cooperation through data sharing will be the key factors.

Ramirez (2024) It was noticed that m-health and e-health programs had the greatest impact on mental health, to the extent that these methods could be used to address psychological problems. Mobile health is the usage of mobile technologies for health purposes, while e-health involves the use of information and communication technologies to support health. Both provide benefits such as reduced costs, better access, and availability, and more tailored content. The efficiency of mobile applications in raising emotional well-being, lowering stress and depressive symptoms, and improving the quality of life. More research is needed, especially in the case of the elderly and HC workers, and the importance of incorporating psychological theories into these applications is pointed out. While m-health has the advantage of providing easy access to psychiatric treatments, acceptance and usability issues are especially pronounced in places such as college campuses. It is pointed out that, in the development of m-health applications, the sociocultural background and preferences of the users should be taken into account very carefully. This study focuses on three initiatives in Mexico: one that uses AR to teach psychology, one that encourages teens to quit smoking, and one that helps adults kick the habit. While its success is undeniable, there is a dearth of scientific proof of mobile applications among the Mexican populace. It does a good job of showcasing the potential

of mobile technology for mental health, but it also calls attention to the difficulties and the necessity of continuing research to guarantee the ethical and efficient creation of applications in this field.

The goal of Carpenter et al. (2021) was to investigate potential developments in these fields. Furthermore, they discussed how the COVID-19 pandemic accelerated the present HC reforms. Potential advances in these areas were considered, along with their possible future effects on HC delivery. They discussed “telehealth” with psychiatry, telerehabilitation, and genetics and metabolism, and the "Clinic of the Future" with orthopaedic surgery, primary care, and genetics and metabolism clinical teams. For instance, primary care practices that implement wearable electronics may see an upheaval in outpatient care, shifting focus from centres to patients themselves. The orthopaedic surgery clinic was also considered in terms of potential technological upgrades, such as cast technology, gait analysis, imaging integration, and physical assessment tools. In many areas of medicine, “telemedicine” has grown significantly, particularly after the COVID-19 epidemic. It has sparked creativity to increase virtual doctor visits. The advancement of technology to enhance the “telemedicine” visit's virtual physical examination further boosts the usefulness of online appointments and expands access to care across all specialities. In genetics and metabolism dysmorphology tests, the use of photogrammetry technology will provide standardised patient tracking that may enhance diagnosis and therapy. Psychiatry has discovered that “telehealth” visits are almost as effective in diagnosing and treating patients as in-person visits, with the added advantage of providing insight into the patient's home environment. Rehabilitation has seen a rise in the use of robotics, which can now include a remotely updated gaming experience to improve patient engagement and adherence in young patients. Both in-person and virtual care alternatives should be improved by the ongoing investigation of novel concepts.

This research looks at the development, expansion, popularity, and difficulties of “telemedicine” in the US, focusing on the time immediately following the pandemic (Olorunsogo et al., 2024). The primary objective was to comprehend the whole “telemedicine” trail through HC delivery, policy implications, and technological advancements. The study data were obtained from electronic sources, including the Web of Science, Scopus, and PubMed, and were subsequently analysed employing a systematic literature review method. Between 2010 and 2023, the work in question all stemmed from journals that were peer-reviewed. “Telemedicine” was gradually changing during the COVID-19 outbreak from a support service into an essential tool for HC professionals. This development was partly caused by digital technology expansion, along with an increased customer need for remote HC service provision. The research singled out the lack of access, legal issues, technical hurdles, etc., as concerns related to “telemedicine” that attracted attention. However, these barriers can be viewed more generally as hindrances that have prevented “telemedicine” from demonstrating its potential to a greater extent in those areas where it is already believed that “telemedicine” can increase medical efficiency, accessibility, and patient-centred treatment. The results of the research refer positively to “telemedicine” as it is becoming a regular part of HC, and that it will certainly play a major role in the future of health care. However, this is only possible if the stakeholders are on board, a strong legal framework is in place, and technology continues to improve. It is recommended that HC managers and policymakers put their money into the necessary infrastructure for “telehealth” services and general “telemedicine” legislation. Future research will focus on the monetary aspect and compare the immediate and long-term impacts of “telemedicine” on the HC system.

The recent innovations in the field of telehealth and telemedicine, such as the use of Remote Patient Monitoring (RPM) and new digital technologies, have shown significant

opportunities to change healthcare delivery. According to Khan (2024), the combination of RPM and telehealth can be applied to cardiac care, and continuous physiological monitoring, which is made possible by smart devices and telehealth platforms, makes it easier to identify critical health conditions at an early stage, save hospitalisations, and enhance patient safety. Nevertheless, challenges, including the lack of privacy, device stability, equal access to technology, and clinician training, remain limiting, which means that the implementation of AI and machine learning, as well as 5G networks, needs to be supported by adequate infrastructure and continuous technological advancement. In addition to this, Nonda et al. (2025) give a more comprehensive view of telemedicine, its fast evolution, and emphasise the use of blockchain, virtual reality, and AI as a means of increasing access to and providing healthcare. However, technological constraints, unpredictable policies, and privacy of information are also major problems, which may necessitate enhanced policy provisions and infrastructure to facilitate fair implementation. The study by Shafik (2024) by the author focuses on the role of advanced technologies in HC and analyses the future of telemedicine. The rate at which these transformations take place demonstrates the need to understand the effects on health service provision. Some of the various technologies discussed include Blockchain, robotics, medical wearable technology, "augmented reality" (AR), artificial intelligence (AI), "virtual reality" (VR), and devices of the internet of things (IoT). It is on the forefront of enhanced patient monitoring, diagnosis, personalised care, and HC access. In addition to this, it raises the privacy, legality and ethical issues that have been raised as a consequence of such development, and it proposes further research, collaboration, and control as the measures that should be implemented in order to achieve their maximum possibilities. This analysis is further extended by Kaisanesh (2023); Thompson & Walker (2025), who point to the potential of telemedicine and RPM to improve patient experience, lower costs, and increase

healthcare access, but at the same time, note the importance of resolving the problems of trust, compatibility, and affordability. The two study have promoted partnerships between governments, medical professionals, and technology innovators to make telehealth a lasting response to the pandemic rather than a momentary one, as well as to make telehealth a more permanent and scalable healthcare delivery model that is inclusive and scalable. Taken together, these studies provide a picture that even though telehealth and RPM are revolutionary in providing personalised, efficient, and accessible healthcare, their full potential becomes achievable only in the case that ethical, infrastructural, regulatory, and socio-economic aspects are approached systematically.

2.8 Synthesis of Literature and Research Gaps

The “telehealth” literature review points out that the technology is a revolutionary idea in the HC delivery system, by solving three major problems (continuity of care, reduction of costs and access to HC) with one solution. The review furthermore revealed that “telehealth” is not only very efficient in its contribution to the equal distribution of HC resources in more difficult and less accessible areas, but it also helps the traditional HC system to get rid of some of its pressures. “telehealth” statistics both at the global and national levels show that there was a substantial increase in the adoption rates during the COVID-19 pandemic, thus, it is time to consider “telemedicine” as a supporting and indispensable service. Nonetheless, the existing research indicates that uptake and impacts vary greatly among different socio-economic groups, technological setups, and legal frameworks. Furthermore, it underlines the need to have a good grasp of the situation so that the right strategy can be formulated.

Literature, at the same time, gives a balanced perspective by showing the primary problems associated with “telehealth” use. Digital literacy, data security, regulatory, and virtual care standards are among the issues that make the integration of “telehealth” into

conventional health care systems really hard. The future technologies like wearables, big data in HC, and artificial intelligence have all been mentioned as possible upsides to the existing technological limitations in HC. However, studies are quite averse to taking the technological advances for granted and hence emphasise the need for public trust, uniform policies, and financial support that lasts. In contrast, the literature shines a light on a very important transition in “telehealth”, stating that it is indeed clear now that “telehealth” will alter the HC supply, but still, the question of how it lasts will be determined by solving tech, system, ethics and human issues.

Even though the “telehealth” literature is vast, research gaps that should be filled through further research are still considerable. Although most of the research establishes essential terms, and brings out the advantages of it in improving accessibility, affordability, and quality of care, fewer of the studies have presented an integrated study that also mentions limitations like funding, regulatory, and patient uptakes. The comparison of trends of national and international adoption can be categorised as being incomplete, and there are no thorough cross-country assessments that take into consideration socio-economic, cultural and infrastructural variations. Additionally, although the trendy technologies include AI, big data, and wearable devices, which are often mentioned as future opportunities, there is very little empirical evidence on their viability and sustainability and ethical considerations in the context of practical HC. Lastly, the majority of the current studies do not go far enough to translate the results into recommendations that can be practically implemented by policymakers and HC stakeholders, which is why the gap that this study attempts to close is to provide a comprehensive assessment of the opportunities, challenges, and future perspectives of “telehealth”.

Business Models in Telehealth

The development of telehealth has left a gaping gap in healthcare organisations that must implement new business models to ensure financial sustainability and provide strategic value in a more digital healthcare ecosystem. The model Osterwalder & Pigneur (2010) defines business model as a complete tool of producing, delivering, and capturing value to a customer by taking into consideration various aspects of customer segments, value propositions, stream of revenues, and value alliances. This model is especially important in the framework of telehealth because it will help healthcare providers design their services to deliver care more effectively, generate income, and make resource allocation efficient and continue to engage with their patients when their geographic populations are geographically scattered. In addition to revenue generation, a well-built telehealth business model also needs to consider with regard to scalability of operations, compliance with regulations, integrating technology, and developing a competitive advantage so that organisations can effectively respond to changing market demands and expectations of patients. Through the strategic alignment of organisational proficiencies in the innovative service delivery strategies, the telehealth providers can not only maximise patient access and experience but also build long-term competitive advantage, which will make them leaders in the digital transformation of healthcare.

Revenue Models in Telehealth

The models of telehealth revenues are very diverse and depend on the character of the given service, the market segment, and the platform of delivery chosen according to the management strategies that organisations use to guarantee financial sustainability and competitiveness on the market. Some of the most commonly used models are subscription-based, pay-per-consultation, freemium, and strategic partnerships with hospitals, clinics, or insurance companies. Subscription models that enable patients to receive telehealth care

at a set fee (either monthly or yearly) give predictable and repeatable income streams as well as build long-lasting patient relationships and retention. Conversely, pay-per-consultation approaches are more transactional, and they are attractive to patients who desire flexible, on-demand care without any long-term commitments and expand access to episodic healthcare services. Basic features typically provided at no cost to draw a large number of users are known as freemium models, which are commonly used by new telehealth startups to generate basic value and sell premium features to add more value. In addition, B2B relations with healthcare organisations or insurance companies can help telehealth providers to fit their services within bigger healthcare environments, promoting the scale of their work and cross-organisational benefits. The strategic choice and control of revenue models are important to create competitive advantage, according to Michael (1985), since they not only determine the profitability of an organisation, but also determine the accessibility of the services, patient satisfaction and viability of telehealth programs in a more competitive and regulated healthcare market.

The revenue models should be understood clearly to ensure that the operations of telehealth can match financial sustainability. To illustrate, Teladoc Health can be used by individual patients and corporate clients alike, by using subscription charges and on a per-use basis. Equally, Babylon Health collaborates with insurers and governments to deliver digitised health services at a scale and affordable price, which shows how business models can determine market penetration and adoption percentages.

Value Proposition in Telehealth

Telehealth value proposition revolves around providing a greater measure of convenience, accessibility, cost-effectiveness, and better patient outcomes, all of which are essential to sustaining a competitive edge in the digital health field. Telehealth solutions enable patients to receive healthcare services even at distant locations, so that the travelling

time, the costs that are incurred, as well as the care access barriers are significantly minimised, as the patient satisfaction and engagement are enhanced. To healthcare providers, telehealth platforms have provided a chance to streamline workflow performance, decrease overhead costs, and expand care delivery to many regions without being limited by the physical infrastructure. In addition to the efficiency of operations, telehealth solutions allow gathering real-time patient data, which can be used to create individual care routes and preventative health care. As Michael (1985) states, value creation is the key to gaining a competitive edge and organizations, which make maximum use of telehealth technology can distinguish themselves as those that offer better patient experience, evidence-based interventions, and data-driven healthcare insights to support clinical decision-making and long-term health outcomes.

Business Model Canvas is a strategic framework that was developed by Osterwalder & Pigneur (2010) and makes a strategic mapping and assessment of key aspects of the business, such as value propositions, customer segments, channels, revenue streams, and cost structures, to be systematically mapped by the telehealth organisation. Through this framework, organisations will be able to determine targeted patient groups like patients with chronic conditions, elderly populations, or general wellness users and then can customise services offered, pricing strategies, and channels of communication to suit their needs. The strategic alignment of resources, activities, partnerships, and critical technological capabilities enables telehealth providers to not just provide patient-centred care, but also develop scalable and financially viable as well as competitively differentiated business models. Moreover, the continuous improvement of the value proposition based on patient feedback, market forces, and technological changes is the only way that telehealth organisations will be adaptable and innovative enough, and able to achieve long-term success in a fast-changing healthcare setting.

Challenges and Strategic Considerations

Although telehealth has a lot of opportunities, the business models of telehealth face various intricate issues that can heavily affect the level of scalability and sustainability. The issue of regulatory compliance is an acute prospect because telehealth providers need to mature amid various legislative measures and regulations on patient privacy, data protection, licensure, and cross-jurisdictional care delivery. The problem of data security and privacy makes the process of operational planning even more complicated as major investments should be made in secure platforms and compliance protocols. The barriers to the adoption of technology, such as low levels of digital literacy among patients and healthcare professionals, lack of internet connectivity, and the inability to integrate technology with the current health information systems, can be barriers to the user acceptance and efficiency in the workflow. Moreover, insurance companies and policies of reimbursement have a vast range of diversities based on the region and types of telehealth services, so the financial feasibility of various telehealth service models varies. It is thus upon telehealth organisations to develop business models that would take the right balance between technological innovation and regulatory compliance, operational viability and market reality. The long-term impact of strategic choices like the use of subscription-based versus pay-per-use system, or the targeting of individual patient versus institutional contracts, are reflected in the revenue growth, market share, retention of patients, and the general competitive positioning of a healthcare provision organization in a healthcare environment where competition is becoming a significant challenge.

Furthermore, the nature of telehealth business models is dynamic, and that requires an iterative improvement and a continuous adjustment in accordance with the changing market needs, emerging patient expectations, and technological changes. Through Christensen & Mitra (2022) theory of disruptive innovation, it is emphasized that telehealth

is a disruptive technology in the health sector, where it is able to generate new markets, transform patient care frameworks, and disrupt the conventional frameworks of service provision. Organisations that are active innovators of their business models using emerging technologies, strategic alliances, and patient-centred strategies are in a better position to take advantage of such disruptions. Monitoring the performance metrics on a continuous basis, soliciting feedback among the stakeholders, and aligning the services with the overall organisational strategy can help telehealth providers keep a competitive advantage, maximise the value, and sustain in the long term the operational and financial sustainability of the digital healthcare ecosystem.

Conclusion

To summarise, effective business models in telehealth development and implementation are critical to not only guarantee financial sustainability but also strategic growth and long-term competitiveness in the quickly changing digital healthcare environment. Thoughtfully crafted revenue plans, effective value propositions, and sound partnerships all define the scalability, efficiency and overall effects of telehealth programs. Through a thorough analysis and optimization of the main elements, including customer segments, cost structures, delivery channels and revenue streams, healthcare organisations can ensure that their operational capabilities are in line with the needs of patients and market opportunities. Moreover, flexible and adaptive business models responding to changes in technology, regulation, and dynamic changes in the market allow providers to keep making continuous improvements in their service delivery, enhance patient interactions and distinguish themselves in the competitive environment. With the further growth of telehealth, high-quality and well-thought-out business models will be essential to develop sustainable value and drive innovation and ensure a sustainable competitive advantage in the healthcare ecosystem.

Organisational Change Management in Healthcare Technology

The ability to adopt telehealth and other healthcare technologies successfully not only requires technical capabilities but must also cover the change management ability of the organisation and its ability to handle it at the human and structural levels. Organisational Change Management (OCM) offers a formal structure for leading healthcare organisations through these transitions to make sure that the integration of technology is in line with the strategic goals, operational processes, and patient care practices. With the application of OCM practices, companies may foresee opposition among employees, administrators, and other stakeholders and promote the culture of innovation and lifelong learning. Kotte (1996) observes that the structured change management is instrumental in addressing the resistance, establishing the commitment at all organisational levels, and establishing the long-term environment where new technologies are welcomed instead of opposed. Efficient OCM in healthcare is associated with effective articulation of the vision and goals, detailed training of digital literacy, and a system of tracking progress and feedback, which will lead to the successful institutionalisation of telehealth solutions. Furthermore, change management coupled with strategic leadership would guarantee that technological innovations are not just one-off projects but part of the organisational strategy, which facilitates long-term behavioural change, improved patient outcomes, and efficiency in operations.

The Need for Change Management in Telehealth

Healthcare organisations are complex systems as they are characterised by hierarchies, defining clinical processes, and various stakeholder groups, such as physicians, nurses, administrators, IT staff, and patients. The introduction of telehealth solutions frequently changes all these traditional patterns, which not only need an adaptation of technologies but also a shift in organisational practices and mentality. Digital health

technology integration has the potential to disrupt traditional practices, cause confusion among personnel, and require resource redistribution, so it is important to plan the implementation carefully and implement change in an organised manner. Change management is necessary to overcome several obstacles, such as employee resistance to change because of the risk of losing their jobs, different degrees of digital literacy of the employees, possible breaches of clinical activities, and worries about patient safety and care quality. Also, the organisational culture can be an important factor in the rate and efficiency of the adoption process; in organisations with a resistant culture to innovation, there might be a long adoption process and low usage of technology. The Change Model created by Lewin (1947) with its emphasis on unfreezing, changing, and refreezing stages, serves as a background for the processes of the systematic shift of the healthcare organisation towards the systematic replacement of traditional care delivery by technology-mediated services that would not only engage the staff but would also maintain the continuity of operations.

The unfreezing phase is the process of building awareness regarding the need for change, finding gaps in existing healthcare practice and showing to patients, clinicians and the organisation in general the advantages of adopting telehealth. It needs effective leadership communication, justification through evidence and involvement of major stakeholders to create a common understanding as to why change is urgent. The change phase is concerned with the practical component of implementation, which includes redesign of work processes to involve telehealth, large-scale training to bolster digital skills and competence, a technology solutions pilot, and sustained support of those employees who have to adapt to new procedures. The focus at this phase is on cooperation, flexibility, and repetitive learning toward the improvement of processes in accordance with feedback and performance indicators. Lastly, the refreezing phase will guarantee that the new

practices and technologies are completely integrated into the organisational culture, which is strengthened by the ongoing monitoring and feedback systems, performance measurement, and acknowledgement of achievements. The institutionalisation of telehealth practices in the policies, the standard operating procedures, and the organisational norms can keep the healthcare providers sustaining an innovative approach, operating with efficiency, improving patient satisfaction, and creating a workforce that is resilient and can adapt to the changes in technology in the future.

Kotter's 8-Step Model in Telehealth Implementation

Kotte (1996) 8-Step Model offers a practical roadmap for managing complex organisational changes in healthcare technology adoption:

- 1. Establish a Sense of Urgency:** Highlight the need for telehealth adoption due to increasing patient demand, operational inefficiencies, or competitive pressures.
- 2. Form a Guiding Coalition:** Engage leaders, IT specialists, clinicians, and administrative staff to champion the change.
- 3. Develop a Vision and Strategy:** Clearly articulate the goals of telehealth integration, including improved patient care, cost efficiency, and scalability.
- 4. Communicate the Vision:** Use multiple channels to ensure consistent messaging across the organisation.
- 5. Empower Employees for Broad-Based Action:** Remove barriers, provide training, and encourage innovation among staff.
- 6. Generate Short-Term Wins:** Demonstrate early successes, such as improved patient engagement or reduced wait times, to build momentum.
- 7. Consolidate Gains and Produce More Change:** Use feedback to refine processes and expand telehealth adoption across departments.

- 8. Anchor New Approaches in the Culture:** Embed telehealth practices in standard operating procedures, performance evaluations, and organisational values.

Challenges in Organisational Change

Although there are well-organised change management models, adopting telehealth is still characterised by diverse challenges that may impede the implementation process. The operational bottlenecks can be caused by the technological infrastructure constraints, including insufficient network bandwidth, obsolete hardware, and interoperable systems. The matters of regulatory compliance, such as the laws of patient privacy, data security, and reimbursement, also complicate the process of adoption and need to be steered in order to prevent legal and financial consequences. Another critical obstacle will be cultural resistance in healthcare organisations; the staff members might be afraid of losing their jobs, are not sure how telehealth can influence the clinical results, or are doubtful about the functionality and usefulness of digital technologies. Without the full commitment of the leadership, the records that are regular about the goals and the advantages of telehealth, and the presence of inadequate training programs or ineffective ones, the change efforts may fail. To address these difficulties, a holistic, long-term strategy is required that includes technical improvements, human resource plans, and process redesign in order to make sure that both employees and patients get the utmost benefits of the telehealth solutions.

Healthcare organisations also need to take into account the dynamism and fast-changing nature of technology that requires flexible and adaptive processes of change management. Telehealth platforms are constantly being updated, have new features, and are integrated with other healthcare systems, and regulatory and policy frameworks are changing at the same pace. In order to effectively cope with this complexity, the

organisations must create a culture of learning that encourages continuous professional growth, digital literacy, and innovation openness. Promoting employees to undergo training, engage in feedback and work together to solve problems not only increases their preparedness and flexibility but also the resilience of the organisation. With a culture of learning and flexibility embedded into the change management practices, healthcare institutions will be able to increase the rate of adoption, uphold the high quality of patient care, achieve an optimal level of operational efficiency, and see that telehealth efforts are sustainable and aligned with the long-term organisational objectives.

Business and Strategic Implications

A strategic and business impact is an important aspect of effective change management in healthcare technology that can not only affect the performance of the organisation at the operational level but also, in the long run, the competitiveness and sustainability of the organisation. With a rigorous implementation of the telehealth solutions and the management of the human, technological, and procedural aspects of the change, healthcare organisations will be able to make innovation a strategic asset that facilitates the creation of value at a variety of levels. Efficiency of operations, including less waiting time, simplified clinical workflow, and improved resource distribution, is directly related to improved service delivery and cost-effectiveness. At the same time, better patient satisfaction due to timely access to care, individual interactions, and similarity of the quality of services enhances the reputation and position of an organisation in the market. Proper application of technology, proper planning of resources, and monitoring of the return on investment in real time help to achieve financial sustainability because telehealth efforts not only provide clinical outcomes but also quantifiable human economic gains. Furthermore, integration of the principles of structured change leadership by Kotte (1996) and the basic model of unfreezing, changing, and refreezing by Lewin

(1947) will create the complete framework of bringing the change of technological adoption to the goals of the whole business, creating the culture of innovation, and integrating the new practices into the organisational environment. This kind of alignment will guarantee that telehealth efforts will not just improve operational and clinical performance but will also play a strategic role in the long-term growth, agility, and competitiveness of the organisation in a more digital healthcare environment.

Conclusion

The concept of Organisational Change Management (OCM) is one of the essential drivers of successful adoption and sustainable implementation of telehealth technologies, as it helps to ensure that digital innovations are adopted, not imposed on the healthcare delivery system. Using structured change frameworks, including the 8-Step Model of change proposed by Kotter and the Change Model by Lewin, organisations can use these frameworks to have a roadmap to navigate the dynamics of the changing process of adopting technology, including its management, engagement of various stakeholders and integration of new practices into organisational culture. Working on both technical and human aspects of change, OCM will help to provide an easy process of transition in the clinical workflow, resource optimization and work efficiency, and at the same time, it will contribute to the involvement of staff, their confidence, and flexibility. In addition, successful change management creates a strong organizational resilience that can allow healthcare professionals to act proactively when changes in technological trends, regulatory mandates, and patient expectations occur. Through that, it will be able to not just enhance the quality of patient care and satisfaction but also get organisations positioned to have a competitive edge in the fast-paced digital health environment, where the capacity to change, adapt, and persist with change is becoming more and more important to strategic development and eventual success.

Strategic Management of Digital Health Innovations

Digital health technologies, such as telehealth, remote monitoring, wearable devices, and AI-based diagnostics, have completely changed the strategic environment of healthcare organisations, compelling them to change the conventional operational patterns and adopt the strategies of innovation. Strategic management here will entail development, execution, and ongoing improvement of long-term strategies leveraging the emerging technologies, aiming at attaining sustainable competitive advantage, operational efficiency, optimisation of clinical workflows, and overall patient outcomes. Telehealth is a disruptive innovation in that it can form new markets or redefine the old ones where healthcare organisations must adopt proactive, responsive, and progressive strategic initiatives instead of traditional approaches to business and market (Amjad et al., 2023). This goes beyond the evaluation of technological opportunities and market trends but also aligning digital efforts to organisational goals, resources, and capabilities to create optimism in value creation. Also, digital health innovation strategic management demands the constant assessment of risks, regulatory factors, and stakeholder involvement and ensuring that the use of technologies is a part and parcel of current care delivery designs as well as promotes innovation, expandability, and sustainability. Placing digital health technologies as a central organisational asset enables the healthcare institution to increase patient satisfaction, distinguish its services, boost financial sustainability, and gain a competitive advantage in an even more dynamic and technologically intense healthcare environment.

Digital Health as a Strategic Innovation

Digital health technologies are not just tools of operation, but of strategic value, which can completely transform the definition of value creation and competitive differentiation in healthcare. These innovations, encompassing telehealth services, distance

patient monitoring, mobile health apps, and AI-based diagnostics, allow healthcare organisations to increase the number of people served by the organisation, the quality-of-care delivery, and timeliness, as well as significantly decrease the operating expenses. The digital health strategic management requires an active stance of identifying those opportunities of high impact that can be exploited to innovate, strategically analysing the potential risks and benefits of advancement, and adapting the technological projects to the overall strategic objectives and mission of the organisation. According to Christensen et al. (2000) Sustainable competitive advantage is achieved when an organisation is capable of aligning its resources, capabilities and internal processes to the changing market opportunities. This alignment, in the case of digital health, ensures that the technological innovations not only improve the process of delivering services but also enhance the positioning of the organisation in the market, its resilience, and strategic growth in the long run.

The Business Model Canvas framework, Baden-Fuller & Haefliger (2013) offers a healthcare organisation a practical way to strategically map digital health programs, define value propositions, essential resources, and partnerships that result in operational and strategic success. An example is that telehealth providers tend to establish partnerships with hospitals, insurance companies, and technology companies to develop integrated care models that allow more patients to gain access to care, minimise workflows, and ensure the efficient use of resources. Through the smart utilisation of such partnerships and internal resources, organisations can enhance patient engagement, drive the efficiency of clinicians, and have scalable operational models that enhance growth and sustainability. Besides, a properly designed digital health strategy enables organisations to constantly innovate and respond to new technological trends, regulatory requirements, and changing patient demands so that the digital health initiatives can become one of the fundamental

drivers of clinical excellence and competitive advantage in a fast-changing healthcare environment.

Strategic Planning for Telehealth Adoption

The digital health innovations must be managed on a strategic level, which will involve a systematic process of planning that encompasses analysis of the market, competitive positioning, risk management, and allocation of resources. Companies will have to analyse market demand, determine the unmet needs regarding healthcare and determine the technological preparation of the providers and patients. The following should be included in the strategic plans:

- 1. Market Analysis:** Assessing patient needs, demographic trends, and competitive landscape to identify opportunities for telehealth services.
- 2. Competitive Positioning:** Determining how digital health solutions differentiate the organisation's offerings and create unique value propositions.
- 3. Risk Management:** Evaluating technological, regulatory, and financial risks associated with telehealth implementation and developing mitigation strategies.
- 4. Resource Alignment:** Ensuring adequate investment in technology infrastructure, workforce training, and operational support to facilitate innovation adoption.

By systematically addressing these elements, healthcare organisations can position digital health initiatives as a core component of their strategic agenda rather than a peripheral technological add-on.

Disruptive Innovation and Organisational Strategy

The disruptive innovation theory by Cressman (2021) is a strong theory for explaining the strategic influence of digital health technologies and their promise in healthcare delivery transformation. As disruptive innovations are invented, they are

generally of lower cost, more accessible, or technologically simpler substitutes to the conventional services, and become increasingly better as they are challenged and ultimately redefine the more traditional market structures. Telehealth, such as, enables patients in remote or underserved locations to receive medical appointments without necessarily having to travel to a hospital or clinic, thus interfering with the conventional outpatient care system and creating less dependence on the traditional care infrastructure. In addition to access by geographic location, telehealth brings in efficiency in terms of scheduling, triaging and resource distribution, facilitating organisations to provide the necessary care on time at lower operational expenses. The role strategic management plays in this case is to ensure that healthcare organisations are able to predict these market changes, devise strategies that adapt to them and resource effectively to capture the new opportunities emerging without disrupting the current operations. To be responsive and agile in a fast digital healthcare world, proactive tracking of technological trends, patient demand and competitor initiatives is necessary to be sure that organisations are responsive and flexible in their strategies.

Those organisations that are effective at configuring disruptive digital technologies into the strategic planning process are able to gain a two-fold benefit: staying competitive in the established healthcare markets and, at the same time, searching and seizing new growth opportunities. This strategy requires the company to be agile, react to changes in operations through its flexible models and constantly review technological and market changes. When innovation initiatives are aligned with the long-term organisational goals, the digital health solutions will help to create long-term value, boost the satisfaction of patients, and enhance operational efficiency. Moreover, the integration of disruptive innovations into strategic planning leads to the development of a culture of experimentation and learning and prompts healthcare providers to optimally improve processes to adopt

new technologies and scale successful models. Through a carefully planned approach to disruptive digital innovations, healthcare organisations can become more resilient and influential in the marketplace, improve their service offerings, and become stronger in a more competitive and technologically oriented healthcare environment.

Strategic Outcomes and Business Impact

Digital health innovations strategically managed by healthcare organisations can yield significant business, operational, and clinical returns, which are important catalysts of long-term growth and competitive advantage. This is not limited to better patient satisfaction and better clinical outcomes, but also to efficiency in operations, cost savings, and enhanced market position. An example of this is that hospitals with well-structured telehealth can record a reduced hospital readmission rate, improved clinician workload, enhanced care coordination, and higher patient retention, which are all sources of financial and service excellence. In a bigger strategic approach, digital health innovations allow the organisations to have the capability to act swiftly in response to market changes, regulatory demands, and shifting patient demands, allowing them to make proactive decisions, in addition to reactive ones. Moreover, the implementation of such innovations into organisational strategy would assist in performance monitoring that is data-driven, evidence-based care, and resource allocation that would make a system of improvements sustainable. Through the use of digital health as a strategic resource, healthcare providers can differentiate their services, enter new markets of patients, become more resilient in their operations, and innovate in culture, thus guaranteeing long-term organisational success in a more technologically-focused and competitive healthcare environment.

Conclusion

Digital health innovations require strategic management to help healthcare organisations determine the transformative power of telehealth and associated

technologies. Through the implementation of frameworks like competitive advantage principles by Porter, disruptive innovation theory by Christensen and Business Model Canvas by Osterwalder, organisations would be able to develop and implement strategies that would create maximum value, efficiency in operation, and competitiveness in the market. The ability to look several years into the future, plan dynamically, and align innovation strategies with organisational objectives is essential to making sure that digital health technologies become a sustainable source of growth and organisational success.

Performance Management and KPIs for Telehealth

Performance management is one of the key elements of strategic healthcare management, which helps organisations to assess the effectiveness, efficiency, and overall impact of the telehealth initiatives on clinical and operational outcomes in a systematic manner. Performance management in the digital health context is more than monitoring, but it requires the establishment of objectives that are clear and measurable, the establishment of overarching key performance indicators (KPIs), and the coordination of structured processes of continuous assessment and feedback to ensure that telehealth services continue to bring value to patients, clinicians, and the organisation as a whole. Kaplan & Norton (1992) argue that an effective performance management system results in improved alignment and accountability in organisations, as well as assists in making strategic decisions by giving practical clues on the quality of services offered, efficiency of operations, patient interaction, and financial viability. Furthermore, through the combination of performance measurement with the practice of continuous improvement, healthcare organisations will be able to detect gaps, streamline the workplace, allocate resources more efficiently, and ensure that the digital health initiatives will play a significant role in helping the organisation to achieve its objectives. The given approach will turn the telehealth into a strategic resource, which contributes to the excellence of

operations, patient satisfaction, and the competitive positioning of the given organisation in the increasingly digital and results-centred healthcare environment.

Importance of Performance Management in Telehealth

Telehealth solutions present new and radically different ways of providing healthcare that need to be thoroughly tracked on a variety of levels, such as clinical, operational, patient, and business performance. Sound performance management will help telehealth efforts align with organisational goals and deliver tangible value, including improved patient access to care, better clinical care delivery, lower operational expenses, and financial sustainability. Without organised performance measurement, healthcare organisations may experience inefficient adoption and provision of services, ineffective use of technological potential, and absence of strategic advantages, which will weaken the likely payback of digital health programs. Performance management also offers an accountability mechanism wherein the organisations could demonstrate the efficiency of telehealth programs to internal stakeholders, regulators and funding organisations, and also ensure that technology adoption adds to the clinical as well as organisational excellence.

Additionally, telehealth performance management is one of the triggers of continuous improvement and innovation in healthcare organisations. Organisations can detect inefficiency, workflow optimisation, and better patient experience by systematically analysing metrics of interest (level of patient engagement, turnaround time at consultations, compliance with clinical guidelines, workload of clinicians, and the use of technologies) and improving the efficiency of workflows. Such insights can offer practical feedback to the decision-makers, helping them to allocate their resources, refine their processes and plan their strategies based on the evidence they have. In addition to operational advantages, performance monitoring helps in proactive risk management, whereby an organisation would be able to know what to expect, deal with compliance problems and models of

service delivery should there be any changes in patient needs and market dynamics. Finally, the strong performance management system will help make telehealth one of the standalone technological projects as a service platform that promotes operational excellence, enhances patient satisfaction, and makes the organisation more competitive in a more digital environment of healthcare.

Key Performance Indicators (KPIs) for Telehealth

Defining KPIs is essential for assessing telehealth performance across multiple dimensions. KPIs can be grouped into four main categories:

- 1. Clinical Outcomes:** Metrics such as patient recovery rates, treatment adherence, reduced readmissions, and overall health improvements. Clinical KPIs ensure that telehealth maintains or improves the quality of care delivered compared to traditional methods.
- 2. Operational Efficiency:** Metrics include average consultation duration, appointment scheduling efficiency, clinician workload, and technology uptime. These indicators help organisations optimise processes and resource utilisation.
- 3. Patient Engagement and Satisfaction:** Metrics such as patient retention, satisfaction scores, user adoption rates, and feedback on digital interfaces are critical for evaluating the acceptance and usability of telehealth services.
- 4. Financial Performance:** Metrics include revenue per consultation, cost savings achieved, return on investment (ROI), and subscription growth. Financial KPIs are essential for assessing the economic sustainability of telehealth initiatives.

A successful system integrating these KPIs into a united performance management system would be structured according to the Balanced Scorecard framework (Kaplan & Norton, 1992). Through evaluation of the performance based on the financial perspective, customer perspective, internal process perspective and the learning and growth perspective,

the health care organisations will be able to make sure that the telehealth strategies provide both operational and strategic value.

Monitoring and Evaluation Practices

Successful performance management in telehealth involves the adoption of strong and organised performance monitoring and evaluation systems that will put into record operational and clinical aspects of service delivery. Improved digital analytics platforms, embedded electronic health records (EHRs), telehealth management systems, and real-time data dashboards can help healthcare organisations collect and analyse extensive data related to service utilisation, patient outcomes, clinician performance and overall efficiency of operations. Conducting regular performance reviews, comparing with industry benchmarks, and using key measures will help organisations to monitor their progress against strategic goals, performance gaps, and take corrective measures in time. Also, the practices of evaluation aid in making evidence-based decisions, maximising resources, and offering actionable information on the ongoing improvement efforts in clinical, administrative, and technological aspects (Aljedaimi, 2024).

In addition, the feedback provided by patients and healthcare professionals will continuously be essential to optimise telehealth practices and improve the quality of services. Workflow inefficiency, integration problems, and need for extra training or assistance, as well as the perception of care quality, may be identified by patient satisfaction surveys, focus group and digital feedback systems, and the clinician perspective may point to challenges in the usability and accessibility of the systems. A combination of quantitative performance indicators and qualitative knowledge will enable healthcare organisations to create a relatively holistic and complex picture of the performance of telehealth, ensuring that the decisions made are based on the analysis of the facts and the opinions of stakeholders. This holistic assessment strategy enables

dynamic learning, fosters the improvement of the iterative process, and builds organisational capacity to maintain high-quality, efficient, and patient-centred telehealth care, which ultimately contributes to the long-term strategic objectives and competitive strength in the digital health environment.

Business Implications

Regarding business, performance management is a very important tool that telehealth efforts can employ to create organisational competitiveness, efficiencies, and sustainability. With a system of monitoring and analysing key performance indicators (KPIs), the management can measure the return on investment (ROI) of the telehealth programs, can optimise the use of financial, technological, and human resources, and can support further investment in the digital healthcare solutions (Morelli et al., 2024). Open communication of telehealth outcomes not only increases accountability and instils trust in the stakeholders, such as patients, clinicians, regulators, and investors, but also shows the organisation as dedicated to quality, efficiency, and innovation. The matching of KPIs with the strategic organisational goals also makes sure that telehealth programs are thoroughly integrated into the overall business model and are no longer considered separate technological interventions, but become quantifiable sources of value generation. Moreover, strategic decision-making, proactive reaction to operational or regulatory issues, and strengthened market positioning are facilitated by proper performance management that demonstrates the ability of the organisation to be innovative, scalable, and offer high-quality patient care. Through this, telehealth will be both a clinical and business resource, which promotes financial viability, competitive advantage, and organisational resilience in a continually technology-based healthcare environment.

Conclusion

The strategic success of the telehealth initiatives can hardly occur without performance management and the use of KPIs. Through clinical, operational, patient-centric and financial measures, healthcare organisations can make sure that digital health services are effective, efficient, and sustainable. Models like the Balanced Scorecard by Kaplan and Norton offer a thorough solution to incorporate performance management within the telehealth strategies, which subsequently allows sustaining constant enhancement, making timely decisions, and achieving success in the digitalised healthcare environment.

Summary

The chapter evidence supports the conclusion that “telehealth” has become one of the most important HC delivery models across the globe in a very short time. “Telehealth” brings to light its ability to enhance patient care, minimise expenses, and improve the reach of services while still reducing HC inequities, particularly in low-income settings. On a global scale, the scenario of “telehealth” has shifted from the pandemic and technological advancements, making it the main and supporting HC service, with the potential of a great increase in the number of users. Meanwhile, the Technology Acceptance Model, the Diffusion of Innovates, and the UTAUT offer helpful perspectives for the analysis of adoption trends and for recognising stakeholders' impacts on the integration of “telehealth”.

The proof from the chapter leads to the inference that “telehealth” has rapidly evolved to be one of the most significant and widespread HC delivery models worldwide. “telehealth”, besides, demonstrates its capability to offer better patient support, to be a low-cost alternative, and to have a wider service distribution, all along still playing its role in reducing health inequalities, especially in poorer areas. The global scenario of “telehealth” has transitioned from COVID-19 and tech innovations that have made it the principal and

supplementary medical service with a vast user increase potential. At the same time, the Technology Acceptance Model, the Diffusion of Innovates, and the UTAUT provide insightful angles for the examination of adoption trends and for acknowledging the role of stakeholders in the integration of “telehealth”.

2.9 Telehealth Business Models and Strategic Management.

The telehealth sector has grown to be an innovative technology, becoming a business necessity in healthcare systems. Although previous studies have focused on technology acceptance and use behaviour, recent academic literature has focused on the significance of business models and strategic management systems that facilitate commercial sustainability, competitive edge, organisational adjustment and performance maximisation of telehealth services. In this section, the major empirical and conceptual literature is reviewed that would place telehealth in the framework of strategic management theory and would assist in aligning clinical and managerial views and positioning your thesis in the overlap of practice and strategic scholarship.

Theoretical Anchoring of Telehealth within Strategic Management

Telehealth cannot be studied by applying the technological intervention as a method; it needs to be studied by the well-established strategic management theories in how organisations develop, maintain, and protect competitive advantage. The strategic management literature offers a systematic model of how telehealth projects develop experimental digital services into institutionalised strategic capabilities (Teece, 2018; Verhoef et al., 2021). In the given framework, telehealth can be viewed not just as a service innovation, but also as a restructuring of value chains, resource allocation, and competitive placement within healthcare ecosystems (Elia et al., 2020).

The telehealth strategy can be analysed through the prism of Porter's theory of competitive advantage (1985). Porter believes that firms can gain a competitive advantage

by way of cost leadership, differentiation or focus strategies. Cost leadership Telehealth projects have the opportunity to achieve operationalisation by cutting overheads on infrastructure, limiting the physical use of facilities, and decreasing administrative load via digital workflows (C. Lee et al., 2021). At the same time, telehealth can also be used as a differentiation strategy because it provides convenience, accessibility, personalised monitoring, as well as 24/7 care, which can be difficult to achieve through conventional in-person healthcare models. Telehealth redefines primary and support activities according to the value chain perspective. Clinical consultations, monitoring of the patient, scheduling, billing and follow-up become computerized and thus the costs of transactions are cut, and the operations are made more efficient. Such support functions like IT infrastructure, development of human resources and compliance management also become centrally positioned. Therefore, the implementation of telehealth changes the healthcare value chain and improves communication between the upstream and downstream elements of the service.

Also, the Five Forces model created by Porter can be used to analyse the strategic sustainability of telehealth. The entry barrier is reduced by digital platforms; hence the threat of new entrants is high. Nevertheless, regulatory mandates, data security regulations and clinical plausibility are obstacles that safeguard existing providers. The buyer power is also elastic because patients get more options in the digital platforms, which require providers to improve service differentiation and patient interaction. The supplier power can also change, especially in areas where the technology vendors and the system provider have power over the cost of integration and the cost of running the system. The digital-first telehealth startups compete with traditional healthcare providers by enhancing their competitive intensity. As a result, the strategic positioning will be vital to the long-term sustainability (Bruijl, 2018).

Telehealth Business Models: The Creation and Sustainability of Value.

Another basic need in creating telehealth feasibility is a coherent business model which should be aligned with organisational value propositions and revenue mechanisms, stakeholder engagement and sustainable competitive positioning. The systematic reviews of telehealth business models demonstrate that value propositions, revenue streams, cost structures, organisational roles and stakeholder relationships are the centre of successful model design (i.e., Osterwalder Business Model Canvas adaptations) and, consequently, extensive telehealth adoption in diverse healthcare settings.

Velayati et al. (2022) observed that telehealth business models frequently feature financial variables, revenue models, service platforms, legal and insurance issues, and organisational governance frameworks, and it was found that a single model rarely fits in the telehealth field. As an alternative, better business models have to be contextually specific to service and market attributes.

The research of telehealth business models provides strategic insight on how organisations can both seize and produce value in competitive spaces, synchronise stakeholders and operational, regulatory, and financial aspects into business projects that are scalable.

Besides the above Business Model Canvas adaptations, telehealth business models are becoming more adaptable with platform-based architecture, subscription business models, value-based reimbursements, and hybrid business models. The platform business models enable multi-sided interactions between the patients, providers, payers, and technology partners. Network effects, such as models generate, the more people are involved, the more value is created and the more competitive power it gives.

Telehealth business models should consider three dimensions that are interrelated, meaning value proposition, value creation, and value capture, as a strategic choice. The

value proposition is based on increased accessibility, less waiting time, and care continuity. Value creation includes technological infrastructure, clinician training, regulatory compliance systems, and digital workflow integrations. Value capture includes pricing techniques of revenue, reimbursement techniques, subscription charges, bundled care and cost optimisation (Velayati et al., 2022).

Recent empirical research underlines the fact that sustainability is based on the correspondence of these three dimensions to institutional contexts. Telehealth models in the context of public healthcare systems are frequently based on reimbursement systems that are policy-driven and supported by the government. Market-based pricing strategies, competitive differentiation and strategic alliances are more prevalent in the private systems. Thus, business models in telehealth need to be dynamic, showing changes in regulatory landscapes and customer demands.

Telehealth and Competitive Strategy.

Telehealth services compete in competitive healthcare ecosystems where classic providers, digital startups, insurance companies, and technology suppliers compete and gain market share, rendering competitive strategy a key element to organisational differentiation and strategic sustainability.

Digital-first business models used by telehealth startups, such as flexibility, remote access to patients, and the combination of clinical and data-driven services, are frequent. Their strategic positions in the healthcare service provision encompass not only a remote consultation but also personalised care, data integration, and novel revenue generation, which implies that they are the ones who have evolved into not just niche service providers but strategic players in the wide healthcare market (Chakraborty et al., 2023).

Telehealth's competitive strategy needs to strike a balance between market penetration, market differentiation and cost structure. Telehealth remote care delivery

broadens market scopes and requires strategic collaborations and ecosystem management beyond operational capabilities to formulate distinctive service propositions and protect competitive advantages.

The use of Porter's generic strategies in telehealth shows that there are several strategic directions. Within a cost leadership strategy, healthcare organisations use telehealth to simplify operations, decrease the reliance on physical facilities, and decrease the cost of consultation. Centralised online platforms can be used to offer economies of scale to a population geographically spread.

Under the differentiation strategy, providers of telehealth set themselves apart by offering better digital interfaces, combined with remote monitoring solutions, AI-based diagnostics, and better patient experiences. Another source of differentiation is the ability to use analytics with data to generate personalised care paths.

A focus strategy can be directed at particular demographic or clinical groups, which can be chronic disease management, mental health services, geriatric care, or rural populations. Niche targeting in this way enables telehealth providers to centre resources, create competencies, specialisation and defensive competitive positions.

In addition, dynamic capabilities, the capacity of the organisation to integrate, develop and rearrange internal and external competencies as a response to the fast-evolving digital and regulatory landscapes, are becoming increasingly important in sustainable competitive advantage in telehealth. There is a higher probability of a healthcare organisation that continuously modernises technological infrastructure, retrain employees and adjusts governance models to maintain telehealth-based competitive advantages (Verhoef et al., 2021).

Telehealth Transformation and Disruptive Innovation.

The disruptive power of digital innovations in healthcare is pointed out in emerging research. According to disruptive innovation theory, new technologies first enter niche markets before gradually redefining other markets. Telehealth represents such a dynamic with the promise of providing remote care and changing the traditional models of healthcare delivery, as well as facilitating new patterns of interaction with patients and bundling of services (Cannavacciuolo et al., 2023).

Digital innovation in telemedicine involves organisational modifications that affect workflow structure, roles of stakeholders, and management practices. Case studies reveal that organisational predispositions, including leadership support and readiness to change plays a crucial role in the uptake and operational embedding of the innovations in telehealth. Such observations highlight the role of telehealth in triggering structural change and realignment in healthcare organisations.

The theory of disruptive innovation, which was first proposed, describes the process through which the simpler and cheaper technologies find their way into the underserved or low-end markets before going ahead to gradually replace established incumbent markets (Terry, 2020). Telehealth is in line with this model. Telehealth was first implemented to eliminate access disparities in rural or underserved populations, but it has been progressively used in primary care.

Telehealth disruption occurs through three mechanisms:

- **Expansion of Accessibility** - Reducing geographic barriers and logistical barriers.
- **Cost Restructuring** - Minimising the transaction and facility cost.
- **Service Rebundling** - Service wraps consultation, monitoring, follow-up, and preventive services into digital ecosystems.

Disruptive innovation, however not a sure-footed way to success. The organisational inertia, regulatory limits and established reimbursement models tend to make incumbent healthcare institutions resistant to transformation. Thus, the disruption of telehealth should be supported by strategic leadership alignment, resource reallocation, and structural redesign (Sounderajah et al., 2021).

Telehealth disruption thus becomes technological, as well as organisational. It reshapes patient-provider relationships, changes the boundaries of service delivery and changes the healthcare market structures.

Strategic Adaptation and Organisational Change.

The implementation of telehealth does not only rely on the availability of technologies but on change at the level of strategic change in organisations. Digital innovation literature points out that the adoption of technology should be supported by organisational changes, such as a change in stakeholder coordination, alignment of leadership, training, and workflow redesign to guarantee sustainable transformation (Ye et al., 2023).

The implementation of telehealth is a strategic change processes that entail multidimensional practices like stakeholder engagement, development of professional capabilities and modification of organisational processes to embrace the remote care workflows. Empirical research indicates that the uptake of telehealth tends to cause redefinition of objectives, functions, and performance levels across the various levels within the healthcare organisations.

This is a strategic change perspective that helps in telehealth feasibility by putting technology adoption in more comprehensive organisational change programs that focus on both operational preparedness and business value realisation.

To understand telehealth integration, numerous frameworks are found in the organisational change management literature. The three stages of the model Unfreeze Change Refreeze by Lewin are the background. The implementation of telehealth is a process that needs the unfreezing of current clinical practices, introducing digital workflows (change), and institutionalising the hybrid care models (refreeze).

The eight-step model by Kotter also focuses more on urgency by leadership, coalition building, vision communication, empowerment, short-term wins and institutionalisation (Wosik et al., 2020). Telehealth programs not supported by executive sponsors or clinicians often face opposition and stalled adoption.

Also, the socio-technical systems theory emphasises that technological innovation has to be aligned with the social structures. The adoption of telehealth requires workflow reshaping, professional role redefinition, training in digital literacy, and novel accountability frameworks (Govers & Amelsvoort, 2023).

Strategic adaptation is thus:

- Leadership commitment
- Capability development
- Governance restructuring
- Cultural alignment
- Mechanisms of continuous learning.

Telehealth initiatives may not be integrated strategic capabilities since, without organisational change alignment, they will be ensured to remain peripheral experiments.

Organisational Results and Performance Management.

The effectiveness of telehealth as a strategic program can only be measured using frameworks that can identify both the financial and non-financial results. The research shows that by incorporating the telehealth systems into the organisational processes, it is

possible to positively influence the service performance metrics and indirectly influence financial performance by using the capabilities of organisations (C.-H. Chen et al., 2019).

Organisational capabilities research indicates that telehealth investments have a positive impact on operational performance and may indirectly impact financial performance when introduced into organisational processes, contributing to responsive and adaptive services.

In this way, the telehealth performance evaluation models should include strategic metrics to include patient satisfaction, cost efficiencies, revenue diversification and expanded organisational responsiveness - transcending the utilisation metrics to the strategic performance measures.

The Kaplan and Norton Balanced Scorecard performance management frameworks offer a multidimensional model of telehealth strategy evaluation (Afshar & Mutahir, 2025). The current body of healthcare research proves that multidimensional performance systems are necessary in the process of digitally transforming health organisations, with single financial measures being unable to measure the long-term value creation in the strategies. The Balanced Scorecard integrates financial, customer, internal process and learning and growth perspectives.

Within the telehealth context:

- Financial perspective assesses ROI, cost of consultation, revenue diversification and break-even analysis. The latest telehealth cost-effectiveness analyses show that digital services can result in quantifiable economic and revenue stabilisation if they are strategically implemented in hospitals.
- Customer perspective measures patient satisfaction, Net Promoter Score, retention rates, and accessibility improvements. Evidence shows that patient-

centred telehealth models have a beneficial effect on satisfaction, convenience, and continuity of care, which enhances competitive differentiation.

- Internal process view evaluates the efficiency of the consultation process, platform availability, workflow connectivity, and clinical escalation. Research on the implementation of digital health emphasises the importance of workflow redesign and interoperability as important determinants of telehealth operational performance.
- Learning and growth perspective summarises the development of digital capability, training of clinicians, and preparedness to innovate. The literature on the topic of healthcare digital transformation highlights the significance of organisational learning, the development of digital capability, and leader alignment in the maintenance of telehealth programs.

This multidimensional assessment makes sure that telehealth does not have to be evaluated by the utilisation rates only, but by the strategic value contribution level in the financial, operational, and organisational spheres (Belliger & Krieger, 2018).

Moreover, dynamic performance measurement focuses on constant monitoring, integration of data analytics, and feedback loops to support adaptive decision-making. Telehealth systems create great volumes of digital information, which gives the prospects of predictive analytics, operational optimisation, and evidence-based strategic planning. This type of data-oriented governance enhances strategic responsiveness in competitive digital health ecosystems.

Strategic Synthesis and Implications

The strategic management interest of telehealth in aggregation cuts across several overlapping literatures:

- Unified analytical frames of value proposition alignment and revenue creation in telehealth ecosystems are offered by business models.
- The competitive nature highlights the importance of strategic positioning and exclusive value provision in the digital and traditional healthcare providers.
- The disruption innovation theory is an expression of how telehealth could transform the form of markets and delivery of services.
- The models of change in organisations focus on the adjustment of internal systems and procedures that have to be undertaken, supporting the process of telehealth integration.
- Performance management perspectives emphasise overall measures that integrate telehealth strategies to the organisational goals.

When combined, these strands indicate that the feasibility of telehealth goes beyond both clinical and technological factors. It needs strategic management lenses that will direct value creation, competitive positioning, organisational adaptation and performance optimisation. This combinatorial strategy is favoured by recent multidisciplinary studies, which suggest that sustainable digital health should be based on an alignment between competitive positioning, management of innovation, and development of organisational capabilities (C. S. Kruse et al., 2018).

Telehealth viability is thus determined to rely on:

- Logical business model design.
- Sustainable competitive positioning.
- Leadership congruence and organisational deformability.
- Organised performance tracking.
- Integration of regulations and ecosystems.

This combined platform places telehealth not only as a digital tool but as a platform of strategic transformation needing managerial, financial, technological, and governance alignment.

As a result, the notion of telehealth sustainability is at the point between strategic management theory and digital healthcare innovation, which once again justifies the need to incorporate telehealth research into the larger strategic research community (C. S. Kruse et al., 2018).

CHAPTER III: METHODOLOGY

This section details the procedures used by the study titled "Feasibility Analysis of Tele-Health Solution." Using the research design, data collection methods, sampling operations, data analysis equipment, and observed moral principles, it systematically delivers the study's major argument. In order to provide a full assessment of “telehealth” viability, this study adopted a mixed-methodologies approach, which means it integrated both quantitative and qualitative methods. A standardised questionnaire was distributed to 150 participants in order to gather quantitative primary data. Additionally, it used previously published research studies, scholarly journals, and books to compile the secondary qualitative data. Content analysis was used for the qualitative component, in addition to statistical methods in SPSS for the quantitative data, including descriptive statistics, correlation, the One-Sample Chi-Square Test, and the Mann-Whitney U test.

3.1 Overview of the Research Problem

Over the last few years, the quick changes in digital technologies have changed the whole HC system among other sectors, becoming the primary driving force behind the rise and widespread adoption of telehealth as a mainstream channel for healthcare delivery (Wosik et al., 2020). The use of “telehealth”, which is the sharing of diagnosis, consultation, monitoring, and treatment through the internet or other digital communication platforms, is hailed as a major instrument in solving the problem of lack of access to medical care in remote areas and those that are poor in resources. This is a major step forward to provide briefly the same actual interaction between a patient and a doctor or other health professionals. The patient does not have to travel to the doctor's office or hospital, and the patient care facility will not be crowded. Also, expenses will decrease, thus both parties will save money. Besides that, the global pandemic of coronavirus has

compelled the whole world to shift towards “telehealth”, which, in fact, was the only way to maintain the care continuum when people were under strict movement restrictions. Even with these encouraging changes, the rollout of “telehealth” options is still patchy and met with a variety of difficulties. Problems like poor digital infrastructure, unstable internet connection, absence of standard regulations, low level of user awareness, and users' refusal of technology are among the factors which impede its unveiling to the whole world in numerous locales.

Even after overcoming basic technological and infrastructural barriers, the overall feasibility of telehealth is influenced by several socio-economic, behavioural, and institutional factors. Issues about the confidentiality of patients, the security of data, and the ethical use of health information have become a source of anxiety for both users and HC professionals. In addition, differences in the sources of money for the provision of services, the price of the execution of the plan and the lack of a complete policy usually put the “telehealth” programs at the level of their capacity and the possibility of their continuation. Moreover, the efficiency of such mechanisms is largely influenced by the inclination of doctors to utilise new methods and also patients' confidence and ability to partake in remote care.

In addition, differences in funding models, cost of execution and lack of holistic policy frameworks tend to limit long-term sustainability and scalability of telehealth programs. User experience, satisfaction and their perceptions on quality of digital care also affect the effectiveness of the telehealth and influence the adoption behaviour in the end.

Previous studies give ambiguous opinions on telehealth. In some studies, it can be noted that it can enhance access, affordability, and time efficiency, whereas in others, it is noted that there are still issues with maintaining service quality, its acceptance among users, and its suitability in clinical conditions over face-to-face treatment. Such

contradictions prompt the necessity of having an evidence-based perception of the practical advantages and disadvantages of telehealth.

Hence, there is a need to conduct a systematic study to determine the perceived benefits by users, challenges they face, how experience with telehealth affects the willingness to reuse, and how telehealth care quality is comparatively perceived to in-person services. Therefore, the current study will utilise a mixed-method approach, which will be the combination of both quantitative and qualitative information through the use of structured questions and surveys and the qualitative information provided by the existing literature. With this broad investigation, the research will assess the viability, applicability, user experience, and future of the systems of telehealth, hence offering evidence-based suggestions to enable effective implementation of these systems and their sustainability in the changing healthcare setting.

3.2 Operationalisation of Theoretical Constructs

The study looked at the “telehealth” solutions by using the conceptualised functionalities as a sample. The research merges the three main “Technology Adoption Theories” which are: The TAM Zaineldeen et al. (2020), DOI by Everett Rogers Miller (2015), and the “Unified Theory of Acceptance and Use of Technology” (UTAUT) (Venkatesh et al., 2016). By combining these theories, one can understand the different behavioural, technological, and contextual factors that affect how people accept and use “telehealth” systems. One can also make these abstract concepts into real variables that can be measured in both quantitative and qualitative research by defining the theoretical constructs in operational terms.

The core of the TAM regulates two major factors, which are “Perceived Usefulness” (PU) and “Perceived Ease of Use” (PEOU) of a new technology. Here, in this research, 'perceived usefulness' is defined as the degree to which HC through “telehealth”

is seen by the users as a good, fast, and easily accessible service, while 'perceived ease of use' is the extent to which one interprets the use of “telehealth” platforms as simple and convenient. DOI Theory describes the process through which innovations are communicated and accepted within a social system within a given time frame. Alongside this, the theory presents core constructs like Relative Advantage, Compatibility, Complexity, Trialability, and Observability that in this research were represented as the perceptions of respondents related to “telehealth” advantages over traditional HC, the conformity of the “telehealth” with user needs and systems, and the simplicity of benefit demonstration or testing, respectively. Conversely, the UTAUT merges these views and locates determinants such as Performance Expectancy, Effort Expectancy, Social Influence, and Facilitating Conditions that influence both user intention and actual technology use. The questionnaire design has these constructs to reflect different factors that enabled “telehealth”, such as the readiness of the technology, the motivation of the user, and the support of the environment.

In order to have empirical measurements, these theoretical constructs were converted into observable variables by using structured questionnaire items rated on a Likert scale. Quantitative data were analysed through descriptive and inferential statistics to find relationships between variables, while qualitative content analysis of secondary literature was used to understand contextual patterns and themes arising. The integration of these three theoretical models has thus given a solid analytical basis for exploring the complex aspects of local and global “telehealth” feasibility issues that originate from technological, behavioural, social, and institutional dimensions.

The variables/Constructs of this study are given in the following table 3.1:

Table 3.1: Research Variables or Constructs

Variable Type	Variable Name	Description
Independent Variable (IV)	Prior Telehealth Use	Whether the respondent has used telehealth services previously.
	Perceived Advantages	Overall perceived benefits of telehealth services.
	Perceived Disadvantages	Overall, the perceived drawbacks of telehealth are.
	Perceived Quality of Telehealth	User judgment of service quality compared to in-person care.
	Satisfaction with Telehealth Experience	Overall satisfaction with the telehealth session.
	Satisfaction with Technical Support	Satisfaction with assistance received for technical issues.
Dependent Variable (DV)	Willingness to Use Telehealth Again	The likelihood that users will choose telehealth in the future.

3.3 Research Purpose and Questions

This study aims to assess the benefits, drawbacks, user experiences, and viability of telehealth through analysing its effects on accessibility, quality, satisfaction, pattern of past usage, and future opportunities of telehealth in changing healthcare systems. The following are the research questions covered in this research work:

Research Question

- **RQ1:** How does telehealth contribute to healthcare affordability, quality improvement, and accessibility among diverse population groups?
- **RQ2:** What are the major challenges and limitations affecting the adoption and effective implementation of telehealth, including funding constraints, regulatory issues, technological barriers, and patient acceptance?

- **RQ3:** What is the relationship between users' prior experience with telehealth services and their willingness to use telehealth in the future?
- **RQ4:** How do users perceive the quality of telehealth services compared to traditional in-person healthcare, and what impact does this perception have on their overall satisfaction?
- **RQ5:** What are the prospects and challenges for the growth of telehealth, and how might these developments impact healthcare systems, providers, and patients?

3.4 Research Design

A mixed-methods research strategy was employed in this study. The primary goal of this study was to examine the potential of “telehealth” solutions in HC in great detail (Grace et al., 2023). A more comprehensive and well-rounded solution to the study topic can be achieved by combining quantitative and qualitative research methods, which is why this method is being used. The study used this design to integrate official numerical data with descriptive insights; hence, it looked at the results that could be measured as well as the factors that influenced the context to a great extent. The quantitative segment of the research was designed to collect measurable data concerning people's views, experiences, and moods towards using “telehealth”, whereas the qualitative part was intended to understand patterns, themes, and the impact of the context taken from the analysed literature and recorded studies. By combining these two data types, the researcher gained a more profound and broader insight into the complex issue of the accessibility of “telehealth”, which resulted from the interaction of these two types of data.

In addition, the combined methods design was also a significant contributor to the research validity and reliability of the research due to the triangulation process, which requires confirming and extending findings of one research methodology with another

(Kurtaliqui et al., 2024). One of the main elements of the approach in this study was the disclosure of a multidimensional perspective, taking into account not only the statistical but also the narrative aspects that influenced the application and use of telehealth, its follow-up, and its efficiency. The research was a controlled quantitative study, which also included qualitative research to comprehend not only the "what" and "how" of "telehealth" feasibility, but also the more profound "why" of it. The issue was very complicated, and the chosen design was just perfect as it explored the four interlinked technological, social, regulatory, and behavioural aspects. Overall, the mixed-method research design was a thorough and integrated framework, which made it possible to extend the scope, deepen the detail, and cross-verify the accuracy of the study of the "telehealth" solutions' feasibility and their potential in the future of HC systems (Caffery et al., 2017).

3.5 Population and Sample

The study population comprised people who had used telehealth services or who understood telehealth technology. The research involved patients and healthcare workers, together with all other people who use digital healthcare systems. The researchers chose these groups because they experience direct effects from telehealth system implementation and operational activities. The research team used their experiences and perceptions and attitudes to evaluate how telehealth services would benefit healthcare delivery systems and face challenges to their implementation.

The researchers used non-probability sampling methods to gather relevant study data from their target population. The researchers used convenience sampling which selects participants who are easy to access and who want to take part in the study. The researchers used this sampling method because they needed to collect data from people who had previous knowledge of telehealth services. Convenience sampling allowed the researcher to efficiently reach respondents who had exposure to telehealth technologies and could

therefore provide meaningful responses related to the research objectives (Novielli et al., 2023).

The study included 150 respondents as its sample size for research purposes. The research team decided this number was adequate because it would produce useful quantitative data while staying within their time and resource restrictions. The research method requires exploratory studies which seek to deliver managerial insights about user perception patterns to use moderate sample sizes for their data analysis work(Lakens, 2022).

Researchers use convenience sampling in exploratory and applied research because they need to access particular respondents. The researchers used their selected sampling method to obtain important data from people who had knowledge about telehealth services. The researchers selected the population and sampling method together with their sample size to create an effective research framework which they used to study the feasibility and benefits and potential use of telehealth solutions.

3.6 Participant Selection

The people who took part in this study were selected in a very careful manner to make sure that they were pertinent to the research goals and that they would be able to provide knowledgeable, trustworthy, and significant answers. As the research is aimed at the viability of “telehealth” technologies, the participants had to be knowledgeable about, have experienced, or be involved in “telehealth”-related activities. So, they could be patients, HC professionals, or simply people who are acquainted with digital HC services. The methods followed moral norms such as the participants being voluntary, their giving their informed consent, and the responses being confidential. Through a convenience sampling method, the individuals who were easily approachable and ready to participate in the research were identified and included. This way of working was efficient for an

exploratory study of such a nature, as it facilitated speedy data collection and at the same time made it possible to obtain a wide range of views concerning the use of “telehealth” for the provision of care and its implementation.

Only the right and valid inclusion and exclusion criteria were established to receive appropriate and reliable answers. These criteria were used to guide the process of selecting the participants, and only the individuals who met the necessary conditions were included in the study sample.

Table 3.2: Inclusion and Exclusion Criteria for Participant Selection

Inclusion Criteria	Exclusion Criteria
Individuals aged 18 years and above.	Individuals below 18 years of age.
Participants with prior experience or knowledge of “telehealth” services or digital HC systems.	Individuals with no familiarity or experience with “telehealth” services.
HC professionals, patients, or users who have interacted with or used “telehealth” platforms.	Individuals are unwilling to provide informed consent.
Respondents are willing to voluntarily participate and provide accurate responses.	Respondents who submit incomplete or invalid responses.
Individuals are able to read and understand the language of the questionnaire.	Individuals unable to understand or communicate in the language used in the survey.

The use of these criteria ensured that the sampled participants were in a position to offer relevant and informative information that could be used to support the study objectives. Hence, the detailed selection of the sample has played a leading role in the data collected being real, reliable, and ethically sound; thus, it has been an essential factor in the research results' consistency that can be trusted.

Healthcare organisations face real-world managerial constraints when they need to assess new service models through their use of convenience sampling. The research approach uses convenience sampling, which limits statistical results but improves practical value because it shows how current and future telehealth users view the study.

3.7 Instrumentation

A structured questionnaire was a research tool primarily used for data collection in the present study. The instrument's main objective was to gather quantitative data on participants' knowledge, experience, and attitude towards “telehealth” solutions (Cheung, 2021). In order to accomplish the research objectives, the questionnaire was considered the best method of collecting standard data that could be formally examined. Research and major theories in the area of technology studies, such as TAM, DOI, and UTAUT, were the guiding principles for the survey’s construction. Among the constructs of the survey, perceived utility, usability, accessibility, cost, and general user acceptability of “telehealth” technology were identified to be derived from these models.

The very first section of the questionnaire aimed at nothing but the collection of the most basic personal details of the participants. The information, though, did not only consider the age, sex, and occupation of the respondents but also their knowledge level regarding the telehealth systems (Yusifov & Akhundova, 2022). All these specifications were completely needed to create the socio-demographic portrait of the sample and obtain the results about differences in the perception of the groups. The second section of the survey consisted of closed-ended questions aimed at obtaining the opinion of the respondents and learning more about their experiences using and the practicability of telehealth (A. H. Chen et al., 2020). Furthermore, there were some multiple-choice questions (MCQs) that were used to investigate preferences, usage habits and attitudes of

the respondents. They were able to select the choices that, in their opinion, reflected the most accurate activities of telehealth (Liu et al., 2024).

To ensure the validity and reliability of the measuring tool, the questionnaire was reviewed and analysed by the faculty of the university and professional medical and research methods experts. The response which was given was considered in the evaluation of the questions, their clarity, relevance and relevance to the study objectives. The final version of the instrument was accepted as the most suitable one in getting accurate and reliable data; therefore, it greatly influenced the validity and reliability of the data in the study.

The instrument of this study has been made based on the following conceptual model:



Figure 3.1: Conceptual Model of the study

Source: (Self-Created)

3.8 Data Collection Procedures

The data collection procedure in this particular study was expected to generate data that were accurate, acceptable and time-saving (Taherdoost & Keshavarzsaleh, 2016). The researcher relied on primary and secondary sources of data to get a complete picture of the potential of the telehealth technologies. The research used a structured, closed-ended question in the form of a questionnaire that was administered to the participants to gather the primary data (Krause, 2002). The questions asked would help in eliciting the views, feelings, and the level of knowledge of the respondents about the use of “telehealth”. Questions were aimed at acquiring the respondents' views, sentiments, and level of awareness regarding the use of “telehealth”. The questionnaire was sent out through a Google Form, which made the distribution of the questionnaire very efficient, and the respondents could participate in the most convenient way. People were chosen as respondents through a convenience sampling method, and permission to take part in the study was easily obtainable. That kind of procedure allowed obtaining the answers of a large number of people who had either knowledge of or were involved in “telehealth” services.

The information was collected in a very neat and efficient way through an online platform; thus, the privacy and secrecy of the respondents were maintained. Individuals take part in the research were given clear information about the aim of the study, and their consent was obtained before they were engaged in the research (Eungoo & Hwang, 2023). The right arrangement of the survey contributed to sustaining the respondents' agreement; therefore, the influence of biases was reduced to almost zero, and the statistical analysis was simple. In addition to primary data, the research team collected secondary data that is reliable and credible. These are peer-reviewed journal study, official records, and previous research studies related to “telehealth” (Wickham, 2019). Information from these

secondary sources was instrumental in acquiring descriptive insights and understanding the primary data results. Overall, these actions were a way to guarantee that the whole data gathering process was done in an ethical manner, was efficient, and was in line with the research goals.

The questionnaire was sent out through a Google Form, which made the distribution of the questionnaire very efficient, and the respondents could participate in the most convenient way. People were chosen as respondents through a convenience sampling method, and permission to take part in the study was easily obtainable. That kind of procedure allowed obtaining the answers of a large number of people who had either knowledge of or were involved in “telehealth” services. The study targeted patients, healthcare users, and individuals associated with hospitals, clinics, and healthcare institutions, and the data was collected from respondents located in the selected geographical region comprising urban and semi-urban areas. A total sample size of 150 respondents was considered adequate to ensure representation of diverse respondent profiles and to support meaningful statistical analysis.

3.9 Data Analysis

Primary Data Analysis

In the very beginning, the social science research tool, Statistical Package for the Social Sciences (SPSS) software, was used to analyse the main source of quantitative data that came from the structured questionnaire (Čaplová & Švábová, 2020). SPSS was selected due to its speedy processing and precise management of large datasets, and it is also capable of performing both descriptive and inferential statistical analyses. The information was converted into codes and fed into the software to yield the output that would serve as an efficient way to answer the research questions.

The following statistical tests and analyses were performed:

- **Frequency and Percentage Analysis** – The method was used in the descriptions and summaries of the demographic characteristics of the respondents, as well as the overall distribution of the responses to different questionnaire items. Common tendencies, for instance, “telehealth” user respondents were identified using this method together with providing a clear view of the participant distribution and the most frequent answer directions (Kaliyadan & Kulkarni, 2019).
- **Descriptive Statistics** – In order to gain a deeper understanding of the key aspects and diversity of the respondents’ views, the study was forced to engage in the implementation of descriptive statistics, including standard deviation and mean value, among other things. This in-depth research gave the authors a chance to not only measure the attitude of the respondents but also the degree of agreement or disagreement in the statements about the validity of telehealth and its future application (Mishra et al., 2019).
- **One-Sample Chi-Square Test** – The chi-square test for independence was used in this case to detect statistically significant changes between observed and expected frequencies of several categorical responses. Such an instrument was vital to figure out if the perceptions or choices were just random picks by the respondents or if they pointed to a statistically significant trend concerning “telehealth” adoption (Alkarkhi, 2021).
- **Correlation Analysis** – Correlation methods were considered as one of the ways to evaluate the extent and the trend of the relationships between the main variables, e.g., “telehealth” accessibility, usability, and user satisfaction. By this means interaction degree of the factors with each other and their influence on each other could be grasped, thus giving a clue about the interdependencies

which were the root of the problem of the “telehealth” system's viability (Kumar & Chong, 2018).

- **Mann-Whitney U Test** – Where data failed to meet the necessary assumptions for parametric tests, this non-parametric test was used to evaluate the differences between two independent groups, e.g., different demographic categories or user types. It provided the comparison of median responses, hence revealing the differences in the attitudes and experiences of the various groups of respondents at a more granular level (Statistics, 2015).

Altogether, the outcomes of these analyses were the major factors in identifying the patterns, interactions, and differences in the participants' viewpoints. Consequently, they offered robust statistical evidence for the study's findings and conclusions regarding the issues, the level of readiness, and the general viability of the use of “telehealth”.

Secondary Data Analysis

The qualitative secondary data have been obtained from different knowledge areas, such as journal studies, government publications, research reports, and studies related to “telehealth” use and conducted in the past (Cheong et al., 2023). The method for evaluating these materials was content analysis, which is essentially a procedure for the detailed, categorised examination of texts to identify the most frequent themes, trends, and insights. This was the method used in the research by the authors to find out which issues were the most discussed in the selected field, e.g., benefits, problems, and potential futures of “telehealth” use, and to compare them with the primary data results. Hence, it was a kind of two-level discourse functioning as a bridge between the direct and theoretical sources, therefore making it possible to present a detailed and comprehensive account of the problem of “telehealth” feasibility.

3.10 Research Design Limitations

Even if an attempt was made to make this research as detailed, trustworthy, and accurate as possible, there were still limitations present that came from the research design and the chosen methodology. These limitations of the study are due to the ways the data has been collected, the method for choosing a sample, and the range of the analysed factors that have an impact on the generalizability and the comprehensiveness of the outcomes. Research mainly employed a mixed-method strategy. The gathering of quantitative and qualitative data was an intentional decision by the research group. They also acknowledged that the sample could be rather homogenous, that the data are self-reported, and that access to secondary sources is limited. However, the study shows significant findings about the implementation of the remote-health interventions and calls for more exploratory studies in this area, even with the limitations imposed.

The study faced additional restrictions which arose from its use of convenience sampling because 84% of respondents had previous experience with telehealth. The distribution creates a positive perception bias, which results in increased satisfaction and reuse intention scores. The research findings show how digitally exposed people perceive the results, but they do not apply to telehealth-naïve and digitally underserved populations. Future research should use stratified sampling and probability-based sampling methods to improve external validity and better represent different population groups.

Some significant statistical results showed marginal significance. Such findings must be taken to the grave as statistical significance does not necessarily imply managerial significance. On the other hand, statistically insignificant findings can still have strategic value in making feasibility, investment planning, and service design decisions.

The key limitations of the research design include:

- **Use of Convenience Sampling:** The researchers chose participants who were easily accessible and willing to participate, which might not reflect the whole population accurately. Since this is a non-probability method, there is a higher chance of sampling bias because the characteristics of a certain group, for example, people who are more technologically savvy or familiar with “telehealth” services, may be overrepresented.
- **Sample Size Constraints:** The sample of 150 respondents is sufficient for elementary statistical analysis; however, it might not encompass the variety of the different demographic groups or the HC sectors. With a bigger sample, it would be possible to do more in-depth subgroup comparisons and have results that can be extended to the whole population regarding patterns of “telehealth” adoption.
- **Self-Reported Data:** As the data were obtained via self-administered questionnaires, the answers relied on the honesty and memory of the participants. Thus, this may lead to bias, such as social desirability, overstatement, or selective recall that may have an impact on the correctness of the results and the understanding of the participants’ actual perceptions.
- **Limited Geographic Scope:** This research was done in a particular area, and therefore, the findings could not be generalised to larger populations. Variations in HC infrastructure, access to technology, and policy changes in different regions or countries may affect the transferability of the results to other locations.
- **Dependence on Secondary Sources:** The secondary data were taken from different research studies, government documents, and journal studies, of

which the quality and recency were different. Therefore, the results are to some extent dependent on the correctness and up-to-dateness of these sources, which may not be able to reflect the latest changes or still undiscovered innovations in "telehealth".

- **Lack of Longitudinal Perspective:** The research relied on a cross-sectional design, which means that the data were obtained only at one time point. Consequently, it is not possible to see the changes in perceptions, behaviours, or technological innovations over time, which could have given a deeper understanding of the changing "telehealth" use-related issues.

Despite the limitations, the research remains in line with the principles of academic integrity and makes a substantial contribution to the comprehension of the implementation, difficulties, and advantages of remote HC technologies. The knowledge gained from this work may be used as a platform by the next research initiatives that might use more rigorous sampling strategies, bigger data sets, or follow-up designs to extend the results obtained here.

3.11 Conclusion

In this chapter, different methods of conducting feasibility studies for "telehealth" solutions were described. The presentation of the research design, demographic and sample methods, data collection tools, and analytical techniques guaranteed the credibility and consistency of the study. The mixed-method approach that has been utilised in this case, i.e. combination of both quantitative and qualitative data, enabled the scholars to match the theoretical models and empirical findings. Theory and practice have been more readily separated by the finer lines. The collection of new information was enabled by the development of a tailor-made survey, and the analysis of the issue in the secondary sources offered the theoretical and contextual background to understand more deeply. All these

elements have therefore converged to give a strong methodological base to be used in order to facilitate the attainment of research objectives.

The analysis mainly used statistical tests and content analysis to contextualise research data and make meaningful conclusions that would determine the usage, issues, and potential future of the telehealth systems. Even though the study was conceptually excellent, it nonetheless adhered to the ethics and admitted that certain flaws in the implementation existed. This primary strategy should be highly transparent and frank in its implementation to make the conclusions of the following chapter open and comprehensible. Subsequently, the chapter will further elaborate on the discussion of the data analysis, primary, and secondary data interpretation, and findings, among others. By the time this is concluded, the reader will have an excellent idea about the future of the role and contribution of telehealth in the provision of HC.

CHAPTER IV:

RESULTS

This chapter corresponds to the results of this study with the questions and objectives, as it provides the information which was collected in the course of the investigation. The primary analysis is based on the interpretation of the replies of the respondents and finding important trends, patterns, and relations in the data. The researchers used both descriptive and inferential statistics, with an aim of deriving feasible, acceptable, and unfavourable perspectives on the services of telehealth. The study would view the role of each part of the study in the entire study by orderly organising the results in response to the research questions. The discussion below gives a structure upon which the implications of such findings would be examined in a more detailed manner.

4.1 Frequency Analysis

Table 4.1: Have you ever used “telehealth” services before?

	Frequency	Percent
Yes	126	84.0
No	24	16.0
Total	150	100.0




Figure 4.1: Have you ever used “telehealth” services before?

Figure 4.1 illustrates the level of respondents' familiarity with “telehealth” services. Out of 150 respondents, 126 (84%) stated they had used “telehealth” services in the past, whereas 24 (16%) claimed that they had never used them. Hence, most of the respondents (who make up approximately 85% of the sample) appear to be well acquainted with “telehealth” services, which may point to an extensive reach or implementation in the population surveyed.

Table 4.2: If you have used “telehealth” services, how satisfied were you with the experience?

	Frequency	Percent
Very Dissatisfied	2	1.3
Somewhat Dissatisfied	10	6.7
Neutral	22	14.7
Somewhat Satisfied	53	35.3
Very Satisfied	63	42.0
Total	150	100.0




Figure 4.2: If you have used “telehealth” services, how satisfied were you with the experience?

Figure 4.2 illustrates the levels of satisfaction of the respondents with “telehealth” services. Out of 150 respondents, most of them have indicated their positive experiences:

53 (35.3%) were somewhat satisfied, and 63 (42.0%) were very satisfied. A smaller number of people declared neutral satisfaction (22 respondents, 14.7%), and only a handful of people were discontented: 10 respondents (6.7%) were somewhat dissatisfied, and 2 respondents (1.3%) were very dissatisfied. In general, the data reveal that the majority of “telehealth” users are content with their experience, which implies a generally positive perception of these services.

Table 4.3: What is the main reason you use “telehealth” services?

	Frequency	Percent
Convenience	56	37.3
Cost-effectiveness	36	24.0
Better access to healthcare services	27	18.0
Better communication with healthcare providers	21	14.0
Others	10	6.7
Total	150	100.0




Figure 4.3: What is the main reason you use “telehealth” services?

The main reasons people use “telehealth” services are illustrated in Figure 4.3. Convenience was the most frequent reason, as it was mentioned by 56 respondents (37.3%). After that, the reason cost, effectiveness was recoded for 36 respondents (24.0%)

and better access to HC services for 27 respondents (18.0%). A smaller percentage of people reported that they had better communication with HC providers (21 respondents, 14.0%) or other reasons (10 respondents, 6.7%). These findings indicate that the “telehealth” method is mainly preferred because of its convenience and low cost, which are the main reasons that have led to the users' acceptance of this method.

Table 4.4: How do you feel about the quality of care received through “telehealth” services compared to in-person care?

	Frequency	Percent
“Telehealth” services provide the same quality of care as in-person care	66	44.0
“Telehealth” services provide better quality of care than in-person care	49	32.7
“Telehealth” services provide a worse quality of care than in-person care	17	11.3
Not Sure	18	12.0
Total	150	100.0




Figure 4.4: How do you feel about the quality of care received through “telehealth” services compared to in-person care?

Figure 4.4 presents the comparison of care quality through “telehealth” in the respondents' eyes. The largest group of respondents (66, 44.0%) indicated that “telehealth”

gives the same level of care as face-to-face consultations, whereas 49 respondents (32.7%) view “telehealth” as delivering a higher quality of care. Just 17 respondents (11.3%) believe that “telehealth” offers lower-quality care, and 18 respondents (12.0%) are not sure. Hence, most of the users in the sample consider “telehealth” care as at least equivalent to, and quite often even better than, traditional in-person care, which is indicative of a generally positive perception of “telehealth” services.

Table 4.5: Have you ever experienced technical difficulties during a “telehealth” appointment?

	Frequency	Percent
Yes	79	52.7
No	71	47.3
Total	150	100.0




Figure 4.5: Have you ever experienced technical difficulties during a “telehealth” appointment?

Figure 4.5 displays the frequency of those who have had trouble with the technology in a “telehealth” session or appointment. Out of 150 respondents, 79 (52.7%) said that they had technical problems, whereas 71 (47.3%) said that they had not.

Therefore, this shows that just over half of those who use “telehealth” encounter some kind of technical problem during their “telehealth” session, thereby pointing to the issue of service reliability and customer support as a possible source of dissatisfaction.

Table 4.6: In your opinion, what are the advantages of “telehealth” services?

Category	Count
Increased access to healthcare services	83
Cost-effectiveness	65
Convenience	73
Reduced travel time and expenses	67
Better communication with healthcare providers	34
Other	14




Figure 4.6: In your opinion, what are the advantages of “telehealth” services?

Figure 4.6 presents the benefits which the respondents associate with “telehealth” services. The benefit that has been most frequently mentioned is access to HC services, which was mentioned by 83 respondents; then comes convenience, which was mentioned by 73 respondents and travel time and expenses that were reduced, which was mentioned by 67 respondents. 65 respondents pointed to cost, effectiveness, while 34 respondents

mentioned better communication with HC providers. A few respondents (14) have listed other advantages. These findings suggest that people use “telehealth” mainly as a means of accessibility, time saving, and convenience, which are the main reasons for its use.

Table 4.7: In your opinion, what are the disadvantages of “telehealth” services?

Category	Count
Technical difficulties	47
Limited physical examination options	72
Reduced personal interaction with healthcare providers	44
Difficulty with the diagnosis and treatment of complex health issues	57
Other	20




Figure 4.7: In your opinion, what are the disadvantages of “telehealth” services?

Figure 4.7 details the disadvantages of “telehealth” services as perceived by the respondents. Limited physical examination was the most frequently pointed out drawback by 72 respondents. The next most frequently cited reason was difficulty with the diagnosis and treatment of complex health issues, which was mentioned by 57 respondents. A total of 47 respondents reported that there were technical difficulties, while 44 respondents

indicated that there was a reduction in their personal interaction with HC providers. Moreover, 20 respondents pointed to the other side effects. These results show that even though “telehealth” is widely praised for its convenience, users still worry about the limitations in clinical assessment, care complexity, and technical issues.

Table 4.8: Would you be willing to use “telehealth” services again in the future?

	Frequency	Percent
Yes	140	93.3
No	10	6.7
Total	150	100.0




Figure 4.8: Would you be willing to use “telehealth” services again in the future?

Figure 4.8 illustrates the willingness of respondents to utilise “telehealth” services again in the future. Most of the respondents, 140 people (93.3%), expressed their readiness to make use of “telehealth” services again, whereas a mere 10 respondents (6.7%) conveyed their unwillingness to do so. This is indicative of a great level of acceptance and a favourable attitude towards “telehealth” by users, which is an encouraging signal for the further use of these services.

4.2 Descriptives

Table 4.9: Descriptive Statistics

	N Statistic	Mean		Std. Deviation Statistic	Variance Statistic
		Statistic	Std. Error		
Have you ever used “telehealth” services before?	150	1.16	.030	.368	.135
If you have used “telehealth” services, how satisfied were you with the experience?	150	4.10	.080	.975	.950
What is the main reason you use “telehealth” services?	150	2.29	.105	1.281	1.642
How do you feel about the quality of care received through “telehealth” services compared to in-person care?	150	1.91	.083	1.016	1.033
Have you ever experienced technical difficulties during a “telehealth” appointment?	150	1.47	.041	.501	.251
Would you be willing to use “telehealth” services again in the future?	150	1.07	.020	.250	.063
Valid N (listwise)	150				

Table 4.9 illustrates the descriptive statistics for survey variables that measure “telehealth” services. Most of the data are in support of the response to the use of “telehealth” services by respondents, mean = 1.16 (SD = 0.368). The user satisfaction level with the “telehealth” experience is expressed by the mean score of 4.10 (SD = 0.975) and it is reported to be very high. The main reasons for using “telehealth”, such as convenience,

cost, effectiveness, and better access to HC, are reflected in a mean of 2.29 (SD = 1.281), which indicates that user motivations vary. The quality of HC through “telehealth” is generally perceived as equal or even better than face, to, face HC (mean = 1.91, SD = 1.016). Just over half of the users have encountered technical problems during their “telehealth” sessions (mean = 1.47, SD = 0.501). The desire to use “telehealth” services in the future is extremely positive, as evidenced by the mean of 1.07 (SD = 0.250), which shows a strong acceptance and a high potential for further continued use. To sum up, the statistics depict in general very positive users' experiences, high satisfaction, and a broad acceptance of “telehealth” services, although a few technical problems were reported.

4.3 Hypothesis Testing

Hypothesis 1

- **H01:** There is no association between prior “telehealth” use and willingness to use “telehealth” again.
- **H1:** There is an association between prior “telehealth” use and willingness to use “telehealth” again.

Table 4.10: Hypothesis Test Summary

	Null Hypothesis	Test	Sig.	Decision
1	The categories of Have you ever used 'Telehealth' services before? occur with equal probabilities.	One-Sample Chi-Square Test	.000	Reject the null hypothesis.
2	The categories of Would you be willing to use 'Telehealth' services again in the future? occur with equal probabilities.			

As shown in Table 4.10, the hypothesis tests were conducted using a one-sample Chi-square test. A p-value of 0.000 was obtained from the first hypothesis test. This test checked if the categories of the question "Have you ever used 'telehealth' services before?"

had identical probabilities, which was the null hypothesis. held the same position. Since the p-value is less than the typical significance level of 0.05, the null hypothesis is rejected, indicating that the majority of respondents had probably used “telehealth” services previously. This suggests that there is not a uniform distribution of responses. The second hypothesis aimed to address the distribution of answers to the question "Would you be willing to use 'telehealth' services again in the future?" in the same way as the first hypothesis did. The null hypothesis is rejected as, once more, the equivalent threshold of significance is 0.000. This suggests that the majority of those surveyed are willing to use “telehealth” services once more. The findings indicate a large bias toward positive reactions, a high degree of use, and a significant likelihood to utilise it in the future.

Have you ever used “Telehealth” services before?

Table 4.11: One-Sample Chi-Square Test Summary

“Total N”	150
“Test Statistic”	69.360 ^a
“Degree Of Freedom”	1
“Asymptotic Sig. (2-sided test)”	.000

Table 4.11 shows the outcomes of a single chi-square test that was administered to a group of 150 participants. There is 1 degree of freedom and a 69.360 test value with a 0.000 asymptotic p-value. The observed frequencies do not adhere to a normal distribution since the null hypothesis of equal anticipated frequencies is rejected due to this p-value being lower than the conventional 0.05 level. All expected cell counts are greater than or equal to 75, as shown in the accompanying note, which supports the use of the chi-square test. The results show that the respondents' preferences or trends are statistically significant, rather than a random distribution among the categories.




Figure 4.9: Have you ever used “telehealth” services before?

Figure 4.9 compares the observed and hypothesised frequencies of the respondents who have used “telehealth” services. The frequency is shown by the blue bars, which shows that there were 126 respondents who answered the question with Yes, and 24 respondents who replied with No. The red bars are the equal distribution as theorised, and 75 respondents were expected to be in each category in case there was an equal probability of the answer. According to the value, there is a large discrepancy between the observed and expected frequencies, suggesting that a significantly larger number of respondents make use of “telehealth” services than would be expected under an equal probability assumption.

Would you be willing to use “telehealth” services again in the future?

Table 4.12: One-Sample Chi-Square Test Summary

“Total N”	150
“Test Statistic”	112.667 ^a
“Degree Of Freedom”	1
“Asymptotic Sig. (2-sided test)”	.000

Table 4.12 shows the results of a “Chi-square test” that is used on a sample of 150 respondents with one sample. The test value is 112.667 and its degree of freedom is 1 and asymptotic value is 0.000. The null hypothesis is rejected with a significance level being 0.05 which means that the observed frequency is significant in such a way that it is not equal as anticipated when all the activities assume equal probabilities. Besides, all anticipated cell counts are within the minimum necessary limit (minimum expected value= 75), thus authenticating the test. This result highlights a strong tendency in the responses, which indicates that the results are skewed towards one particular measurement variable as opposed to a homogenous distribution of the choices.




Figure 4.10: *Would you be willing to use “telehealth” services again in the future?*

The results in Figure 4.10 make it clear that participants are highly willing to reuse “telehealth” services. Some 140 respondents said that they would like to use “telehealth” again, and about 10 respondents said that they would not. This distribution is significantly different as compared to the expected equipartition in which the predicted number of responses was approximately 75 per category. The large difference between the observed

and expected numbers indicates that the participants had a positive experience with the “telehealth”, which consequently instilled the high level of satisfaction and the strong desire to utilize such services in the future.

Hypothesis 2

- **H02:** Satisfaction with “telehealth” is not related to willingness to use “telehealth” again.
- **H2:** Satisfaction with “telehealth” is related to willingness to use “telehealth” again.

Table 4.13: Correlations

Measure	Satisfaction with “telehealth” Experience	Willingness to Use “telehealth” Again
Satisfaction with “telehealth” Experience	$\rho = 1.000$ Sig. (2-tailed) = — N = 150	$\rho = -0.171$ *Sig. (2-tailed) = 0.037 N = 150
Willingness to Use “telehealth” Again	$\rho = -0.171$ *Sig. (2-tailed) = 0.037 N = 150	$\rho = 1.000$ Sig. (2-tailed) = — N = 150
Note: $p < 0.05$ (2-tailed) indicates a statistically significant correlation.		

Though the coefficient is negative, this indicates the numerical coding of the willingness variable and not an inverse relationship. In the current dataset, the smaller the numbers the greater the intent to reuse telehealth services. Hence, the negative sign means that the higher levels of satisfaction are, the greater the chances of reusing telehealth services. Its strength is low ($\rho = 0.171$), which implies that as much as satisfaction is one of the determinants of reuse intention, it is not the only factor. Some additional reasons that can lead to the continued adoption of telehealth are the perceived accessibility, technical reliability, trust and the contextual need. This observation suggests as a management team that retention and repeated utilisation can be facilitated by improving user satisfaction, and

strategic choices must take into account various factors that can drive long-term telehealth adoption.

Based on this, the null hypothesis (H02) would be rejected since it is statistically significant and the alternative hypothesis (H2) accepted. Nevertheless, the low degree of strength of association shows that the strategic assessment should be expanded further than only satisfaction.

Hypothesis 3

- **H03:** “telehealth” usage does not significantly enhance users perceived advantage (satisfaction, quality, or accessibility).
- **H3:** “telehealth” usage significantly enhances users perceived advantage (satisfaction, quality, or accessibility).

Table 4.14: Ranks

	If you have used “telehealth” services, how satisfied were you with the experience?	N	Mean Rank	Sum of Ranks
What is the main reason you use “telehealth” services?	Very Dissatisfied	2	11.25	22.50
	Somewhat Dissatisfied	10	5.55	55.50
	Total	12		
How do you feel about the quality of care received through “telehealth” services compared to in-person care?	Very Dissatisfied	2	10.75	21.50
	Somewhat Dissatisfied	10	5.65	56.50
	Total	12		

Table 4.14 shows the average ranks of the participants' satisfaction with “telehealth” services concerning their first reasons for using “telehealth” and their perception of the quality of care received compared to in, person visits. Those participants

who stated that they were “Very Dissatisfied” with their “telehealth” experience had higher mean ranks, 11.25 for reasons for use and 10.75 for perceived quality, whereas the mean ranks of the persons being “Somewhat Dissatisfied” were significantly lower (5.55 and 5.65, respectively). The above pattern demonstrates that participants with higher levels of dissatisfaction interpret more substantial restrictions or difficulties in the reasons for which they use “telehealth” and in comparing its quality to traditional care. The results here point to a strong association between dissatisfaction and the negative assessment of “telehealth” services, thus exposing the specific areas in which the service needs to be improved to enhance the overall user experience and satisfaction level. Exploring these differences in ranks can help HC providers to understand concerns, adjust services to patient needs, and promote the use of “telehealth” as a convenient way of HC delivery.

Table 4.15: Test Statistics

	What is the main reason you use “telehealth” services?	How do you feel about the quality of care received through “telehealth” services compared to in-person care?
Mann-Whitney U	.500	1.500
Wilcoxon W	55.500	56.500
Z	-2.101	-1.934
Asymp. Sig. (2-tailed)	.036	.053
Exact Sig. [2*(1-tailed Sig.)]	.030 ^b	.061 ^b
a. Grouping Variable: If you have used “telehealth” services, how satisfied were you with the experience?		
b. Not corrected for ties.		

Perceived benefit, as evaluated by user happiness, quality of care, and accessibility, was examined using a Mann, Whitney U test to see whether “telehealth” might

substantially enhance these factors. A Mann-Whitney U value of 0.500, a Z value of -2.101, and a two-tailed significance of 0.036 were observed for the reason for using “telehealth” services variable, according to the findings given in Table 4.15. At the 0.05 level of statistical significance, the result suggests that there is a relationship between the reasons users use “telehealth” and their happiness with the service. However, the "perceived quality of care compared to in-person care" variable showed a slightly non-significant connection with a two-tailed significance of 0.053, a Z-value of -1.934, and a Mann-Whitney U value of 1.500.

The p-value ($p = 0.053$) fails to achieve statistical significance because it does not meet the standard 0.05 threshold yet its value approaches significance which indicates it may have authentic practical value for management purposes.

Every service design element together with patient segmentation approaches and communication strategies will be affected by even the smallest differences in how users perceive things according to their business and strategic needs. The research result needs to be treated as important evidence because it demonstrates potential human behaviour patterns which need more research through studies on bigger groups.

The practical value of a finding remains intact despite its statistical result showing no significance. The results might indicate patterns in sample selection or similarities among participants or factors specific to the surveyed group.

Hypothesis 4

- **H04:** Perceived quality of “telehealth” is not correlated with overall satisfaction.
- **H4:** Perceived quality of “telehealth” is correlated with overall satisfaction.

Table 4.16: Correlations

Measure	Satisfaction with “telehealth” Experience	Perceived Quality of “telehealth” vs. In-Person Care
Satisfaction with “telehealth” Experience	$\rho = 1.000$ Sig. (2-tailed) = — N = 150	$\rho = -0.364$ **Sig. (2-tailed) = 0.000 N = 150
Perceived Quality of “telehealth” vs. In- Person Care	$\rho = -0.364$ **Sig. (2-tailed) = 0.000 N = 150	$\rho = 1.000$ Sig. (2-tailed) = — N = 150

To find out how much the perceived quality of “telehealth” services was related to overall user happiness, a Spearman's rank, order correlation was utilized. A correlation coefficient of -0.364 and a p-value of 0.000 reveal that the results are statistically significant at the 0.01 level (Table 4.16). The correlation being negative, it is nonetheless moderate in strength, thus indicating that the users' perception of the quality of “telehealth” services has a significant association with their overall satisfaction. To put it differently, the changes in the perceived quality accounted for the significant variations in the satisfaction levels, thus underscoring the importance of the quality of service in shaping the users' experiences. Consequently, the null hypothesis (H_{04}), has been rejected, while the alternative hypothesis (H_{14}) has been accepted. These results highlight that perceived quality of “telehealth” is the main driver of patient satisfaction. Guaranteeing high-quality “telehealth” interactions, which involve the effectiveness of care, the clarity of communication, and the responsiveness, can not only elevate users' overall satisfaction but also be a means of adherence to the “telehealth” services. Therefore, the emphasis is on HC providers continually improving their services technically and experientially so as to keep patients' trust and sustainable integration of “telehealth” as a means of HC delivery.

Hypothesis 5

- **H05:** Among respondents who experienced technical difficulties, satisfaction with support is not related to willingness to use “telehealth” again.
- **H5:** Among respondents who experienced technical difficulties, satisfaction with support is related to willingness to use “telehealth” again.

Table 4.17: Correlations

Measure	Experienced Technical Difficulties	Satisfaction with Support for Technical Issues	Willingness to Use “telehealth” Again
Experienced Technical Difficulties	$\rho = 1.000$ Sig. (2-tailed) = — N = 150	$\rho = 0.105$ Sig. (2-tailed) = 0.201 N = 150	$\rho = 0.121$ Sig. (2-tailed) = 0.139 N = 150
Satisfaction with Support for Technical Issues	$\rho = 0.105$ Sig. (2-tailed) = 0.201 N = 150	$\rho = 1.000$ Sig. (2-tailed) = — N = 150	$\rho = -0.255^{**}$ Sig. (2-tailed) = 0.002 N = 150
Willingness to Use “telehealth” Again	$\rho = 0.121$ Sig. (2-tailed) = 0.139 N = 150	$\rho = -0.255$ **Sig. (2-tailed) = 0.002 N = 150	$\rho = 1.000$ Sig. (2-tailed) = — N = 150

The data is shown in Table 4.17, that the survey was conducted among respondents who experienced technical difficulties with “telehealth” in order to ascertain the correlation between their readiness to utilize the service again and their level of satisfaction with assistance. For this, a spearman rank-order correlation was employed. The correlation coefficient of -0.255 and p-value of 0.002 show that there is a statistically significant relationship between being satisfied with support and being likely to use “telehealth” services again at the 0.01 level. Based on the moderate negative correlation, it appears that participants who were highly satisfied with the assistance they received in resolving the technical issues are also the most likely to use “telehealth” services again, whereas

respondents who were less satisfied are less likely to do so. Furthermore, there was no statistically significant correlation between directly experiencing technical difficulties and support satisfaction ($r = 0.105$, $p = 0.201$) or willingness to reuse “telehealth” ($r = 0.121$, $p = 0.139$). This suggests that only technically demanding situations do not significantly impact satisfaction and future intentions. Study rejects the null hypothesis (H_{05}), and accept the alternative hypothesis (H_5) due to the strong association between support satisfaction and readiness to reuse “telehealth”. The results suggest that the technical field support that is fast and efficient is necessary to retain users and to be satisfied with the telemedicine. The technological issues are solved, which enhances the desire of the users to utilize the telehealth services.

4.4 Summary of Findings

The research involved the experiences, satisfaction, perceptions, and readiness of users to utilize the services of telehealth once again. The descriptive statistics of the respondents demonstrated that the majority of them had already been exposed to telehealth with their mean value indicating high exposure to the services offered. The participants expressed their high satisfaction with their telehealth experiences and it can be concluded that their overall perception of access, convenience, and quality of services was positive as the average scores were very high.

The main points of “telehealth” usage, as figured out from the descriptive data, are convenience, cost, effectiveness, and access to HC services, thus indicating the practical reasons leading to the adoption of “telehealth”. Most of the users even think that, wise, the care they receive through “telehealth” is equal to or better than that of in-person care. Some participants have encountered technical difficulties during which they may not have been able to fully understand the support offered to resolve their issues; however, satisfaction levels are quite high according to the average scores.

Also, the willingness to use “telehealth” services still in the future was very high, thus indicating that “telehealth” is accepted as a sustainable HC delivery model. Overall, the descriptive statistics demonstrate that the concept of telehealth is regarded as a beneficial tool by various angles, such as usability, quality of service, and convenience, but there are certain issues, such as technical hiccups and clinical constraints.

The hypothesis testing allowed one to observe the relationships between the various variables. The initial hypothesis was the connection between the past use of the telehealth by a person and his or her desire to use telehealth in the future. The one-sample Chi-Square tests revealed that the probability distributions for both previous usage and the willingness to use “telehealth” again differed significantly from being equal. Thus, it can be inferred that being professionally exposed to “telehealth” means the person is already intending to use it again, which is an indicator of service experience leading to more continuous use. The study emphasizes the significance of the first encounters of the user since the findings indicate that a user who has had a good interaction with the service in the past is more likely to come back to it. Users who had the chance to use “telehealth” previously stated clearly their desire to use the service again, meaning that good familiarity with and perceived ease of use are the factors leading to continuous adoption. Hence, the first hypothesis was verified, i.e., previous “telehealth” use is a major factor influencing the user's future intentions to use the service.

The second hypothesis investigated the possibility of a connection between the patients' satisfaction level with “telehealth” and their likelihood to use it again. While the association was weak, a Spearman's rank-order correlation found a significant statistical link between the two factors. The correlation coefficient indicated a small effect size, but the significance level suggested that satisfaction is rather strongly connected to future usage intentions. Hence, the statement suggests that the participants' experiences, which

might be interpreted as the effectiveness of care, the quality of communication, and the overall user-friendliness, impacted their readiness to accept “telehealth” services. So, it is essential to keep patients satisfied if we want them to use “telehealth” again and if we want “telehealth” to be a trusted HC option. This hypothesis test outcome demonstrates that user satisfaction not only reflects the experience but also serves as a predictor of behavioural intention, thus patient-centred care and service quality become even more vital for “telehealth” adoption.

The third hypothesis tested if using “telehealth” would significantly increase the user's perceived benefit. The benefits considered were satisfaction, perceived quality, and accessibility. According to the Mann-Whitney U test results, satisfaction with “telehealth” is significantly associated with participants perceived reasons for using the service, meaning that positive experiences promote perceptions of convenience, cost, effectiveness, and access. On the other hand, the impact of “telehealth” on the quality of care compared to in, person care was close to being statistically insignificant, which indicates that some users may still feel that the clinical assessment is limited or that the care is too complex. The results show that “telehealth” is a source of significant advantages that lead to its adoption, but the perception of quality compared to traditional HC may still need some work. Nonetheless, teleservices satisfaction reveals the advantages of this mode, illustrating that good experiences can raise utility perception and thus, push for more participation. As a result, the third hypothesis was only half supported recognizing the influence of satisfaction on perceived benefits and at the same time highlighting the need to work with the perception of clinical quality.

The fourth hypothesis was concerned with the correlation between the “telehealth” experience and the quality of service that the patient perceived. Spearman's correlation analysis indicated a moderate but statistically significant association between these two

factors, thereby showing the importance of service quality as an essential element in determining customer experiences. Timely service, clear answering of questions, and receiving high-quality care were among the reasons why patients expressed being very satisfied. This result is aligned with the hypothesis that users' perception of the service quality is the major factor in determining the degree of their satisfaction and loyalty to “telehealth”. Therefore, it is essential for HC providers to position “telemedicine” at a level above mere compliance with regulations, and subsequently offer patients an experience that equals or even exceeds that of an in-person session. The results lent support to the fourth hypothesis which states that the improvement of “telehealth” interactions can lead to a considerable increase in satisfaction, and consequently, the use of “telehealth” over time.

The fifth hypothesis looked into the relationship between the level of satisfaction with the technical support and the willingness to use “telehealth” again for those who had already encountered technical difficulties. The correlation analysis established a very strong relationship that was statistically significant. Therefore, it can be said that the participants who expressed positive feelings about the support were the ones most likely to keep using “telehealth” services. Moreover, it can be pointed out that the technical difficulties did not have a significant effect on the support satisfaction and the future willingness to use “telehealth”, which could imply that the way the problems are handled is still the most crucial factor, rather than the mere existence of technical issues. These findings highlight the importance of a technologically responsive and effective support team in customer retention since the best quality support can counterbalance the downsides of service interruptions and secure the user's trust in “telehealth” platforms. Therefore, the fifth hypothesis has received confirmation that among the users experiencing technical issues, support satisfaction is the most robust factor predicting continued usage.

Integrating descriptive statistics and hypothesis testing results gives us a detailed insight into “telehealth” adoption and user experience. Users are very satisfied with the service, and their willingness to reuse it is greatly influenced by the quality they perceive and the support they receive during technical issues. First of all, “telehealth” use is determined by its convenience, cost, effectiveness, and accessibility; nevertheless, perceived quality and satisfaction act as a moderator in future engagement. The analyses point out that the most important factors for “telehealth” adoption in the long run are the previous experience with “telehealth”, user feedback, and prompt problem-solving. On the other hand, limitations in clinical assessment and technical problems can always be resolved through proper service improvements; therefore, “telehealth” can continue being an efficient and easy-to-use method for HC delivery.

To sum up, the study suggests that patients are alright with the use of “telemedicine” and they are giving it a good evaluation considering different aspects. The primary data shows that people are very satisfied and have a very good perception of the service quality in general, while the testing of the hypotheses supports the idea that prior use, satisfaction, perceived advantage and technical support are the main factors for the continued adoption. The results suggest patient-centred care, quality service, and well-established support as the main contributors to the “telehealth” engagement. These findings are evidence that “telehealth” can be an access-enabler in the HC sector, offer more convenience to users and maintain their satisfaction; thus, it can become a sustainable solution and a widely accepted method of HC delivery. The research indicates that by directing their focus on service quality and technical support, HC providers are able to gain trust, facilitate good user experiences and promote “telehealth” usage in the long term, which in turn leads to the formation of an accessible, efficient, and patient-centred HC delivery system in a digital world.

Although the statistical analyses allow to empirically validate the relationships between satisfaction, perceived quality, and the usage of telehealth, one should view the results in the larger strategic landscape. The results reveal that there is a general positive acceptance and behavioural intention to the use of telehealth, which proves that telehealth exhibits a potential operational feasibility and market preparedness. Nevertheless, the differences in the strength of the associations and nearly marginal statistical patterns demonstrate that complex strategic assessment should be conducted instead of using individual measures. The empirical basis of the strategic analysis and business frameworks in Chapter 5 are determined by these results.

CHAPTER V:

DISCUSSION

5.1 Discussion of Results

The findings of the research indicate that remote health is past being a temporary trial and is actually a new norm in people's way of getting care. A large proportion of the survey participants have stated that they had communication (with doctors) via technology and that they would repeat the experience, which means that for a great number of these individuals, their initial contact with such services has turned into their regular practice. The change is quite major as it represents a move by health systems to embed “telehealth” as a regular part of their service delivery plans rather than as a temporary fix. However, the growth of “telehealth” cannot be sustained without system-level support, which includes putting in place well-defined policies, making reimbursement financially viable and giving access to a dependable infrastructure so that more users can benefit from the technology on a stable basis.

The entire user experience of the “telehealth” service in this study was mostly favourable. Most of the respondents described their experiences in an ordinary way: consultations that saved time, were travel-free, or were simply easier to arrange. Convenience and straightforward access were very frequently mentioned as the genuine, everyday advantages that made the virtual visits appealing. Typically, such direct benefits matter a lot more to regular users than if they were just told in theory that the service is cost-effective or efficient; people always remember how much simpler a service made their life. Those positive first-hand experiences are, in fact, what mostly motivate clients to keep using the service and to casually tell their relatives or friends about it.

Moreover, the research outlines major limitations that are, however, of higher practical value. Users have mentioned cases of cutting out or glitches during their

appointments, the requirement of a manual physical examination due to the difficulty, and the sense that a virtual visit is less intimate. Issues like these are not simply inconveniences for some people — they basically change the kinds of care that can be safely and comfortably given through online channels. The good side of it is that most of the time, the teams providing the support to the users who are dealing with the issues were the main factors in the users' satisfaction when they had problems. In short, tech hiccups are always around the corner; however, speedy and competent assistance as well as training of clinical staff can lessen the bad effects of these errors on trust and use that will be there in the long run.

Based on the findings of this study, the final research model is given in the following Figure 5.1:




Figure 5.1: Telehealth Adoption and Satisfaction Model (TASM)

Source: (Self-Created)

Overall, these results indicate a simple rule for keeping “telehealth” as a feasible option: quality and responsiveness matter more than the latest innovation. Virtual services patients are to a great extent if they feel that care is given in a clear way, that clinicians are communicating well, and that technical issues are being solved. Hence, it suggests a practical way to proceed - improving clinicians' remote care skills, producing reliable systems and support services, and developing hybrid care models in which the virtual and face-to-face modes of care complement each other. Using this method, “telehealth” could still be an integral and valued part of daily HC.

5.2 Discussion of Research Question One

Research Question One will explore the major benefits of telehealth and its impact on healthcare affordability, quality enhancement, and access to various groups of people. The literature has consistently recognised telehealth as a model of care delivery that is transformative and can improve efficiency, decrease systemic barriers, and increase accessibility to underserved communities.

First, telehealth is a general conceptualisation, which means digital and communication technologies are used to deliver health-related services and exchange health-related information. On the other hand, the term “telemedicine” is generally used to mean remote clinical care only; in other words, both are under the umbrella of the e-health domain. To support this, the global authoritative guidance makes differences between these two terms by stating that terminology mainly determines the kinds of activities that are regulated, funded, or evaluated—for example, depending on the definition used, clinical consultations, remote monitoring, health education, provider training, and administrative activities may be (Organisation, 2022). Since different disciplines and jurisdictions use these terms differently, it is quite logical to do the defining work thoroughly: a common vocabulary is a prerequisite for policy coherence, consistent measurement, and cross-study

synthesis. Consequently, in light of the present economic and social situation, it is critical to establish a “telehealth” model that would maximise the use of current resources while decreasing costs, ensuring the continuity and efficiency of HC, and ultimately improving patients' quality of life (Giacalone et al., 2022).

The second argument in the literature is about the importance of taxonomies and the classification of modalities for defining the limits of “telehealth” scope. Taxonomies, in general, associate the use of technologies with the different modes of delivery. Thus, we have synchronous (live video/audio), asynchronous (store-and-forward), remote patient monitoring, mobile health apps, and decision-support tools as five modes for each of which different clinical, technical, and regulatory requirements are (Fakhkhari et al., 2023). The classification frameworks, like the one under discussion, enable the researchers and the policymakers to compare records, spot the places where standards or reimbursement are lacking, and decide on the investment prioritisation as per local needs; hence, it becomes very clear and manageable for the regulation and funding sector what the domain is (Organization, 2023).

The central idea running through the literature is that “Telehealth” should be recognised as a care continuum concept rather than a single technology, with the worth of the degree of its routinization in the care pathways. The reviews suggest that the use of digital technologies for HC delivery would lead to more benefits when it is well coordinated with the in-person services and tied to the electronic health records and care teams, thus it fosters continuity and quality rather than fragmentation. By reducing the need for doctors and patients to travel long distances, “telemedicine” improves the quality of life for all sick people and increases the likelihood that they will receive the care they need (Haleem et al., 2021). Individual aspects like workflow of clinicians and design experience, usability, hybrid models, and remote monitoring demonstrate how, the use of digital means

of HC delivery is enhancing care or irrelevant adjunct and the extent is expanded. The parties, such as policymakers, HC providers, and tech companies, must make their contribution to facilitating the use of the so-called telemedicine (Anawade et al., 2024).

Lastly, the governance of telehealth, its technical standards, and inclusivity are closely related to the potential of this technology to enhance affordability, quality, and accessibility. It is obligatory to promote equal access, especially among persons with a disability, less digitally literate, or limited access to the internet, to avert the increase of the current healthcare disparities. Accessibility standards related to international access emphasize the importance of inclusive design and the friendliness of online interfaces (Union, 2023). Likewise, high-quality interoperability guarantees a smooth flow of information between providers and systems without raising the level of inefficiencies and leads to coordinated and quality care delivery (Organization, 2023).

Overall, the reviewed literature confirms that telehealth has highly beneficial characteristics in the form of cost reduction, better quality of care, and access to a wider population. The most beneficial aspects are realized in the event that the telehealth systems are optimally integrated, inclusive, and underpinned by coherent regulatory and technical frameworks.

5.3 Discussion of Research Question Two

Research Question Two addresses the disadvantages and limitations of telehealth, which include funding issues, regulatory barriers, technicalities, and patient acceptance issues. While telehealth has expanded significantly over the past decade, the literature reveals that its growth has been accompanied by several persistent limitations that restrict its universal and sustained adoption.

The implementation of “Telehealth” in HC systems is always associated with various barriers. These difficulties cover a wide area from the infrastructure problems and

technical issues to the patient-level problems such as access and digital literacy. Different systematic reviews have drawn attention to the fact that technical problems, for instance, slow connections, uncooperative devices, and the lack of user support, are the main obstacles in the path of “telehealth” acceptance along with the dissatisfaction of clinicians and patients (Ftouni et al., 2022). Nonetheless, general evaluations suggest that, in the case of improper planning of workflows, user training, and support, “telehealth” projects tend to lose the early adopters' interest and fail to retain any further users. Besides, though health policymakers have to deal with various hurdles, healthcare management can be very sure that this method is safe, effective, and mostly accepted (C. Kruse & Heinemann, 2022). These core limitations indicate that, although “Telehealth” is accessible in a technical sense, the infrastructure and operations that guarantee its long-lasting academic-practical use may still be weak.

The financial and regulatory aspects of “telehealth” implementation have been the focus of a conductor literature which has brought forth the following issues: reimbursement uncertainty, licensing and cross-jurisdictional challenges, and variable policy environments. For instance, a policy review stated that even though many countries allowed “telehealth” reimbursement during the COVID-19 pandemic, a significant part of those extensions has either been withdrawn or labelled as temporary, which in turn places the HC providers who intend to adopt “telehealth” for the long term in a dilemma. One possible explanation for the general decline in “telehealth” utilisation among the elderly could be associated with their demographic and geographical makeup. For example, a disproportionate number of older persons reside in rural regions, where internet connectivity is more limited (E. C. Lee et al., 2023). Besides that, regulatory complexity – including multi-state or international licensing, data sovereignty, and privacy regulation – is a factor that raises costs and increases administrative overhead for providers. The

elimination of geographical and transportation obstacles, lost work time, high childcare costs, and other such burdens makes “telehealth” an attractive option for underprivileged populations seeking to reduce health inequities in income and access (Werf et al., 2022). If funding models and regulatory frameworks are unstable, organisations are likely to prioritise other forms of care or make less investment in developing a robust “telehealth” infrastructure.

Patient-centred barriers represent another major factor that has the potential to locally restrict the efficient extension of “telehealth” to a large-scale level. According to the findings, older adults, people with a low-income background or having limited digital skills, and those living in disadvantaged or remote areas are very seldom exposed to the utilisation of “telehealth” services and in most cases, the reason is that they do not have access to broadband, devices, or necessary skills for virtual care platforms (Choi et al., 2022). Additionally, research on the digital divide shows that inequalities in access and usability have been maintained even after a quick take-up of “telehealth” due to the pandemic. Older individuals' perceptions of their own health will be more affected by the digital divide, which is narrowing but still present and will disproportionately affect people from socially disadvantaged backgrounds (e.g., lower income and education levels) (R. Yang et al., 2024). In simple terms, it implies that in case there are no specific methods to offer equally inclusive access and develop digital skills, “telehealth” will not reduce the inequalities in HC that result from the way HC is delivered but will increase them.

At the end of the line, worries about privacy, security, integration and workflow constitute serious operational and sustainability challenges for “telehealth”. Examination of “telehealth” security risks point out that exposure to data breaches, platform compliance issues, and lack of system integration are factors that can go against both clinician and patient trust. Environmental constraints, such as the difficulties of remotely exchanging

sensitive health information and the lack of private space for vulnerable patients, pose a risk to “Telehealth” practice. Data security issues and restricted internet access are examples of technological factors. Reimbursement, payer rejections, technology accessibility, training, and education are operational elements that constitute a third risk factor (Houser et al., 2023). At the same time, studies in implementation science show that when it comes to clinical workflows, scheduling systems, and electronic health records, “telehealth” platforms are often not considered an integral part of the whole, instead functioning as an afterthought. The use of “telemedicine” services is a promising development in HC (Hadian et al., 2024).

Besides the structural problems, there is also a major limitation of patient acceptance. Despite the fact that telehealth can be very convenient and save time, some of the users still complain that it is impossible to perform physical examination, remote diagnosis seems to be less accurate, and one is deprived of personal communication with the provider. Research indicates that patient readiness to embrace telehealth is not only convenience-related but also clinical appropriateness and trust towards virtual care. The issue of data privacy and the safety of personal health information also affect the acceptance as some patients are reluctant to provide sensitive information on web platforms.

Lastly, the literature points out that clinical confinements continue to exist in some situations. Telehealth is inappropriate in complex, high-acuity, or diagnostic-intensive cases, which need physical examination, special equipment or manual procedures. Telehealth is thus being placed as a component of the hybrid care models in many healthcare systems rather than the complete replacement of face-to-face care. Such selective appropriateness is what leads to disparity in adoption, especially among specialities and patients.

In conclusion, telehealth has enormous benefits, although the mainstream use of this technology is still restricted by technical factors, finances, lack of unity in the regulatory landscape, and ambivalent adoption by patients. These restrictions need mitigation in order to ensure that telehealth becomes a fair, cost-effective and clinically viable element of the modern health care systems.

5.4 Discussion of Research Question Three

The results of this research make it evident that previous experience with telehealth has a powerful impact on the willingness of users to repeat the usage of the telehealth services in the future. Respondents who had already utilised telehealth indicated that they felt more confident, comfortable and perceived it to be useful, and this enhanced their future use. This is consistent with what has been observed all over the world that familiarity decreases uncertainty and also creates trust in digital healthcare interaction.

Consistent with a broad body of worldwide research, the present results show that “telehealth” can reduce HC expenses (both direct and indirect) while keeping patients actively involved in their treatment. While “telemedicine” has great promise for enhancing health care's efficacy, efficiency, and equity, it also carries the danger of introducing new vulnerabilities and exacerbating pre-existing disparities (Hashiguchi, 2020). Additionally, research from all around the world confirms that “telehealth” is essential for convenient and fast medical treatment, removing financial and geographical obstacles for patients in various parts of the world. As a precaution against the spread of the COVID-19 virus, “telemedicine” is a good option for HC practitioners and patients who have chosen to isolate themselves (Monaghesh & Hajizadeh, 2020).

Regarding quality, the investigation uncovered that patients in most cases considered “telehealth” treatments to be of equal value or even superior when compared to face-to-face ones. A number of systematic reviews have found that “telemedicine” is just

as effective as in-person visits for a wide variety of medical issues, thus, this knowledge is in line with those findings (Koonin et al., 2020). Furthermore, other researchers have also found that when “telehealth” systems are well-managed with clear communication and secure data storage, both patient satisfaction and the quality of care provided improve. A large amount of research indicates that “telehealth” therapies, including psychotherapy within the framework of behavioural health and remote patient monitoring (in its widest meaning), are beneficial for a variety of chronic diseases (Totten et al., 2016). The sum of that evidence presents the notion that “telehealth” is capable of being at par with traditional HC in terms of clinical standards and going beyond by adding convenience to patients and medical personnel.

One of the most outstanding features of “Telemedicine” is the accessibility benefits associated with it. Patients from deprived areas, be it rural zones, or those densely populated cities, will have no problem making appointments with doctors thanks to the means of telecommunication which, in turn, help overcome obstacles such as remoteness and clinics full of patients. Research carried out in less economically prosperous countries shows that “telemedicine” is an effective bridge in the delivery of services and, as such, helps increase access to specialists in remote areas. Regulatory, legal, and reimbursement hurdles have prevented “telehealth” from being widely adopted, despite the benefits it has brought to both patients and clinicians (Gajarawala & Pelkowski, 2021). Studies which focus on HC equity also reveal that introducing virtual/hybrid care as an option makes joining the local community easier, and more people can benefit from the programs. For example, elderly people or individuals having some type of impairment as well as those who don’t have a means of transport or a local clinic, and so on. “Telemedicine” provides a safe, accessible, and convenient medical care (Jnr, 2020).

Moreover, the affordability and accessibility results that are observed here are deeply connected with the live discussion about the evolution of environmentally friendly HC systems. Besides that, “Telehealth” which is basically aimed at making patients' lives more comfortable by cutting down their expenses, can also be a great factor in achieving cost reductions at the system level. For example, by reducing hospital readmissions and alleviating unnecessary emergency visits. Using “telemedicine” for chronic illness management and preventative care programs can help both patients and providers save a lot of money. Medical professionals have been able to help patients of all ages and with a wide range of ailments by utilising remote health services (Almathami et al., 2020). In the same way, “telemedicine” treatments have made it easier for patients to get medical services and, at the same time, have lowered the total cost of treatment and the number of visits to hospitals and emergency rooms. These additional benefits of “telemedicine” can guide the decision of policymakers, insurers, and health care providers, especially in the current scenario of disrupted access to health care (J. S. Lee et al., 2024).

Hence, given these advantages and prior experiences, the findings highlight that prior telehealth exposure plays a critical role in shaping future usage intentions, emphasising the need for health systems to promote introductory telehealth experiences, patient onboarding support, and ongoing awareness initiatives. Such measures can help build long-term digital health engagement and enhance the sustainability of telehealth adoption.

5.5 Discussion of Research Question Four

In this study, it has been found that the general attitude of a user towards the quality of telehealth care is like and in the majority of cases equivalent to face-to-face care. They indicated that telehealth visits aided in solving their health problems, were able to receive adequate medical guidance and overall communication with the provider was very good.

These impressions contribute to achieving the growing acceptance of telehealth as an effective and reliable way of care delivery.

The statistical analysis also indicated that there is a strong positive correlation between the perceived quality of care and general user satisfaction. The respondents who rated the quality of telehealth positively were also more likely to experience greater levels of satisfaction with their telehealth. This implies that perception of quality is one of the determinants of satisfaction, which confirms past models of service quality in healthcare, claiming that effective communication, clear diagnosis and attention of the provider to the patient determine patient satisfaction.

Interestingly, users tended to stress that even though some aspects of physical examination may be more appropriate to be performed during the in-person visit, telehealth counters with the convenience, less waiting time, and more accessibility. These non-clinical quality dimensions, including responsiveness, scheduling ease, and comfort, had an impact on the development of user satisfaction and helped to form the idea that telehealth care quality is not lower than receiving a physical consultation.

Following the trends in the global literature, the findings reveal that a high level of perceived quality in telehealth correlates directly with the increase in the level of satisfaction, which justifies telehealth as a valid and reliable continuation of all modern healthcare services. This highlights the need of healthcare providers to uphold the standard of communication, technical dependability, and quality virtual interaction to uphold the level of user trust and satisfaction.

5.6 Discussion of Research Question Five

The future of “Telehealth” is bright with several technologies and care-delivery domains. The advancements in remote monitoring, wearable sensors, AI for triage and decision support, and improved video/virtual care platforms provide the means to transition

“telehealth” from the usual episodic visits to continuous care models, which enable chronic disease management and prevention. Essentially, “telemedicine” through the use of online HC delivery systems cuts back HC expenses and encourages a prompt execution of tasks in the system, eventually leading to the gaining of health equity and the rise of good health conditions (Anawade et al., 2024). The need for a patient to come to a HC facility is thus drastically reduced, and the patients can be given the care that they need at any time. If these innovations are also combined with predictive analytics that signal deterioration, patient convenience would be greatly enhanced, and intervention could take place at an early stage (S. Hu et al., 2025). But a precondition for these “telehealth” advancements to be possible is using clinical pathways as a medium for these tools rather than somehow incorporating them as an add-on; otherwise, the next-generation technologies will be underused.

One of the largest potentials is remote patient monitoring (RPM) and its ability to move care out of hospitals and into neighbourhoods, thereby possibly reducing expenses and raising the quality of care for chronic conditions (Dalloul et al., 2023). A patient who is monitored remotely can check in with a doctor online, and thus, the patient will be physically present only for a rare visit, and the usual medical care can be done virtually. As a result, RPM is potentially a very effective tool for hybrid care, which, on the one hand, will help to avoid unnecessary hospitalisations and, on the other, facilitate value-based care agreements. Policy makers should grasp the fact that strong governance structures, continuous evaluations and inclusive engagement are among the crucial factors that make sustainability and integration of “telemedicine” possible (Keelara et al., 2025). In a way, these changes would reduce the pressure on the health care system and make the larger input of experts more efficient and scalable; however, they would require investment in the procurement of devices, data transfer, setting up the payment models that recognise

monitoring services, and creating the HC personnel's workflows that turn constant data into meaningful actions.

Notwithstanding such outlooks, there are still significant hurdles that have the power to impact the extent of “telehealth” usage. The uncertainty over policy and payment — e.g., the varying reimbursement rules following pandemic waivers — is still the most important factor that could limit provider investment and long-term planning and thus be a serious bottleneck. FQHCs “federally qualified health centres” may see improvements in workflow, compliance, and no-show rates with the use of “telehealth”. Payment parity and more grants to solve “telehealth” infrastructural needs (such as Internet access, equipment, and literacy) are urgently needed, nonetheless, in order to reform current funding systems (Porteny et al., 2025). Besides regulatory questions (licensure, cross-jurisdiction care), medico-legal exposure, and workforce readiness (training, role redesign) also contribute to the friction that hampers the rate of adoption and the emergence of fragmented services (Shaver, 2022). The conclusion is that unless payers, regulators, and health systems coordinate incentives and clarify rules, many promising “telehealth” models will face difficulties in going beyond the pilot stage and sustaining operations over time.

Moreover, the combined impacts at the system, provider, and patient level rely on the factors of equity, interoperability, and trust. If investment decisions are made with the priority of ensuring access for all, securing data, and enabling smooth EHR integration, “telehealth” would be a tool that could facilitate access and lower the costs for many groups of patients; otherwise, it would contribute to the deepening of disparities and the emergence of siloed care (Association, 2025). Developing thorough training and capacity-building programs for HC providers is necessary to overcome medical professionals' opposition and guarantee the efficient utilisation of “telehealth” technologies (Ndwabe et al., 2024). Therefore, the reforms in policy (reimbursement that is stable), technical standards

(privacy, interoperability), workforce training, and the creation of targeted programs for the promotion of equity are the levers that most probably will determine whether or not the future benefits of “telehealth” will be realised at scale.

5.7 Strategic Analysis Using Business Strategy Frameworks

The findings of this study align strongly with core components of SWOT analysis:

Table 5.1: SWOT Analysis

Strengths (S)	Weaknesses (W)
High patient satisfaction and positive perceived quality	Technical difficulties, device issues, poor connectivity
Convenience, reduced travel, improved accessibility	Limited physical examination and interpersonal interaction
High willingness for future use among experienced users	Unequal digital literacy and access gaps among elderly and low-income groups
Opportunities (O)	Threats (T)
Integration with AI, RPM, hybrid care	Uncertain reimbursement and inconsistent regulation
Expansion into rural outreach and chronic care	Data privacy risks and medico-legal liability
Value-based care and system cost reduction	Fragmentation due to varying platform standards

Rather than assisting in descriptive assessment and aiding in strategic decision-making, the outcomes of the given research were analysed through the prism of a broader SWOT-to-TOWS strategic framework that enables integrating the results and dynamically interpreting them. This methodology is a more systematic analysis of the interaction between internal organisational strength and external environmental influence to derive tactical, situation-specific strategic options to telehealth adoption and scale-up, as opposed to viewing SWOT as a fixed checklist comprising strengths, weaknesses, opportunities,

and threats. The TOWS-based analysis can be used to transform empirical insights into the strategic oriented decisions to help in making informed managerial decisions on service positioning, resource allocation, risk reduction of risks and long-term sustainability. This model can be particularly relevant under the telehealth environment, where there is technological readiness, regulatory uncertainty, and market rivalry that cut across user acceptance in order to establish the consequences of adoption. The augmented SWOT-to-TOWS analysis, in its turn, provides a systematised framework in which the user can discover the ways in which the healthcare organisations may utilise the existing strengths to capitalise on the newly exposed opportunities, address the internal limitations, cope with the external threats, and incorporate telehealth into the stable, sensible, and purposeful care delivery models.

Strategic Interpretation of SWOT Dimensions

The internal strengths that the healthcare organization could leverage to achieve strategic advantage include high patient satisfaction, positive perceived quality, convenience and a high desire to continue using the product, which are the strengths that were witnessed in the current study. These strengths are good indicators that telehealth has attained a certain level of user legitimacy and trust thereby minimising behavioural-level resistance and adoption factors. This form of acceptance will provide a stable platform through which institutionalisation of telehealth can be done within normal care pathways, service differentiation, patient retention, and better continuity of care. All these advantages are bundled in a strategic plan that opens an opportunity of telehealth emerging as a support tool but a major ingredient in the paradigms of patient-centric and value-based healthcare services delivery.

The identified weaknesses (in their turn) that implied the lack of technical stability, the lack of physical examination scopes, and the differences in digital literacy indicates the

presence of internal factors that constrain the universal and unilateral use of telehealth services. These constraints have illuminated the necessity to add the element of telehealth with hybrid service design so as to have a balance of care through virtual and face-to-face to ensure a feeling of clinical relevancy and quality of service. Among the opportunities in the external environment, there is the introduction of artificial intelligence, remote patient monitoring (RPM), rural outreach, and compatibility as value-based care models, which offer colossal growth, scale, and innovation potential. Nevertheless, all these opportunities are accompanied by structural threats, such as uncertainty of reimbursement, a patchy regulatory landscape, data privacy, and platform compatibility, among others, that would weaken long-term sustainability unless they are addressed through governance, investment, and strategic coordination.

TOWS-Based Strategic Options for Telehealth

Building on the SWOT findings, a TOWS analysis enables the formulation of concrete strategic options:

- **SO (Strength–Opportunity) Strategies:** Healthcare organisations can use high patient satisfaction and usability to build on telehealth as a chronic disease management strategy, preventive care, and rural outreach strategy. By combining telehealth and AI-supported decision support and RPM, organisations can transition to more continuous and value-based care and enhance differentiation.
- **ST (Strength–Threat) Strategies:** Threats to telehealth services associated with regulatory uncertainty and privacy issues can be addressed through strong user trust and satisfaction by placing telehealth services in the context of well-developed governance, compliance, and security systems. Brand credibility

and institutional trust can be employed by established providers to lead competitive advantage in the increasingly regulated digital health markets.

- **WO (Weakness–Opportunity) Strategies:** The problem of technical constraints and digital illiteracy may be resolved by means of specific investments into infrastructural development, user education, and inclusive design with the assistance of collaborating with technological companies and government agencies. These plans enable organisations to turn their weaknesses into opportunities to build capability.
- **WT (Weakness–Threat) Strategies:** Defensive strategies are necessitated in the regions where internal weaknesses along with external threats exist simultaneously, i.e., low digital preparedness and unstable reimbursement. These involve a step-wise approach, a pilot-scaled approach, and the selective application of telehealth to clinically suitable cases in order to prevent operational and financial risk.

5.8 Feasibility Evaluation Using TELOS Models

To assess whether telehealth adoption is practical, sustainable, and strategically aligned with healthcare needs, the TELOS feasibility framework was applied. This model evaluates the technical, economic, legal, operational, and scheduling viability of telehealth based on the study’s empirical findings.

Table 5.2: TELOS Feasibility Evaluation

TELOS Component	Study Findings & Implications	Feasibility Rating
Technical	Connectivity issues, platform glitches, need for integration	Moderate
Economic	Cost savings for patients & system; uncertain reimbursement	Moderate

Legal	Data privacy, licensing, and medico-legal concerns	Low–Moderate
Operational	Good patient satisfaction; variable provider readiness	Moderate–High
Schedule	Rapid deployment is possible but dependent on policy timelines	Moderate

In order to further develop the feasibility analysis, the TELOS framework was used not as a single checklist but rather as a complementary strategic feasibility tool, as the technical, economical, legal, operational, and scheduling aspects of adoption affect each other mutually. The analysis is not conducted based on individual dimensions, but on the dynamic interplay of the individual dimensions- such as how technical infrastructure constraints can increase operational inefficiencies, how regulatory uncertainty can decrease economic feasibility, or how readiness of the workforce can influence the timeframes of implementation. The combination of TELOS allows a more realistic evaluation of the adoption of telehealth since systemic risks, organisational readiness and implementation constraints are captured, facilitating informed managerial decision-making on the prioritisation of investments, the implementation phases, and scalability of the telehealth solutions in healthcare organisations over time.

Technical Feasibility

Though the research establishes that telehealth platforms are largely functional and acceptable to the users, lack of consistency in connectivity, and device variability, and lack of interoperability with the current health information systems remain a problem to diminish overall reliability. Strategically, technical feasibility will be termed as the availability of the platform, besides the resiliency, capacity, and soundness of the underlying digital infrastructure. Telehealth implementation should be well integrated with electronic health record (EHR), clinical processes, and other fundamental healthcare

systems to make sure that virtual care is integrated fully into clinical operations and not an isolated service. In addition, the technical aspect will include the elements of cybersecurity and data integrity and standardisation of the platform as the key elements that should be established to build clinical and patient trust. The above systemic technical challenges will need to be overcome in order to make telehealth viewable as an addition or even experimental and, thus, unable to alter the care delivery paradigm, maintain hybridisations, and influence healthcare organisations in the long run.

Economic Feasibility

The economic feasibility can be justified by the fact that telehealth results in a significant change of patients in terms of saving money due to travelling, reducing opportunity costs, and the need to visit a hospital, and enhancing health care systems in terms of resource optimisation, hospital readmission, and care coordination. Nonetheless, telehealth has limited long-term financial viability due to the lack of consistent reimbursement frameworks, payer insecurity, and inconsistent coverage policies, all due to an unstable economic background. Strategically, this is a paradox: as economically beneficial as the concept of telehealth is, its implementation is prone to external financial and policy changes. In order to realise sustainable deployment, healthcare organisations need to combine operational cost reduction with high-quality revenue collection through the mechanisms of a value-based payment model, bundled care, and incentives based on performance. In that way, telehealth can turn into a cost-saving addition to the modern healthcare delivery and become a more affordable and strategically incorporated part of it.

Legal and Regulatory Feasibility

The most difficult dimension of TELOS is legal feasibility, which indicates the difficult and dynamic nature of the regulatory environment of telehealth. The main limitations are data privacy obligations, medico-legal liability, limitations on the

professional licence, and cross-jurisdictional regulations that introduce significant uncertainty and add administrative overhead for healthcare organisations. Notably, these legal and regulatory variables are very interdependent with economic and operational viability; unless there are clear guidelines and standards that need to be enforced, organisations would hardly invest heavily in infrastructure, workforce training, and integration of the platform. Strategically, it is not a side thing but a fundamental prerequisite of a sustainable scale that the legal barriers should be addressed, proactive governance reform, alignment of licensing and reimbursement policies, and standardised protocols of data security and patient consent. Well-developed legal and regulatory frameworks, therefore, constitute the basis of telehealth adoption that allows for minimising risks, establishing trust between stakeholders, and contributing to the overall viability of virtual care services in terms of their operations and financial sustainability.

Operational Feasibility

The telehealth operational feasibility is relatively high as the patient satisfaction levels are high, the user experience is positive, and the platform usability has been demonstrated, suggesting that virtual care can be used in the current service delivery. Nonetheless, the readiness of the providers is not balanced, and there are significant disparities in the personnel training, workflow redesign, role adjustment, and clinical confidence in virtual modalities. Strategically and management-wise, operational feasibility is not just about the implementation of technology, but it depends highly on the ability of an organisation to change, the willingness of the leadership and organizational culture of institutional learning. Health facilities that take the initiative to invest in workforce development, develop definite telehealth models, and create flexible and participatory conditions will be at a better position to incorporate virtual services into regular practices. On the flipside, those organisations that have poor change management

practices face the risk of underutilization, poor quality, and employee resistance, and as it is, the success of telehealth implementation depends not only on technological capability but also on organisational preparedness and strategic governance.

Scheduling and Implementation Feasibility

Telehealth implementation and schedule have been characterised as moderate regarding their complexity since they demand organisation of adoption at both organisational and external levels. Though the technical part of telehealth solution implementation may be rather short in duration, integration at the enterprise-wide levels requires a considerable compromise with regulatory approvals, budget cycles, human resource accessibility, and the work processes. The pace of adoption and its effectiveness is all contingent on policy schedules and reimbursement options and the preparedness of the institution; therefore, fragmented rollouts are risky and disruptive. This is strategically important as it emphasises the significance of implementation schemes that are performed in phases, in a step-by-step manner and are dynamic in nature, where pilot programs are grown over time, lessons learned in the process and optimisation of resource utilisation are made. By ensuring equal technical preparedness to organisational capability, adherence to regulations and financial planning, healthcare organisations stand chances of successfully adopting telehealth as sustainable and efficient and resilient.

Integrated Strategic Feasibility Interpretation

The extended TELOS analysis shows that telehealth may be regarded as a strategically viable option, but its sustainability at the long-term level depends on the possibility of addressing the issues associated with the problem on a multidimensional level. Technical strength and operational fitness form the strong basis, which implies that the platforms are accessible, and the experiences with patients are positive in general. However, the fact that the economy is unpredictable with changing reimbursement

systems, legal and regulatory uncertainty are significant growth and long-term investment constraint. The interrelationship between TELOS dimensions highlights that piecemeal development in one dimension, such as the modernisation of technology, without centralising the unemployment facilitation and compliance with legal requirements, will barely deliver profitable outcomes. Thus, healthcare organisations need to seek systematic and coordinated interventions, including governance changes, uniform reimbursement models, the development of digital infrastructure, employee education, and workflow redesign. Such an integrated method is the only way that telehealth adoption can transform into more than a technically feasible pilot project and a strategically integrated, scalable, and resilient element of a contemporary healthcare delivery system that can produce a long-term value to patients, their providers, and the health systems alike.

5.9 Financial Feasibility Analysis of Telehealth Implementation

The adoption of telehealth should be considered in its strategic viability calculated not only in terms of user satisfaction and intention to use but also in terms of financial utilisation. To ensure the sustainability of telehealth services in healthcare institutions, a systematic examination of the return on investment (ROI), the cost/benefit proportion, and the break-even timelines is necessary.

1. Cost Structure Contemplations

The main elements of costs of telehealth implementation are:

- Primary technology infrastructure (software platforms, secure communication systems, servers)
- Hardware (cameras, surveillance equipment, digital diagnostic equipment)
- Employee empowerment and online ability.
- Data protection compliance and cybersecurity.
- Maintenance and subscription licensing fees.

These are more about fixed upfront investments, with moderate recurring operating costs thereafter.

Telehealth, however also saves some of the traditional healthcare expenses, including:

- The use of physical infrastructure.
- Overheads of administration associated with face-to-face visits.
- Patient no-show losses
- Travel and logistical coordination expenses.

In this way, telehealth will change the cost model in the facility-intensive spending to the delivery of services based on technology.

2. Cost–Benefit Analysis

A cost benefit analysis indicates that telehealth produces both direct and indirect financial benefits.

Direct Financial Benefits:

- More consultations because they were able to schedule.
- Lower overhead per patient encounter.
- Broadened market geographically.
- Reduced the marginal cost of extra consultation.

Indirect Financial Benefits:

- More patient retention (satisfaction results in Chapter 4)
- Improved brand positioning as a digital progressive provider.
- Hardiness in operations in times of interruptions (e.g., pandemics)
- Less expensive to acquire patients in the long-term.

Though Chapter 4 results show a statistically significant but weak relationship between satisfaction and reuse intention, small changes in repeat usage can have significant cumulative revenue implications over time as they increase.

Taking on the managerial side of the issue, the financial incentive is heightened more in case telehealth is introduced as a hybrid care model instead of being implemented as a single service.

3. Return on Investment (ROI)

Telehealth Return on Investment (ROI) may be described as follows:

$$\text{ROI} = (\text{Net Financial Gain} - \text{Total Investment Cost}) / \text{Total Investment Cost} \times 100$$

During the initial stage, ROI can be moderate because of the cost of infrastructure setups. Nevertheless, with the growing volume of consultation and the maturing process associated with digital processes, the marginal costs decrease, but the revenue per digital encounter is constant.

With empirically measured positive perception of patients and their readiness to continue using telehealth services, the possibility of protracted utilisation enhances the long-term ROI prospects.

ROI is maximised when:

- The utilisation of digital appointments is above a certain level.
- Automation of administrations decreases the redundancy of labour.
- The scalability of the platform enables it to increase without increasing costs accordingly.

Therefore, telehealth is not to be considered in the short-term quarterly performance but in the medium- to long-term ROI horizon (12-36 months).

4. Break-Even Analysis

Break-even analysis is used to identify the stage where the total income of the telehealth consultations is equal to the total expenditure.

The break-even volume may be determined as:

Break-Even Point = Fixed Costs/Contribution Margin per Consultation

In the case where the contribution margin is determined as:

Consultation Fee–Variable Cost per Digital Visit

Since telehealth minimises variable operational expenses when compared to face-to-face visits, the contribution margin per consultation can be comparatively high after the infrastructure is developed.

According to strategic modelling assumptions, in case the uptake of digital consultation has become a recurring stable level, and variable costs are checked by technological efficiency, the break-even level may be reached at the initial operation stage, especially in cities and semi-urban populations that are highly digital.

5. Strategic Financial Implications

The financial feasibility study shows telehealth as a patient convenience program and a service model that can expand strategically.

Financial sustainability, however, depends on:

- Sufficient patient adoption rates.
- Connection with already established healthcare processes.
- Constant check of performance indicators.
- Cybersecurity and regulatory compliance risk mitigation.

Telehealth shows its potential for sustainable long-term economic viability when researchers combine empirical satisfaction data with structured financial models and conduct monitoring during staged implementation.

5.10 Telehealth Readiness Assessment Tool (TRAT): A Practical Framework for Healthcare Organisations

Although the current analysis shows that telehealth solutions are viable and accepted by an increasing number of people, healthcare organisations do not have systematic tools to assess their internal preparedness to implement telehealth solutions. The professional healthcare leaders need to evaluate the technological feasibility and the organisational preparedness, the capability of the workforce, the alignment with the regulations, and financial sustainability. To fill this practical gap, the current study offers an original Telehealth Readiness Assessment Tool (TRAT) that will help to make an evidence-based decision in the practical healthcare environment.

The framework is made up of seven organisational dimensions, which are interrelated and measured on a five-point standardised rating scale of very low readiness (1) to very high readiness (5). A combination of these dimensions describes the technological, organisational, human, regulatory, and financial needs to have a successful deployment of telehealth.

Table 5.3: TRAT: Organisational Dimensions (7 Dimensions)

Dimension	Description	Rating Scale (1–5)
1. Technological Infrastructure Readiness	Internet reliability, hardware availability, EHR interoperability, cybersecurity	1 = Very Poor → 5 = Excellent
2. Leadership & Strategic Commitment	Senior management support, vision alignment, funding commitment	1 = No support → 5 = Strong commitment
3. Clinical Workflow Integration	Alignment of telehealth with existing clinical processes	1 = No integration → 5 = Fully integrated
4. Human Capital & Digital Skills	Staff digital literacy, training availability, change readiness	1 = Unprepared → 5 = Highly skilled

5. Patient Readiness & Acceptance	Patient digital access, willingness, trust in telehealth	1 = Low acceptance → 5 = High acceptance
6. Regulatory & Compliance Preparedness	Licensing, data privacy (HIPAA/GDPR), reimbursement clarity	1 = non-compliant → 5 = Fully compliant
7. Financial Sustainability	Budget allocation, ROI clarity, reimbursement mechanisms	1 = Unsustainable → 5 = Sustainable

Scoring Interpretation

- 7–17 → Low readiness (High implementation risk)
- 18–27 → Moderate readiness (Pilot recommended)
- 28–35 → High readiness (Scale-up feasible)

The cumulative TRAT score gives a clear picture of the level of telehealth readiness at an organisation. Low scores mean very high risk of implementation and the necessity to make improvements that would be laid in the foundation, whereas high scores indicate the possibility to scale the telehealth services. Using this tool, healthcare leaders will be able to perform internal diagnostics, compare readiness between departments, and work out individual implementation plans in accordance with the organisational capabilities.

Telehealth Implementation Success Factors Model

Telehealth readiness is a necessary condition of adoption, but it does not necessarily indicate success and a long-term implementation process. Numerous telehealth projects end up failing even following a proper investment in technology because of organisational challenges, behavioural forces of human beings, and the environment. To close this gap between adoption and long-term success, the current research study introduces a novel original Telehealth Implementation Success Factor Model (TISFM).

The TISFM is a practitioner-based framework that is systems-oriented and explains why certain telehealth projects are successful and others perform poorly or fail despite the

same technological environment. In this model, the basic findings of the study are incorporated with the literature on the implementation and adoption of research to offer healthcare leaders a comprehensive execution model.

The following Table 5.4 illustrates the key constructs of this Framework:

Table 5.4: Telehealth Implementation Success Factors Model (TISFM) Factors

Domain	Key Factors	Managerial Implications	Outcome Factors Influenced
Technological	Platform usability, system reliability, data security, interoperability	Ensure stable, secure, and user-friendly telehealth platforms	Clinical quality, cost efficiency, and operational reliability
Organizational	Leadership support, strategic alignment, process redesign	Embed telehealth into core strategy and workflows	Adoption sustainability, operational efficiency
Human	Clinician acceptance, patient trust, training, digital literacy	Invest in training and structured change management	Patient satisfaction, service continuity, quality of care
Environmental	Regulatory support, reimbursement policies, digital infrastructure	Monitor policy changes and ensure compliance	Financial viability, scalability, long-term sustainability

The model has four interactive domains as seen in the table above: technological, organisational, human, and environmental, all of which are paramount in shaping the outcome of telehealth. Instead of considering telehealth as an entirely technological intervention, the TISFM conceptualises the implementation success as a multi-level alignment between the systems, people, and policy environments.

Figure 5.2 illustrates the Telehealth Implementation Success Factors Model (TISFM).




Figure 5.2: Telehealth Implementation Success Factors Model (TISFM)

Source: (Self-Created)

5.11 Strategic Implications for Healthcare Organisations

Telehealth adoption has profound strategic implications for healthcare organisations, influencing **competitive positioning, market entry barriers, and strategic growth options**.

Competitive Positioning: Telehealth shifts the historical nature of the competition by transferring the differentiation base to the avenue of clinical expertise knowledge to the digital access, convenience of patients, and patient-to-patient care channels. Through innovation, organisations that succeed to incorporate telehealth in their service delivery

can be differentiated in a highly crowded market since they will introduce hybrid solutions that combine both face-to-face and remote care. Such positioning not only accelerates the retention of patients but also improves brand image, a card that is a sign of innovation and sensitivity to the changing patient needs. Telehealth also provides geographic expansion in which inaccessible or underserved populations could be reached by the provider, which expands the market more without a corresponding capital expenditure in the physical infrastructure.

Entry Barriers: Despite the fact that telehealth creates new opportunities, it creates a threat of barriers to entry strategy. The entry barriers are high with the large initial investment in the technologies, the compliance with the extremely complicated system of regulations and the use of the electronic health records and the training of the workforce. The organisations that have worked out a clinical and IT infrastructure are in close contact with payers, and have previous experience in negotiating regulatory mandates, have the first-mover advantage. Contrastingly, a smaller provider, or resource-constrained providers, have the potential to fail to scale or interoperate; it is possible to mention that strategic partnerships or outsourcing solutions matter.

Strategic Options: Healthcare organisations can adopt multiple strategic approaches to leverage telehealth:

- 1. Market Penetration:** Telehealth expansion in the new patient bases to enhance use and satisfaction.
- 2. Service Diversification:** The use of specialised remote services (e.g., chronic disease management, mental health, post-operative follow-up) as a means of attracting new patient groups.

- 3. Alliances and Partnerships:** Partnering with technology providers, insurance companies and remote monitoring devices manufacturers in order to improve service quality and scale.
- 4. Value-Based Care Integration:** Integrating telehealth into value-based payment models, outcomes, preventive care, and decreased hospital readmissions.

It is important to note that these options have to be determined strategically, in the framework of regulatory, financial, and operational feasibility, where telehealth adoption not only remains technically viable but also competitively beneficial and sustainable in the long run.

5.12 Business Model Innovation Opportunities

Telehealth adoption offers healthcare organisations the chance to reimagine traditional business models, moving from volume-driven to value-oriented and patient-centred models.

Revenue Models: Telehealth establishes new sources of revenue, including subscription plans of remote care, pay-per-consultation services, remote chronic illnesses monitoring, and schemes of bundled payments with potential reimbursement depending on the outcome. The other approach through which organisations can consider partnering with insurers is to ensure that telehealth is not excluded in the coverage to allow patients to pay lower out-of-pocket rather than providers of telehealth receiving unpredictable cash flows.

Value Propositions: Telehealth will create value for the patients and the healthcare organisations. The benefits of the patients would be convenience, time saving, continuity of care and increased accessibility of experts. Telehealth enables the providers to achieve cost reduction, operational efficiency and scalability to an extent that a provider can manage a larger number of patients without a corresponding growth in the physical

infrastructure. The strategic value of the patient-centric and data-driven care experience based on analytics, remote monitoring, and customised care plans development is its strategic value.

Partnerships and Ecosystem Development: The cross-sector partnerships will most likely prove efficient in the context of telehealth adoption. The partnership with the technology vendors, artificial intelligence developers, device creators, and cloud infrastructure companies could ensure the stability of the platform, its lack of interoperability and a steady flow of innovation. The access of underserved groups can be improved through cooperation with the community organisation and insurers, and the reimbursement systems can be improved.

Scalability and Expansion: Telehealth models are inherently scalable as well, and thus, healthcare organisations can reach an increased number of people without being restricted by their location. Hybrid models Telehealth and face-to-face care combinations, can assist the organisations in allocating clinicians effectively, reducing patient wait time, and utilising resources better. Having platforms based on modular technology, standardised protocols, and strong data governance is strategic because scalability is enhanced and organisational requirements are met and services can be scaled on demand and regional needs.

5.13 Change Management Framework for Telehealth Implementation

Ready Assessment: Before scaling telehealth, organisations need to assess their readiness in terms of technologies, operations, and human resources. Among others, the critical factors are the reliability of the infrastructure, the skill of clinicians in virtual care delivery, the integration of workflows, and the level of digital literacy of personnel and patients. Assessment tools, surveys, pilot programs can reveal the gaps, and then specific interventions could be implemented before the full deployment.

Stakeholder Analysis: Telehealth has various stakeholders, namely, patients, clinicians, administrators, IT staff, and payers. It is important to map the stakeholders and their interests and where they may be resisting. Using the example, clinicians can be afraid of workflow disruption, liability, or absence of interaction with the patient but patients can be afraid of privacy, digital illiteracy, or distrust in virtual care. These attitude insights assist the organisations to tailor the communication, training and incentives to ensure that the stakeholders are oriented towards the adoption goals.

Resistance Management: Resistance is natural during transformational initiatives. Strategies to mitigate it include:

1. **Communication and Education:** The advantages of telehealth, both to the patient and to the provider, should be well explained, as well as continuous training should be offered to increase confidence.
2. **Incentive Alignment:** tying the telehealth adoption to performance metrics, reimbursement incentives, and professional development.
3. **Incremental Implementation:** Telehealth services can be gradually rolled out in phases, starting with the low-risk, high-acceptance implementation cases so that staff and patients can adjust.
4. **Feedback Loops:** Ongoing monitoring, gathering feedback and repetitive advancements are used to make sure that operational, clinical, and technical obstacles are tackled in time, which supports trust and dedication.

Introducing a change management framework will also make the adoption of telehealth not just possible through technology, but it should also be sustainable on an organisational level, and digital care should be ingrained in the working culture and processes of the organisation.

Table 5.5: Summary of Feasibility & Strategy Findings

Domain	Study Insight	Strategic Implication
Technical Feasibility	Good platforms but inconsistent connectivity	Invest in infrastructure & integration
Economic Feasibility	Cost benefits proven	Need stable reimbursement
Legal Feasibility	Privacy & compliance barriers	Strengthen governance
Operational Feasibility	High user satisfaction	Expand hybrid care
Strategic Fit	Aligns with value-based care	Adopt telehealth in care pathways

The strategic analysis of the feasibility results indicates that telehealth would fit the current healthcare priorities; however, it needs special measures to be expanded sustainably. Digitally, interoperability and improvement of digital infrastructure is necessary in order to overcome the connectivity and platform disparities. On the economic front, telehealth helps to save money, but the stability of reimbursements is essential to guarantee the long-term involvement of the providers. There should be stricter governance structures at the legal level to deal with concerns of privacy, compliance and cross-border licensing. The level of operational feasibility is high because users are satisfied, which means that they are ready to further integrate hybrid care. Telehealth is a strategic fit within a value-based model of care and must be integrated into its standard care pathways to enhance efficiency and accessibility.

CHAPTER VI: SUMMARY, IMPLICATIONS, AND RECOMMENDATIONS

6.1 Summary

The primary aim of the research is to identify and analyse the diverse factors that have led to the successful acceptance of “telehealth” and its increasing importance as a revolutionary innovation in HC distribution. The initial consideration centred around the question of how the HC actors i.e. providers and patients can change their relations via the adoption of digital and communication technologies. This would result in better access to HC, continuous care, and higher productivity. This study aims to interpret “telehealth” in a wider sense, other than just the technology supporting it. It envisioned a system that could address some of the issues that have been hindering HC today, such as cost limitations, long distances to travel, and insufficient resources. Medical wellness. The study took the quantitative and qualitative approaches. This research gave a complete picture of Telehealth's current usage, the obstacles in its implementation, and its possible to change HC globally through the findings of the study.

The results of the research indicate that “telehealth” has evolved significantly beyond the expectations set initially and is now considered as a main HC practice rather than a temporary one or secondary service. It has even been able to provide HC to the remote and low-income communities, to enhance patient satisfaction, and to make better use of clinical hours. Both the survey results and the literature have shown that “telehealth” is helpful in the treatment of chronic diseases, it cuts down the time from the doctor's appointment to the patient, and it relieves the pressure on hospitals and clinics. Moreover, the limited scope for “telemedicine” to grow was indicated by the study as the major factors that hold it back are unequal internet connectivity, lack of trained medical staff, and different regulations in different regions. These difficulties show that the road is still very

long to travel and the partnership between the government, health care providers, and the technology sector is paramount if we have to deliver “telehealth” services in a way that is both equitable and efficient.

The studies have identified the reality that the perseverance and success of Telemedicine depends on the combination of the following factors: technology, easy to use, and institutional preparedness. The system under consideration should be supported at all times by investing in physical and technological facilities, training the population, and digital literacy to establish the needed level of confidence between users and providers in order to implement a fully working system of telehealth, as the latter can be a variation of the traditional model of HC delivery. The transition is not only technological but also carries the connotation of the HC system being more inclusive, more accessible, and more patient-centred, as the transition is in line with the changing demands of the contemporary society.

6.2 Telehealth Performance Measurement Framework

Although telehealth adoption and implementation can be considered essential, these measures will only guarantee success in the long-term upon constant performance monitoring and evaluation of outcomes. Telehealth service providers, including healthcare leaders, need developed, data-oriented instruments to monitor clinical outcomes, patient experience, financial sustainability, and efficiency. To address this need, the present study proposes an original Telehealth Performance Dashboard Template (TPDT) as a practical performance management framework.

The following table shows the key constructs and KPIs of the dashboard of measurement for telehealth performance:

Table 6.1: Telehealth Performance Dashboard KPIs

Performance Dimension	Key Performance Indicator (KPI)	Description	Measurement Frequency
Clinical Quality	Diagnostic accuracy rate (%)	Percentage of correct diagnoses delivered via teleconsultations	Monthly
	Follow-up compliance rate	Proportion of patients completing recommended follow-ups	Monthly
	Clinical escalation rate	Percentage of telehealth cases escalated to in-person care	Monthly
	Readmission rate (telehealth vs in-person)	Comparative readmission outcomes between telehealth and physical visits	Quarterly
Patient Satisfaction	Telehealth satisfaction score	Average patient satisfaction rating for teleconsultations	Monthly
	Net Promoter Score (NPS)	Patient likelihood to recommend telehealth services	Quarterly
	Average wait time	Time from appointment booking to consultation	Monthly
	Consultation completion rate	Percentage of successfully completed teleconsultations	Monthly
Financial Performance	Cost per consultation	Average operational cost incurred per teleconsultation	Monthly
	Revenue per teleconsultation	Average revenue generated per teleconsultation	Monthly
	Reimbursement realization rate	Percentage of claims successfully reimbursed	Quarterly
	ROI of telehealth program	Return on investment from telehealth initiatives	Annually
Operational Efficiency	Appointment no-show rate	Percentage of missed telehealth appointments	Monthly

	Provider utilization rate	Proportion of clinician time effectively utilized	Monthly
	Average consultation duration	Mean duration of teleconsultations	Monthly
	Platform downtime (%)	Percentage of system unavailability	Monthly

Based on these, the following Figure 6.1 presents the Telehealth Performance Dashboard Template (TPDT).




Figure 6.1: Telehealth performance dashboard

Source: (Self- Created)

6.3 Implications

The results of this study should have serious implications for leaders, hospitals, the creators of technology, and the rest of the human race. The transition to “telehealth” among the patients is a sign that health care is no longer the same; it is very much different and is simply going through a digital transformation. A regulatory framework is one of the key

messages that this research will pass to the top management, that the framework should be the first and most urgent thing that the management does. such rules should consider different parts of the situation, for example, how the licenses are issued, how the services are paid for, how the information is protected, and the confidentiality of the patient. The enactment of policies of this nature will very much contribute to the integration of “telemedicine” into the universal health care system and thus bring patients' communication with doctors closer, making it easier for everyone to access the service, and ensuring health care continuity. A properly monitored telehealth setting will surely become a source of confidence among patients and practitioners, attract investments of both government and the private, and open the entire health care sector to the digital health solutions.

Judging by the evidence that the organisation offers and the clinic itself, the HC professionals are not able to become spectators of the new digital world but have to enter it. To ensure the health workers receive adequate training and encouragement on how to use technology, hospitals, clinics and a doctor in particular must embrace the concept of telehealth as a routine in treating patients. The main focus should be on upholding virtual consultations to the highest standard, ensuring patient safety, and maintaining good communication between doctors and their patients. In addition, medical executives need to consider obstacles, for example, a lack of digital skills, inadequate facilities, and an unwillingness to accept technological changes. Increasing the supply of “telehealth” services at the hospital will not only make the flow of activities more efficient but also increase patient delight and better health results in various groups of people.

On a societal and technological level, the consequences go as far as inclusivity and accessibility issues. According to the study, it is very important to make the digital divide by increasing the areas with internet connection, lowering the prices of digital devices, and

guaranteeing that the “telehealth” platforms are easy to use and any culture can use them without problems. The developers and innovators of technology ought to direct their efforts towards the creation of mutually compatible systems that not only harmonise with the presently available electronic health records but are also in agreement with the data protection norms. From a social perspective, the use of “telehealth” is a great way to level the playing field in access to HC and, in particular, can be of immense help to those living in rural areas, the aged, and people with some kind of physical limitation. If the HC sector were to address these areas together, it would not be able to use “telehealth” just as a convenient way, but rather as a highly effective tool for achieving health equity, saving resources, and even being resistant to future global health problems.

6.4 Strategy Recommendations-Stakeholders.

The results of the empirical study indicate that the perception of patients, their satisfaction levels, which are statistically significant, and moderate behavioural intentions to reuse telehealth services are positive. Nevertheless, both in the short and long term, sustainable development is based on concerted effort in leadership, finance, technology, entrepreneurship, payer systems, and regulatory frameworks. Consequently, the recommendations have been designed to provide importance of stakeholder roles so as to be clear in terms of strategic responsibility and practical applicability.

Recommendations for Hospital CEOs

The adoption of telehealth should be a strategic change at the enterprise level instead of a technological project of the Hospital Chief Executive Officer (CEO). Through the study results, it can be recommended to the CEOs to formally implement telehealth into the core clinical and operational strategy of the hospital within a short-to-medium period, namely by hybrid care models integrating virtual and in-person services to follow-up care and chronic disease management, mental health services, and rural outreach. The key

element is leadership commitment, so CEOs are advised to develop executive governance processes, invest in clinician education, and align organizational culture, workflow and performance incentives to minimize resistance and to be prepared to change. Competitively, telehealth must be capitalized upon to further position in the market, increase ability to reach more patients than ever before due to geographical limitations, and enhance patient retention by making services more accessible and convenient. Lastly, CEOs should institutionalize the constant performance tracking with the help of the key indicators of patient satisfaction, utilization rates, clinical outcomes, and the efficiency of operations to sustain telehealth initiatives, make them compliant, and in accordance with the long-term organizational objectives.

1. Strategic Actions and Leadership Priorities

The hospital leadership is decisive in making sure that telehealth initiatives can become sustainable organizational value and not a collection of technological experiments. Strategically, the telehealth should be incorporated into the long-term vision and model of service delivery in the hospital, and the senior management should become the strong promoter of its implementation in clinical, administrative, and technological spheres. A good leader must be able to put in place good governance mechanisms, align telehealth objectives with institutional performance indicators and promote cross-functional interaction to reduce silos. Besides that, the leaders are supposed to overcome change resistance by inculcating a culture of learning, invest in the development of clinician capabilities and ensure that the telehealth initiatives are adequately staffed and supported. Hospital management can turn telehealth into a basic quality, access, and competition facilitator in healthcare provision with the assistance of frequent executive management and strategic alignment.

Key Strategic Actions and Leadership Priorities:

- **Integrate telehealth into the hospital's long-term strategic plan and core service portfolio:** It includes the creation of a permanent role for telehealth in healthcare delivery rather than a temporary one. Hospitals can ensure that virtual services align with organisational objectives, such as patient-centred care, service growth, and efficiency, when telehealth is incorporated into strategic planning.
- **Establish executive-level governance and accountability structures for telehealth initiatives:** Some of the governance structures, such as steering committees or executive sponsors, should be used to give ownership, decision-making power and accountability. This helps to coordinate the work of the departments and enables to resolve the organisational and regulatory problems within the required time.
- **Promote cross-functional coordination among clinical, IT, legal, and administrative teams:** Telehealth implementation engages different sections of the organisation. Effective coordination will ensure that the clinical process, technology platform, legal mandates, and administrative functions are coordinated to reduce fragmentation and maximise service delivery.
- **Invest in leadership and clinician training to enhance digital readiness and acceptance:** Resistance is commonly caused by poor digital abilities or being uncertain about the practices of virtual care. Confidence, adoption encouragement and high-quality delivery can be enhanced through structured training and leadership development programs.
- **Align telehealth performance indicators with organizational quality and efficiency goals:** The tracking of the indicators of patient satisfaction,

utilization rates, clinical outcomes, and cost efficiency allows the leadership to assess performance and implement evidence-based changes. Congruency with organizational objectives can be used to make sure that telehealth is an important part of institutional success.

2. Implementation Timeline and Strategic Outcomes

The successful implementation of telehealth should be a gradual and time-limited process that identifies the strategic intent and operational implementation. Strategic alignment and organizational readiness should be considered by the leadership of the hospital in the short run (56 months). The main steps in this stage are to officially introduce telehealth as the strategic direction of the hospital, to create executive governance and responsibility frameworks, to identify the primary areas of service that should be offered through virtual means, and to establish a program of clinician and staff training. The desired results of the stage are the increase in alignment of leadership, decreased change resistance and developing an institutional roadmap of integrating telehealth.

The middle-term (6-12 months) should focus on the operational integration and performance optimization. The hospitals are advised to combine the use of telehealth with the existing electronic health records, reorganize the clinical workflow to facilitate hybrid care delivery systems, and extend telehealth services to specific patient groups like populations with chronic illnesses and rural or underserved areas. At this stage, the management is supposed to initiate systematic or planned monitoring of performance based on key indicators in terms of utilization, patient satisfaction, and efficiency of operations. The expected results are a higher adoption rate, better continuity of care, and quantifiable service access and cost-efficiency.

Over the long term (12-18 months), telehealth needs to shift up to an implementation initiative to a source of competitive advantage. The key strategic areas to

focus on during this stage are to scale the successful telehealth services, enhance partnerships with technology providers and payers, and align telehealth delivery with value-based care and reimbursement models. The processes of continuous improvement would be institutionalized to ensure that the changes of regulations and fluctuating demands of the patients are followed. The efficiency of the operational work, improved market positioning, expanded coverage of the range of services, and transforming telehealth into a stable and value-generating component of the model of hospital delivery are desired outcomes of this step.

Recommendations for Chief Financial Officers (CFOs)

The Chief Financial Officers has the responsibility of ensuring the financial sustainability and necessary viability of telehealth programs using a systematic, data-driven finance solution. The CFOs are encouraged to develop comprehensive financial evaluation models that have potential to include cost benefit analysis, ROI measurement and break-even analysis in order to justify telehealth investments that would not be likely to generate instant revenue. Particular attention should be paid to the calculation of not only direct financial results, such as an increase in volumes of served and a decrease in overheads, but also indirect value creation, such as an increase in patient retention, a reduction in readmission rates, and operation efficiency. In addition to this, CFOs are supposed to assume the risk of managing financial risks as a result of uncertainties in reimbursements, regulatory reactions and costs of maintaining technology with scenarios planning and sensitivity analysis. Revenue stability will also be achieved by investing in telehealth based on the value-based care model and negotiating sustainable reimbursement plans with payers. This is attainable by CFOs through a continuous monitoring of financial performance indicators and aligning telehealth spending to organizational strategy, which

will ensure that telehealth adoption is contributing value to the overall financial resilience and competitive advantage.

1. Financial Evaluation and ROI Framework

In the case of CFOs, telehealth strategic management must be thoroughly planned and analyzed financially to be a long-lasting investment. Short-term, CFOs are to conduct detailed cost-benefit analyses of telehealth infrastructure, technology expenses, and staffing expenses against the expected returns in the operations at the hospitals, including fewer patient admissions, fewer missed appointments, and less utilization of the emergency room. The investment appraisal must focus on those projects that have maximum potential of scaling and those that fit in the organization strategy. The computation of ROI should also include direct and indirect benefits that encompass patient satisfaction and better clinical outcomes which may increase the reputation of the hospital and subsequent revenue. Break-even should determine the break-even point at which telehealth investments will pay off their initial costs and financial risk analysis should take into consideration uncertainties of the dynamics of the reimbursement rates, cost of maintaining technology and probable regulatory compliance costs. All these actions are aimed at helping CFOs have a proper financial roadmap that justifies and manages investments in telehealth.

Cost-Benefit Analysis: It involves the balancing of all the costs incurred in implementation of technology in the telehealth (technology, training, maintenance of the platform, staffing, etc.) to the consequences. A reduction in patient readmission, a dwindling cost of operation, and enhanced clinical efficiency are some of the benefits. This will be to ensure that all investments are contributed to the overall organization performance.

Investment Appraisal: CFOs must concentrate on those projects that can be scaled and can be aligned with the hospital strategy. With this in mind, it may turn out to be more profitable to conduct telehealth with large amounts of chronic care patients than pilot programs. Appraisal entails consideration of financial viability, operational impact as well as long- term strategic fit.

Calculations of ROI: ROI should not just count on direct revenue increases but should also count on the indirect revenue of the increased patient satisfaction, retention and reputation of the hospital. These are not so concrete but can increase revenue streams in future and can justify the investment.

Break-Even Analysis: Determines the at which the total revenue or cost savings of telehealth is equal to the initial investment and operation costs. This will enable CFOs to know how much time will it take before telehealth can be made financially viable.

Financial Risk Assessment: Evaluates uncertainty in occurrence of variable reimbursement rates, cost of technology maintenance or unforeseen compliance cost. CFOs are able to predict these risks and thus make plans in the event of contingencies, which will minimize the chances of financial losses.

2. Revenue Sustainability and Reimbursement Strategy

CFOs need to make sure that the streams of telehealth revenues are predictable and sustainable. This would include solving the problem of reimbursement uncertainty by simulating different possibilities, such as different payer policies, low approvals, or temporary pandemic-related allowances. Scenario planning assists hospitals to know the cost of various reimbursement situations and deal with contingency plans. It is important to align with value-based payment models, attaching the outcomes of telehealth to the quality indicators and performance rewards instead of just sticking to the fee-for-service. A long-term sustainability also implies the active work with payers negotiating the

conditions of reimbursement of remote consultations, remote monitoring, and hybrid care programs. Integrating preemptive scenario planning and alignment with value-based models can help the CFOs to guarantee a stable revenue stream and encourage the effective and quality delivery of care.

Reimbursement Scenario Planning: As telehealth reimbursement policies can differ by payer, location or type of service, CFOs have to plan various financial scenarios. This will assist in forecasting the possible changes in revenues and drawing up mitigation measures. Indicatively, what would be the situation where some of the remote monitoring services are no longer paid at the prevailing rates?

Value-Based Payment Alignment: Telehealth revenue should be based on value-based models that use quality and outcomes to reward hospitals other than merely using a fee-for-service model. The correlation of telehealth services to the outcomes of patients guarantees sustainability of financial encouragement in the long-term and an effective care provision.

Payer Engagement: CFOs are supposed to actively negotiate insurance deals with companies to cover telehealth services, such as virtual consultations, remote monitoring of patients, and hybrid care. Active involvement will help remove revenue risk and make adoption rates higher.

Revenue Diversification: In addition to the conventional reimbursement, CFOs have an opportunity to consider other venues of revenue, i.e., subscription-based chronic care packages, bundles, or collaborating with digital health providers. Revenue diversification will decrease their reliance on one payer systems and increase financial sustainability.

Recommendations for Chief Information Officers (CIOs)

Chief Information Officers play a crucial role in the successful implementation as well as scalability of telehealth systems because they play a role in ensuring that digital infrastructure is aligned with clinical and strategic goals. CIOs are expected to look at building a solid, interoperable and secure technology ecosystem that is able to provide the seamless delivery of virtual care across the platforms and devices. This involves proper network capacity, integration of network with the existing hospital information systems and electronic health records, and compliance with data privacy and cybersecurity. Moreover, CIOs ought to have a gradual technological adoption strategy that would enable them to test, optimise and scale down without causing any significant nuisance to operations. It is important to constantly monitor the performance of systems, user experience, and technological risks in order to ensure the quality of services and the trust of clinicians. CIOs can ensure a long-term viability of telehealth operations by focusing on interoperability, data security, and future-ready digital architecture, which can improve the continuity of care and efficiency of operations within organizations and their competitiveness.

1. Technology Infrastructure and Integration Requirements

Telehealth services can only be effectively implemented with the help of a robust and highly integrated technology infrastructure, which is the primary task of the Chief Information Officer. The third issue that CIOs need to focus on is the choice of scalable and user-friendly telehealth platforms that would fit into the current hospital systems, especially electronic health Records (EHRs), laboratory systems, and billing systems. The ability of various digital tools to interoperate guarantees continuity of clinical processes and data silo that can threaten care quality and efficiency will be avoided. Moreover, the reliability of the system and its performance stability are of the essence because the

disruption of the service may have a direct influence on the clinician trust and patient safety. A cloud-based and modular infrastructure can make healthcare organizations flexible to meet the growing demand, integrate new technologies as they emerge, and be operationally resilient in the long run.

Selection of a platform: Technological and clinical assessment of the telehealth platform and not short-term cost should be applied to select an appropriate platform. CIOs should evaluate the usability of the platform by both clinicians and patients, the compatibility of the platform with the current IT architecture, the vendor technical support and whether the platform supports any future upgrades. The inappropriately chosen platform can lead to the rise of clinician resistance, disruption of work flows, and underutilization of the telehealth services, but an aligned platform will contribute to the efficiency of operations and clinical acceptance.

Interoperability: Interoperability is crucial towards facilitating a smooth flow of data between various healthcare platforms and digital tools. The telehealth platforms should have the capability to interact with laboratory information systems, pharmacy systems, imaging platforms, and hospital information systems through standardized data formats. Interoperability causes discontinuity in the information, overlap in documentation, and clinical inefficiency, which eventually compromise the quality of care and patient safety.

EHR integration: The successful integration with Electronic Health Records will see to it that telehealth encounters are completely integrated into regular clinical operations. By having instant access to patient history, diagnostic outcomes, and treatment plans, clinicians are able to make precise judgments when conducting virtual consultations. Also, telehealth visits are automatically documented in EHR systems, which decreases

administrative workload and data entry mistakes as well as promotes continuity of care in a physical and virtual healthcare environment.

Scalability: The telehealth infrastructure should be in such a way that it can scale according to increasing number of patients, new line of service and emergencies such as a health crisis. Scalable systems enable healthcare organizations to add users, services and functionality without significant system redesign or performance deterioration. Cloud-based and modular systems have proven to be useful especially in providing flexibility, cost effectiveness and long-term digital transformation objectives.

System reliability: The system reliability is also important in telehealth environment, as technical failures can impact the care delivery and undermine the trust between a patient and a clinician. CIOs should have good network bandwidth, system redundancy, and frequent testing of the system, and bouncy technical support systems. Reliable systems will increase the level of user trust, improve the level of service interruption, and minimize the clinical error risk associated with system downtime or data loss.

2. Data Security, Compliance, and System Governance

The regulatory compliance and data security are among the essential factors that determine the trust and sustainability in telehealth ecosystems, which demand the CIOs to build the effective governance structures. Telehealth services are sensitive to the patient information, and without an adequate security strategy, these services will be prone to cyber threats and privacy breaches. CIOs are urged to install a detailed cybersecurity, including encryption, access control, and frequent monitoring of the system in order to protect patient information. The healthcare regulations and data protection laws also matter in order to make sure that no legal risks and financial penalties are introduced. Besides, it ought to have distinct governance frameworks that define accountability, risk management

processes and decision-making to ensure that the digital health technologies are always used consistently and ethically. Best system governance is more than risk mitigation but also increases trust between the stakeholders and telehealth services.

Data privacy security: Data privacy security is a pre-requisite of telehealth provision because e-consultations presuppose that medical data of sensitive nature transfer and storage. Good encryption protocols, role-based access controls, and secure authentication mechanisms should be employed by the CIOs as a way of preventing unauthorized access. The high level of privacy will ensure compliance with legal standards and the enhancement of the degree of trust in patients and their willingness to engage the telehealth systems.

Cybersecurity precautions: Ransomware, phishing, and data breaches are among the cyber-attacks that are becoming a common issue in the telehealth systems. To overcome these risks, CIOs must take active measures to secure the system, including constant monitoring of the system, regular vulnerability checking, and cybersecurity training of health workers. The best incident response strategy also ensures the organizations have the ability to contain and recover security attacks in a short time with minimum interruption.

Regulatory compliance: Telehealth services must have a legal and ethical operation that is founded on the observance of the healthcare regulation and data protection legislation. The CIOs ought to ensure that the telehealth systems do not contravene the national and international laws on data storage, patient consent, transfer of data across national boundaries, and the maintenance of digital records. Compliance with the regulations minimizes the legal risks, institutional credibility, and the ability to establish relationships with insurers and technology suppliers.

Risk mitigation: Well-developed risk mitigation implies the ability to recognize the possible technical, operational, and legal risks related to telehealth systems and mitigate

them in advance. CIOs are expected to perform risk assessment on regular basis, create contingency plans and come up with specific procedures in the event of failure of the system and data breach. Through an organised approach to risk management, healthcare institutions can secure continuity of services and safeguard their image in a more digitised healthcare setting.

System governance: Good system governance provides the authority in decision making, accountability systems and performance control on the telehealth program. CIOs should have a policy that governs the use of data, access to systems, and the vendors and assessment of technologies. Open governance systems assist in avoiding unethical application of digital health technologies, transforming telehealth programs into organisational objectives, and long-term implementation.

Recommendations for Telehealth Entrepreneurs

The entrepreneur in telehealth must possess a balance in innovation, regulatory sensitivity and business sustainability as regards to market development. Successful ventures should consider targeting well-developed market areas where telehealth can provide measurable clinical and access-related value, rather than competing on a general basis in well-saturated markets. A compliance market entry planning, differentiated value proposition, and flexible business models should be included in the entrepreneurial strategy, which would be consistent with the incentives of healthcare providers and patients, along with the incentives of payers. The emphasis of the scalable digital infrastructure, the strategic partnership with the healthcare entities, and insurance companies, and variable pricing mechanisms is crucial to the growth and operational risk reduction. By combining the technological potential and good governance, financial discipline, and customer-oriented design, the telehealth entrepreneurs can establish healthy

businesses that can not only compete well in the long run but also contribute immensely to the healthcare delivery systems.

1. Market Entry and Competitive Positioning

In the telehealth market, entrepreneurs must embrace entry modes that are strategic in nature and which consider regulatory complexity, competitive intensity, and heterogeneous healthcare requirements. Instead of targeting wide market coverage during the early phases, telehealth enterprises need to determine specific clinical or demographic segments where demand is not met, there are barriers to access, or the lack of service efficiency is the most significant. Competitive positioning requires getting ahead of healthcare regulations, establishing a reputation with healthcare providers and payers, and making differentiations based on the quality of services offered, specialisation, or even technological innovation. The concept of sustainable market entry does not therefore concentrate only on technological novelty but rather on regulatory compliance, clinical value creation and defensible differentiation strategies.

Entry strategies: Telehealth entrepreneurs must implement phased or pilot-based entry strategies, i.e. start small in a geographical area or service line. This method enables companies to test models of operation, perfect service delivery and show clinical and economic value on a preliminary scale and then scale. Virgin's introduction also minimises regulatory and financial risk and allows learning from the experience of early adopters.

Niche targeting: Specialising in specific segments, such as treatment of chronic diseases, mental health services, rural population, or post-discharge follow-up care, can allow new companies to not enter direct competition with large and well-established platforms. Niche focus increases the clarity of value proposition, user adoption is accelerated, and reputation building in areas where telehealth offers recognizable and quantifiable advantages is achieved.

Navigating regulations: To enter the market, it is necessary to talk to the healthcare regulators, legal counsel, and accreditation agencies early. Entrepreneurs need to learn about the licensing rules, data protection regulations, and the eligibility of reimbursement in various jurisdictions. Active regulatory adherence minimises the entry lag, creates institutional confidence and makes it more appealing to healthcare partners and insurers.

Differentiation strategies: Competitive differentiation can be in the form of better user experience, clinical specialisation, integration with a wearable or AI-based product, or care coordination. This kind of clear differentiation minimises competition based on price and enhances brand position to allow telehealth ventures to compete based on value, but not scale alone.

2. Business Model and Pricing Strategies

The ability of the telehealth ventures to be sustainable lies in the choice of flexible business models and pricing approaches that balance cost, financial sustainability, and expandability. This is because there is no universal revenue model applicable to telehealth users and payers due to the diversity of users and payers. The most common combination of pricing mechanisms, value propositions that are congruent with stakeholder incentive, and the use of strategic partnership strategies to improve reach and lower the operation costs are common in successful telehealth entrepreneurs. Business models should thus be versatile, data-driven, and should be flexible to change with regulatory and reimbursement landscapes.

Subscription-based models: Subscription models provide stable flows of revenues as they charge healthcare facilities, employers, or patients a recurrent fee to have access to telehealth services. These models tend to be effective especially in chronic care

management and preventive services where in-service would be a must. The subscription pricing helps to improve financial stability and planning of the long-term services.

Pay-per-use prices: Fee-for-service or pay-per-use models offer flexibility and reduced barriers to entry by patients who only need consultation on a limited number of occasions. The given method is ideal in acute care or episodic services but can lead to fluctuations in the revenue. The entrepreneurs have to create a balance between access and pricing models that will be able to capture the cost of service and clinical value.

Strategic collaboration: Collaborations with hospitals, insurers, pharmaceutical firms, or digital health solutions will allow telehealth enterprises to share infrastructure, grow their user base, and enhance access to reimbursement. The collaborative models also lower the cost of acquiring customers and increase the service legitimacy, especially in extremely controlled markets in the healthcare industry.

Scalability: Business models are to be developed to scale the business effectively without directly proportional changes in the operational costs. Scalability is enabled by cloud-based delivery, automation, and standardised clinical protocols, and does not compromise the quality of the services. Application Scalable operating models and pricing mechanisms can make telehealth enterprises sustainable and competitive in the long term.

Recommendations for Insurance Companies and Payers

Healthcare payers and insurance companies can be central to the cost, viability, and the success of telehealth adoption by using reimbursement mandates and policies and their regulatory frameworks. In order to facilitate integration in the long term, payers must leave short-term, volume-based reimbursement models behind and develop payment models that encourage quality, continuity of care, and measurable health outcomes. This necessitates the integration of telehealth reimbursement with value-based care principles, preventive health goals and population health management strategies. Moreover, insurers shall strive

to ensure policy coherence by standardizing coverage requirements, make administrative requirements more straightforward, and assist in data-driven review of telehealth performance. Payers can empower financially sustainable, equitable, and outcome-based virtual care ecosystems by encouraging the alignment of incentives among the providers, patients, and technology platforms and by incorporating telehealth into the mainstream insurance benefit models.

1. Reimbursement Model Design and Incentive Alignment

Payers and system-financing. The long-term viability of telehealth relies on the ability of reimbursement models to respond to financial incentives by responding to care quality, efficiency and outcomes. The traditional fee-for-service systems offer incentives for utilisation in the short-term and can facilitate utilization based on volume but without any related patient outcomes. In comparison, outcome-linked and value-based reimbursement models establish a more effective connection to the purpose of telehealth adoption because they provide incentives to preventive care, continuity, and quantifiable performance outcomes. Risk-sharing models also stimulate providers and telehealth systems to invest in quality and data integration and engage patients. As such, to achieve sustainable telehealth financing, insurers and policymakers should formulate reimbursement systems that give providers an incentive to participate and at the same time contain cost and quality to guarantee quality assurance.

Fee-for-Service Reimbursement Models: Fee-for-service is the most common model that has been adopted because of its simplicity and easy implementation. Nevertheless, its application to telehealth can motivate too many virtual visits without necessarily being clinically effective and therefore can be used only in the short term and not long-term sustainability.

Value-Based and Outcome-Linked Payments: Value-based reimbursement is a payment method that relates payment to health outcomes and patient satisfaction, and the affordability of telehealth. This promotes the adoption of telehealth by providers as an integrated component of care pathways and not as a single service.

Risk-Sharing Arrangements: Financial risk is shared by insurers and providers, telehealth platforms and will encourage accountability and innovation. The models promote investing in preventive care, data analytics, and quality monitoring to reduce long-term healthcare expenditures.

Incentive Alignment Among the Stakeholders: An effective reimbursement design will offer incentives to the patients, providers, and payers and harmonise their interests to reduce resistance to adoption and add to coordinated care delivery.

2. Policy Integration and Long-Term Sustainability

Policy consistency, regulatory transparency, and governance congruence in the healthcare institutions and insurance subsidies level are highly interrelated with telehealth system sustainability. Fragmented legislation and disparate licensing criteria and variations in information protection principles may harm trust in the suppliers and payers to restrict investment and extent. To realise long-term health sustainability, harmonised policy frameworks that reflect the integration of telehealth into the important health strategies in countries and regions, clarify the compliance criteria, and support innovation in terms of safeguarding patient rights should be developed. Telehealth can be made a consistent and consonant component of healthcare delivery by policymakers and healthcare insurers by including it as part of broader healthcare governance systems and financial models.

Governance Coherence in the Health Systems: Coherent governance modalities spur the reduction in the complexity of administration, and easier alignment of the providers, insurers, and technology platforms.

Regulatory Compliance and Standardisation: There should be clear guidelines for licensing, data privacy, and clinical accountability to decrease legal uncertainty and increase access to telehealth services for more individuals.

Integration into National and Regional Health Policy: Inclusion of telehealth in the institutional framework of the public health policy will provide it with institutional support, continuity of funds, and integration into the long-term systems.

System-Wide Sustainability: Policy-based sustainability will guarantee that the telehealth growth does not create inequalities but rather leads to equitable access, quality enhancement, and cost management within the healthcare systems.

12–18 Month Telehealth Implementation Roadmap

Telehealth strategy implementation should occur during 12-18 months to change the strategy into measurable organisational outputs and operational, financial and regulatory risks. The implementation of technology should be accompanied by governance preparedness through the phases of this road map, labour competencies and reimbursement schemes. First stage is founded on simplistic preparedness, e.g., infrastructure inspection, policy consistency and stakeholder engagement; Second stage brings out pilot application, workflow incorporation and performance assessment; and the third phase aids in scale-up, optimisation and institutionalisation of telehealth in the normal care paths. By performing sequencing actions according to specific milestones and continuously tracking such performance indicators as utilisation, patient outcome, cost-efficiency, and compliance, healthcare organisations can impose disciplined implementation, minimise implementation disruption, and receive sustainable competitive and operational advantages of telehealth implementation.

1. Phased Implementation Plan

Implementation of telehealth over a 12 to 18-month period enables the healthcare entities to handle complexity, reduce risk, and ensure that the strategic direction works in concert with operational preparedness. Phase I (06 months) will involve organisational preparedness through examining the infrastructure, examining regulatory compliance, aligning the stakeholders and designing the pilot program. Phase II (6-12 months) will emphasise on controlled deployment, including pilot implementation, integration clinical workflow, staff and early performance assessment. Phase III (12-18 months) supports scaling up and institutionalisation to expand the service coverage, to streamline the technology and processes, and to incorporate telehealth into the regular care trajectories and reimbursement frameworks. The incremental approach will render the integration of telehealth systematic during the experimentation phase and integration with the whole enterprise and maintenance of quality, compliance and financial soundness.

- **Phase I (0–6 Months): Strategic and Operational Readiness**

The tasks undertaken in this stage are technology audits, regulatory and legal compliance testing, stakeholder analysis, and pilot use cases selection. The goal is to put a solid base that will reduce the risk of implementation.

- **Phase II (6–12 Months): Pilot Deployment and Integration**

The action plan at this stage includes starting telehealth pilot programs, connecting platforms with electronic health records, redesigning clinical processes, and educating clinicians and administration staff. Constant surveillance will see technical or operational problems detected sooner.

- **Phase III (12–18 Months): Scale-Up and Optimisation**

The last stage is the extension of telehealth services throughout the various departments, performance metrics are utilized to optimize care delivery, and telehealth becomes a standard operating procedure and model of care.

2. Milestones, Resource Requirements, and Success Metrics

The telehealth implementation should include certain milestones, an adequate amount of resources, as well as well-developed systems of performance measurement to ensure responsible and improve the program. Goals should be in keeping with each step of implementation and possess the necessary financial, human and technological resources. The measures of success must not be limited to the rate of utilisation but must include clinical outcomes, patient experience, cost efficiency and compliance measures, which will enable the organisations to not only measure strategic performance but also operational performance.

- **Implementation Milestones:** Milestones are the completion of pilot launch, system integration milestones, regulatory compliance and scale-up preparedness evaluations, which will give a clear progress indicator at each stage.
- **Human Resource Requirements:** Specialists (telehealth coordinators), IT support teams, clinician champions, and training experts are needed to facilitate adoption and change management.
- **Financial and Technological Resources:** It needs to be allocated financial resources to license the platform, cybersecurity, system integration, training, and continue maintenance to make it reliable and scalable.

- **Success Metrics and KPIs:** Patient satisfaction ratings, clinician adoption, cost savings, and no-show rates reduction, clinical outcome measures, and regulatory compliance rates are among the key indicators.

6.5 Conclusion

According to the research, “telehealth” is a game-changing method of delivering HC that, on the whole, greatly improves accessibility, convenience, and patient happiness. Previous “telehealth” users are more likely to have a positive outlook on continuing to use “telehealth”, according to the statistics; hence, first impressions are crucial for future perceptions.

The main factors that influenced the patients' participation and trust in “telehealth” were patient satisfaction, quality of service as perceived by the patients, and good technical support. The issues involving technology and fear of poor quality of care have not overshadowed the positive side of “telehealth,” which is safety, reliability, and convenience in addressing the challenge of patients not getting access to in-person health care. This is especially the case with digitisation being done more and more, and the need for health services that are more flexible being loudest.

Altogether, these role-specific recommendations show that telehealth sustainability requires a concerted effort by leadership, finance, technology, market innovation, reimbursement systems, and public policy. The empirical results of the study prove the feasibility of telehealth, but to be successful, the project should be implemented with organised governance, financial controls, technological reliability, and clarity of regulations.

Managerial implications for healthcare organisations and policymakers.

Based on the findings of this study, several managerial implications can be identified for healthcare organisations, administrators, and policymakers involved in the implementation and management of telehealth services.

1. Investment in Digital Infrastructure

Healthcare organisations should prioritise their funding for dependable digital infrastructure which encompasses secure telehealth systems and fast internet access and unified electronic health record (EHR) networks. The study findings show that technical disruptions together with connectivity problems lead to decreased user satisfaction and decreased telehealth service effectiveness. Organizations need to establish their technological systems as stable and secure to deliver full remote healthcare services which maintain high standards.

2. Training and Capacity Building for Healthcare Professionals

Healthcare managers should implement training programmes that enhance clinicians' ability to deliver effective virtual care. The process of telehealth consultations demands clinicians to have expertise in three specific areas which include digital communication and remote patient assessment and telehealth platform usage. Continuous professional development together with technical training helps clinicians build their confidence in using telehealth technologies which results in better patient satisfaction and improved service quality.

3. Development of Clear Regulatory and Reimbursement Policies

Policymakers need to create permanent regulatory systems together with reimbursement models which will enable telehealth services to be used throughout the long term. Healthcare organizations face barriers to telehealth service implementation because they lack certainty about three key factors which include payment models and licensing

regulations and cross-jurisdictional practice rules. The presence of defined guidelines together with reliable reimbursement systems will motivate healthcare organizations to adopt telehealth as a standard part of their patient treatment workflows.

4. Adoption of Hybrid Healthcare Delivery Models

Healthcare organisations should implement hybrid care models which combine telehealth services with traditional face-to-face healthcare provision. The findings demonstrate that telehealth functions best for follow-up consultations and routine health assessments and treatment of minor medical problems, although some health conditions still need actual medical tests. Hybrid models enable healthcare systems to achieve maximum operational efficiency while delivering excellent clinical care standards.

5. Strengthening Technical Support and Patient Assistance Services

Healthcare providers should establish dedicated technical support systems to assist patients and clinicians when technical issues arise during telehealth consultations. The user experience improves through complete system functionality which requires immediate technical support during operational disruptions. The support provides help desks along with instructional materials and patient guidance for effective telehealth platform usage.

6. Promoting Digital Inclusion and Accessibility

The digital gap in healthcare access needs to be tackled through special programs which should be developed by both policymakers and healthcare organizations. Telehealth services present accessibility challenges for certain population groups which include older adults and residents of rural areas and people who have restricted digital skills. The achievement of equitable telehealth distribution across different population groups requires three essential components which include digital literacy training programs and enhanced internet access and development of user-friendly telehealth systems.

7. Enhancing Patient Engagement and Awareness

The digital gap in healthcare access needs to be tackled through special programs which should be developed by both policymakers and healthcare organizations. Telehealth services present accessibility challenges for certain population groups which include older adults and residents of rural areas and people who have restricted digital skills. The achievement of equitable telehealth distribution across different population groups requires three essential components which include digital literacy training programs and enhanced internet access and development of user-friendly telehealth systems.

REFERENCES

- Abitha, S. B., Santhoshi, R., Lavanya, G., Sanjeev, R., Bharanika, S., & Mohan Prakash, Y. (2024). A Home driven Physiological Parameter Monitoring Healthcare Teleapp System. *2024 2nd International Conference on Advancement in Computation & Computer Technologies (InCACCT)*, 823–828. <https://doi.org/10.1109/InCACCT61598.2024.10550996>
- Afrihyia, E., Forkuo, A. Y., Mustapha, A. Y., Chinonso, E., Chianumba, & Omotayo, O. (2025). *Telemedicine and Remote Patient Care: Expanding Access and Reducing Inefficiencies in the Healthcare System*. https://www.researchgate.net/publication/393975531_Telemedicine_and_Remote_Patient_Care_Expanding_Access_and_Reducing_Inefficiencies_in_the_Healthcare_System
- Afshar, M. Z., & Mutahir, H. (2025). *Performance Evaluation Using Balanced Scorecard Framework- Insights From A Public Sector Case Study*. 5, 40–47. https://www.researchgate.net/publication/388876005_Performance_Evaluation_Using_Balanced_Scorecard_Framework-Insights_From_A_Public_Sector_Case_Study
- Aljedaimi, M. (2024). Significance of Artificial Intelligence and Machine Learning on efficiency and automation of audit. *International Journal of Scientific and Research Publications*, 14(11), 204–205. <https://doi.org/10.29322/ijsrp.14.11.2024.p15527>
- Alkarkhi, A. F. M. (2021). Chi-square test for one sample variance. In *Applications of Hypothesis Testing for Environmental Science*. <https://doi.org/10.1016/b978-0-12-824301-5.00003-4>
- Alkattan, A., Albokhadaim, I., Almofti, Y. A., Alrasheed, M., & Kandeel, M. (2025). Evolution and emerging directions in occupational therapy for autism: A bibliometric

- evaluation of global literature. *International Journal of Innovative Research and Scientific Studies*, 8(3), 3431–3445. <https://doi.org/10.53894/ijirss.v8i3.7253>
- Almathami, H. K. Y., Win, K. T., & Vlahu-Gjorgievska, E. (2020). Barriers and Facilitators That Influence Telemedicine-Based, Real-Time, Online Consultation at Patients' Homes: Systematic Literature Review. *Journal of Medical Internet Research*, 22(2), e16407. <https://doi.org/10.2196/16407>
- Almuhaideb, M., Bandaru, D., Chawareb, E. A., Yafi, F. A., & Hammad, M. A. M. (2025). Ethical perspective on telemedicine usage in sexual medicine: A friend or a foe? *UroPrecision*, 3(1), 47–53. <https://doi.org/10.1002/uro2.111>
- Amjad, A., Kordel, P., & Fernandes, G. (2023). A Review on Innovation in Healthcare Sector (Telehealth) through Artificial Intelligence. In *Sustainability (Switzerland)*. <https://doi.org/10.3390/su15086655>
- Anawade, P. A., Sharma, D., & Gahane, S. (2024). A Comprehensive Review on Exploring the Impact of Telemedicine on Healthcare Accessibility. *Cureus*. <https://doi.org/10.7759/cureus.55996>
- Andrews, J. C., Bogliatto, F., Lawson, H. W., & Bornstein, J. (2016). Speaking the Same Language: Using Standardized Terminology. In *Journal of Lower Genital Tract Disease*. <https://doi.org/10.1097/LGT.0000000000000157>
- Anggo, S., Rahman, W., Chai, S., & Pao, C. (2025). Telemedicine in the Digital Era: Changing the Face of Health Services with Virtual Technology. *Journal of World Future Medicine, Health and Nursing*, 3(1), 30–41. <https://doi.org/10.70177/health.v3i1.1897>
- Ansarian, M., & Baharlouei, Z. (2023). Applications and Challenges of Telemedicine: Privacy-Preservation as a Case Study. *Archives of Iranian Medicine*, 26(11), 654–661. <https://doi.org/10.34172/aim.2023.96>

- Anthony, D. L., Campos-Castillo, C., & Lim, P. S. (2018). Who isn't using patient portals and why? Evidence and implications from a national sample of US adults. *Health Affairs*. <https://doi.org/10.1377/hlthaff.2018.05117>
- Anwari, F., Hutomo, G. S., Supinganto, A., Utary, D., & Rita, R. S. (2024). Implementation of Telemedicine in Medical Practice: Opportunities and Barriers. *Global International Journal of Innovative Research*, 1(2), 146–153. <https://doi.org/10.59613/global.v1i2.16>
- Association, A. H. (2025). *Fact Sheet: Telehealth* (pp. 1–3). <https://www.aha.org/fact-sheets/2025-02-07-fact-sheet-telehealth>
- Asua, J., Orruño, E., Reviriego, E., & Gagnon, M. P. (2012). Healthcare professional acceptance of telemonitoring for chronic care patients in primary care. *BMC Medical Informatics and Decision Making*, 12(1), 139. <https://doi.org/10.1186/1472-6947-12-139>
- Baden-Fuller, C., & Haefliger, S. (2013). Business Models and Technological Innovation. *Long Range Planning*. <https://doi.org/10.1016/j.lrp.2013.08.023>
- Bashshur, R. L., & Shannon, G. W. (2009). History of Telemedicine: Evolution, Context, and Transformation. *Healthcare Informatics Research*, 16(1), 65. <https://doi.org/10.4258/hir.2010.16.1.65>
- Bashshur, R., Shannon, G., Krupinski, E., & Grigsby, J. (2011). The taxonomy of telemedicine. *Telemedicine and E-Health*. <https://doi.org/10.1089/tmj.2011.0103>
- Belliger, A., & Krieger, D. J. (2018). The Digital Transformation of Healthcare. In *Knowledge management in digital change: new findings and practical cases* (pp. 311–326). Springer. https://doi.org/10.1007/978-3-319-73546-7_19
- Berlilana, Hariguna, T., & Nurfaizah. (2017). Understanding of Public Behavioral Intent to Use e-Government Service: An Extended of Unified Theory of Acceptance Use of

- Technology and Information System Quality. *Procedia Computer Science*, 124, 585–592. <https://doi.org/10.1016/j.procs.2017.12.193>
- Bestsenny, O., Gilbert, G., Harris, A., & Rost, J. (2021). *Telehealth: A quarter-trillion-dollar post-COVID-19 reality?* Mckensey&Company. <https://www.mckinsey.com/industries/healthcare/our-insights/telehealth-a-quarter-trillion-dollar-post-covid-19-reality#/>
- Bitar, H., & Alismail, S. (2021). The role of eHealth, telehealth, and telemedicine for chronic disease patients during COVID-19 pandemic: A rapid systematic review. In *Digital Health*. <https://doi.org/10.1177/20552076211009396>
- Bober, T., Rollman, B. L., Handler, S., Watson, A., Nelson, L. A., Faieta, J., & Rosland, A.-M. (2024). Digital Health Readiness: Making Digital Health Care More Inclusive. *JMIR MHealth and UHealth*, 12, e58035. <https://doi.org/10.2196/58035>
- Bruijl, G. H. T. (2018). The Relevance of Porter’s Five Forces in Today’s Innovative and Changing Business Environment. *SSRN Electronic Journal*. <https://doi.org/10.2139/ssrn.3192207>
- Burrell, D. N. (2023). Understanding telemedicine strategies through field-based observations, interviews and benchmarking activities. *Health Economics and Management Review*, 4(4), 58–68. <https://doi.org/10.61093/hem.2023.4-05>
- Burrell, D. N. (2024). Telehealth Technologies, Disruptions, and Health Literacy. *Health Economics and Management Review*, 5(4), 33–46. <https://doi.org/10.61093/hem.2024.4-03>
- Butt, S. A., Naseer, M., Ali, A., Khalid, A., Jamal, T., & Naz, S. (2024). Remote mobile health monitoring frameworks and mobile applications: Taxonomy, open challenges, motivation, and recommendations. *Engineering Applications of Artificial Intelligence*, 133, 108233. <https://doi.org/10.1016/j.engappai.2024.108233>

- Caffery, L. J., Martin-Khan, M., & Wade, V. (2017). Mixed methods for telehealth research. *Journal of Telemedicine and Telecare*. <https://doi.org/10.1177/1357633X16665684>
- Cannavacciuolo, L., Capaldo, G., & Ponsiglione, C. (2023). Digital innovation and organizational changes in the healthcare sector: Multiple case studies of telemedicine project implementation. *Technovation*, 120, 102550. <https://doi.org/10.1016/j.technovation.2022.102550>
- Čaplová, Z., & Šváblová, P. (2020). IBM SPSS statistics. In *Statistics and Probability in Forensic Anthropology*. <https://doi.org/10.1016/B978-0-12-815764-0.00027-7>
- Carpenter, A. B., Sheppard, E., Atabaki, S., Shur, N., Tigranyan, A., Benchoff, T., Snyder, A., Fisher, A., & Cleary, K. (2021). A Symposium on the Clinic of the Future and Telehealth: Highlights and Future Directions. *Cureus*. <https://doi.org/10.7759/cureus.15234>
- Carroll, C. (2023). Telehealth's Role in Modern Healthcare: Accessibility, Adherence, and Challenges. *Journal of Innovations*. <https://doi.org/10.62470/4f245146>
- Chakraborty, I., Edirippulige, S., & Ilavarasan, V. (2023). The role of telehealth startups in healthcare service delivery: A systematic review. *International Journal of Medical Informatics*, 174, 105048. <https://doi.org/10.1016/j.ijmedinf.2023.105048>
- Chau, P. Y. K., & Hu, P. J. (2002). Examining a Model of Information Technology Acceptance by Individual Professionals: An Exploratory Study. *Journal of Management Information Systems*, 18(4), 191–229. <https://doi.org/10.1080/07421222.2002.11045699>
- Chen, A. H., Bakar, N. F. A., & Lam, C. S. Y. (2020). Comparison of open-ended and close-ended questions to determine signs and symptoms of eye problems among children. *Journal of Optometry*. <https://doi.org/10.1016/j.optom.2019.07.002>

- Chen, C.-H., Lan, Y.-L., Yang, W.-P., Hsu, F.-M., Lin, C.-L., & Chen, H.-C. (2019). Exploring the Impact of a Telehealth Care System on Organizational Capabilities and Organizational Performance from a Resource-Based Perspective. *International Journal of Environmental Research and Public Health*, 16(20), 3988. <https://doi.org/10.3390/ijerph16203988>
- Cheong, H., Lyons, A., Houghton, R., & Majumdar, A. (2023). Secondary Qualitative Research Methodology Using Online Data within the Context of Social Sciences. *International Journal of Qualitative Methods*, 22. <https://doi.org/10.1177/16094069231180160>
- Cheung, A. K.-L. (2021). Structured Questionnaires. In *Encyclopedia of Quality of Life and Well-Being Research* (pp. 1–3). Springer International Publishing. https://doi.org/10.1007/978-3-319-69909-7_2888-2
- Choi, N. G., DiNitto, D. M., Marti, C. N., & Choi, B. Y. (2022). Telehealth Use Among Older Adults During COVID-19: Associations With Sociodemographic and Health Characteristics, Technology Device Ownership, and Technology Learning. *Journal of Applied Gerontology*, 41(3), 600–609. <https://doi.org/10.1177/07334648211047347>
- Christensen, C. M., Bohmer, R. M. J., & Kenagy, J. (2000). Will disruptive innovations cure health care? *Harvard Business Review*, 78(5), 102–112,199. <https://hbr.org/2000/09/will-disruptive-innovations-cure-health-care>
- Christensen, C. M., & Mitra, P. (2022). The Innovator's Dilemma: When New Technologies Cause Great Firms to Fail. In *SWE Magazine*. <https://doi.org/10.2307/40252749>
- Chu, C., Cram, P., Pang, A., Stamenova, V., Tadrus, M., & Bhatia, R. S. (2021). Rural telemedicine use before and during the COVID-19 pandemic: Repeated cross-

- sectional study. *Journal of Medical Internet Research*. <https://doi.org/10.2196/26960>
- Cressman, D. (2021). Disruptive innovation: An organizational strategy and a technological concept. *Handbook on Alternative Theories of Innovation*, 182–196. <https://doi.org/10.4337/9781789902303.00022>
- Dalloul, A. H., Miramirkhani, F., & Kouhalvandi, L. (2023). A Review of Recent Innovations in Remote Health Monitoring. *Micromachines*, 14(12), 2157. <https://doi.org/10.3390/mi14122157>
- Davis, F. D. (1989). Perceived Usefulness, Perceived Ease of Use, and User Acceptance of Information Technology. *MIS Quarterly*, 13(3), 319–340. <https://doi.org/10.2307/249008>
- Demeke, H. B., Merali, S., Marks, S., Pao, L. Z., Romero, L., Sandhu, P., Clark, H., Clara, A., McDow, K. B., Tindall, E., Campbell, S., Bolton, J., Le, X., Skapik, J. L., Nwaise, I., Rose, M. A., Strona, F. V., Nelson, C., & Siza, C. (2021). Trends in Use of Telehealth Among Health Centers During the COVID-19 Pandemic — United States, June 26–November 6, 2020. *MMWR. Morbidity and Mortality Weekly Report*, 70(7), 240–244. <https://doi.org/10.15585/mmwr.mm7007a3>
- Dhabarde, D. M., Maske, M. P., Baheti, J. R., & Shahare, H. V. (2023). A Tele-Healthcare System For Diagnosis and Treatment of Patients During A Pandemic. In *Handbook of Research on Artificial Intelligence and Soft Computing Techniques in Personalized Healthcare Services* (pp. 279–299). Apple Academic Press. <https://doi.org/10.1201/9781003371250-16>
- Dickinson, J. K., Guzman, S. J., Maryniuk, M. D., O'Brian, C. A., Kadohiro, J. K., Jackson, R. A., D'Hondt, N., Montgomery, B., Close, K. L., & Funnell, M. M. (2017). The use of language in diabetes care and education. *Diabetes Care*. <https://doi.org/10.2337/dci17-0041>

- Dorsey, E. R., & Topol, E. J. (2016). State of Telehealth. *New England Journal of Medicine*, 375(2), 154–161. <https://doi.org/10.1056/NEJMra1601705>
- Drummond, M. F., Sculpher, M. J., Claxton, K., Stoddart, G. L., & Torrance, G. W. (2015). *Methods for the economic evaluation of health care programmes*. Oxford university press. <https://global.oup.com/academic/product/methods-for-the-economic-evaluation-of-health-care-programmes-9780199665884?cc=in&lang=en&>
- Elia, G., Margherita, A., & Passiante, G. (2020). Digital entrepreneurship ecosystem: How digital technologies and collective intelligence are reshaping the entrepreneurial process. *Technological Forecasting and Social Change*, 150, 119791. <https://doi.org/10.1016/j.techfore.2019.119791>
- Eungoo, K., & Hwang, H.-J. (2023). The Importance of Anonymity and Confidentiality for Conducting Survey Research. *Journal of Research and Publication Ethics*, 4(1), 1–7. <https://doi.org/10.15722/jrpe.4.1.202303.1>
- Ezeamii, V. C., Okobi, O. E., Wambai-Sani, H., Perera, G. S., Zaynieva, S., Okonkwo, C. C., Ohaiba, M. M., William-Enemali, P. C., Obodo, O. R., & Obiefuna, N. G. (2024). Revolutionizing Healthcare: How Telemedicine Is Improving Patient Outcomes and Expanding Access to Care. *Cureus*, 16(7), e63881. <https://doi.org/10.7759/cureus.63881>
- Fagherazzi, G., Goetzinger, C., Rashid, M. A., Aguayo, G. A., & Huiart, L. (2020). Digital Health Strategies to Fight COVID-19 Worldwide: Challenges, Recommendations, and a Call for Papers. *Journal of Medical Internet Research*, 22(6). <https://doi.org/10.2196/19284>
- Fakhkhari, H., Bounabat, B., & Kassou, I. (2023). Digital Health Taxonomy. *Proceedings of the 6th International Conference on Networking, Intelligent Systems & Security*, 1–9. <https://doi.org/10.1145/3607720.3607780>

- Fatehi, F., & Wootton, R. (2012). Telemedicine, telehealth or e-health? A bibliometric analysis of the trends in the use of these terms. In *Journal of Telemedicine and Telecare*. <https://doi.org/10.1258/jtt.2012.GTH108>
- Flickinger, T. E., Saha, S., Moore, R. D., & Beach, M. C. (2013). Higher quality communication and relationships are associated with improved patient engagement in HIV care. *Journal of Acquired Immune Deficiency Syndromes*. <https://doi.org/10.1097/QAI.0b013e318295b86a>
- Ftouni, R., AlJardali, B., Hamdanieh, M., Ftouni, L., & Salem, N. (2022). Challenges of Telemedicine during the COVID-19 pandemic: a systematic review. *BMC Medical Informatics and Decision Making*, 22(1), 207. <https://doi.org/10.1186/s12911-022-01952-0>
- Gajarawala, S. N., & Pelkowski, J. N. (2021). Telehealth Benefits and Barriers. *The Journal for Nurse Practitioners*, 17(2), 218–221. <https://doi.org/10.1016/j.nurpra.2020.09.013>
- Garfan, S., Alamoodi, A. H., Zaidan, B. B., Al-Zobbi, M., Hamid, R. A., Alwan, J. K., Ahmaro, I. Y. Y., Khalid, E. T., Jumaah, F. M., Albahri, O. S., Zaidan, A. A., Albahri, A. S., Al-qaysi, Z. T., Ahmed, M. A., Shuwandy, M. L., Salih, M. M., Zughoul, O., Mohammed, K. I., & Momani, F. (2021). Telehealth utilization during the Covid-19 pandemic: A systematic review. In *Computers in Biology and Medicine*. <https://doi.org/10.1016/j.compbiomed.2021.104878>
- Giacalone, A., Marin, L., Febbi, M., Franchi, T., & Tovani-Palone, M. R. (2022). eHealth, telehealth, and telemedicine in the management of the COVID-19 pandemic and beyond: Lessons learned and future perspectives. *World Journal of Clinical Cases*, 10(8), 2363–2368. <https://doi.org/10.12998/wjcc.v10.i8.2363>
- Govers, M., & Amelsvoort, P. van. (2023). A theoretical essay on socio-technical systems

- design thinking in the era of digital transformation. *Gruppe. Interaktion. Organisation. Zeitschrift Für Angewandte Organisationspsychologie (GIO)*, 54(1), 27–40. <https://doi.org/10.1007/s11612-023-00675-8>
- Grace, H., Banson, K., & Saraf, A. (2023). Mixed-methods research. In *Translational Radiation Oncology*. <https://doi.org/10.1016/B978-0-323-88423-5.00029-7>
- Greenhalgh, T., Wherton, J., Papoutsis, C., Lynch, J., Hughes, G., A’Court, C., Hinder, S., Fahy, N., Procter, R., & Shaw, S. (2017). Beyond Adoption: A New Framework for Theorizing and Evaluating Nonadoption, Abandonment, and Challenges to the Scale-Up, Spread, and Sustainability of Health and Care Technologies. *Journal of Medical Internet Research*, 19(11), e367. <https://doi.org/10.2196/jmir.8775>
- Gregory, N. S., Shukla, A. P., Noel, J. J., Alonso, L. C., Moxley, J., Crawford, A. J., Martin, P., Kumar, S., Leonard, J. P., & Czaja, S. J. (2023). The feasibility, acceptability, and usability of telehealth visits. *Frontiers in Medicine*, 10. <https://doi.org/10.3389/fmed.2023.1198096>
- Hadian, M., Jelodar, Z. K., Khanbebin, M. J., Atafimanesh, P., Asiabar, A. S., & Dehagani, S. M. H. (2024). Challenges of Implementing Telemedicine Technology: A systematized Review. *International Journal of Preventive Medicine*, 15. https://doi.org/10.4103/ijpvm.ijpvm_48_23
- Haleem, A., Javaid, M., Singh, R. P., & Suman, R. (2021). Telemedicine for healthcare: Capabilities, features, barriers, and applications. In *Sensors International*. <https://doi.org/10.1016/j.sintl.2021.100117>
- Hashiguchi, T. C. O. (2020). *Bringing health care to the patient*. 116(116), 1–104. <https://doi.org/10.1787/8e56ede7-en>
- Herasevich, V., & Subramanian, S. (2019). Tele-ICU Technologies. In *Critical Care Clinics*. <https://doi.org/10.1016/j.ccc.2019.02.009>

- Holden, R. J., & Karsh, B.-T. (2010). The Technology Acceptance Model: Its past and its future in health care. *Journal of Biomedical Informatics*, 43(1), 159–172. <https://doi.org/10.1016/j.jbi.2009.07.002>
- Hong, W. (2024). Analyzing the Public Health Impact of Adopting Telehealth after a Pandemic: A Business Model. *Advances in Economics, Management and Political Sciences*, 116(1), 123–129. <https://doi.org/10.54254/2754-1169/116/20242457>
- Houser, S. H., Flite, C. A., & Foster, S. L. (2023). Privacy and Security Risk Factors Related to Telehealth Services: A Systematic Review. *Perspectives in Health Information Management*, 20(1). <https://pmc.ncbi.nlm.nih.gov/articles/PMC9860467/>
- Hu, P. J., Chau, P. Y. K., Sheng, O. R. L., & Tam, K. Y. (1999). Examining the Technology Acceptance Model Using Physician Acceptance of Telemedicine Technology. *Journal of Management Information Systems*, 16(2), 91–112. <https://doi.org/10.1080/07421222.1999.11518247>
- Hu, S., Song, D., Wan, S., Zhang, S., Luo, C., Li, N., Liu, G., Graça Espírito Santo Vasconcelos, J. da, de Carvalho, L. L. C., Neobísi, E., da Costa, M. L. B., Etchu Takounjou, J., Neves, K. M. Das, dos Ramos da Conceição, L., da Costa Encarnação, M., & Zhao, L.-Y. (2025). Digital health: current applications, challenges, and future directions for enhancing healthcare quality and safety. *Frontiers in Public Health*, 13. <https://doi.org/10.3389/fpubh.2025.1646802>
- Jnr, B. A. (2020). Use of Telemedicine and Virtual Care for Remote Treatment in Response to COVID-19 Pandemic. *Journal of Medical Systems*, 44(7), 132. <https://doi.org/10.1007/s10916-020-01596-5>
- Kaisanesh, J. (2023). *Telemedicine and Remote Patient Monitoring: The Future of Healthcare Delivery*. <https://doi.org/10.31219/osf.io/xugsy>

- Kaliyadan, F., & Kulkarni, V. (2019). Types of variables, descriptive statistics, and sample size. *Indian Dermatology Online Journal*, 10(1), 82. https://doi.org/10.4103/idoj.IDOJ_468_18
- Kaplan, R. S., & Norton, D. P. (1992). The balanced scorecard--measures that drive performance. *Harvard Business Review*. <https://hbr.org/1992/01/the-balanced-scorecard-measures-that-drive-performance-2>
- Karahbolad, H., & Nasrullah, N. (2025). Assessing the Efficiency and Patient Satisfaction of Telemedicine in Managing Chronic Health Conditions. *Global Journal on Quality and Safety in Healthcare*, 8(3), 121–126. <https://doi.org/10.36401/JQSH-24-38>
- Kathiravelu, P., & Moxley, D. (2024). Telehealth in the Circumpolar North: A Perspective of Access and Connectivity. *2024 IEEE Canadian Conference on Electrical and Computer Engineering (CCECE)*, 780–786. <https://doi.org/10.1109/CCECE59415.2024.10667263>
- Katoch, O. R. (2024). *Reshaping Public Health through Technology: Exploring Innovations, Challenges, and Future Avenues*. 1–17. <http://capdr.org/wp-content/uploads/2023/12/18.-Om-Raj-Katoch.pdf>
- Keelara, R., Sutherland, E., & Almyranti, M. (2025). *Beyond the Pandemic: Leading Practices for the Future of Telemedicine* (173rd ed., OECD Health Working Papers). <https://doi.org/10.1787/98863928-en>
- Khan, U. (2024). Remote Patient Monitoring and Telehealth: The Future of Cardiac Care. *International Journal of Engineering Technology Research & Management*, 8(6), 1–8. <https://ijetrm.com/issues/files/Feb-2024-13-1739469414-June2024-20.pdf>
- Kobeissi, M. M., & Hickey, J. V. (2023). An Infrastructure to Provide Safer, Higher-Quality, and More Equitable Telehealth. *The Joint Commission Journal on Quality and Patient Safety*, 49(4), 213–222. <https://doi.org/10.1016/j.jcjq.2023.01.006>

- Koonin, L. M., Hoots, B., Tsang, C. A., Leroy, Z., Farris, K., Jolly, B., Antall, P., McCabe, B., Zelis, C. B. R., Tong, I., & Harris, A. M. (2020). Trends in the Use of Telehealth During the Emergence of the COVID-19 Pandemic — United States, January–March 2020. *MMWR. Morbidity and Mortality Weekly Report*, 69(43), 1595–1599. <https://doi.org/10.15585/mmwr.mm6943a3>
- Kotte, J. P. (1996). Leading Change. In *Harvard Business School* (pp. 1–11). [https://irp-cdn.multiscreensite.com/6e5efd05/files/uploaded/Leading Change.pdf](https://irp-cdn.multiscreensite.com/6e5efd05/files/uploaded/Leading%20Change.pdf)
- Krause, N. (2002). A comprehensive strategy for developing closed-ended survey items for use in studies of older adults. *Journals of Gerontology - Series B Psychological Sciences and Social Sciences*. <https://doi.org/10.1093/geronb/57.5.S263>
- Kruse, C., & Heinemann, K. (2022). Facilitators and Barriers to the Adoption of Telemedicine During the First Year of COVID-19: Systematic Review. *Journal of Medical Internet Research*, 24(1), e31752. <https://doi.org/10.2196/31752>
- Kruse, C. S., Karem, P., Shifflett, K., Vegi, L., Ravi, K., & Brooks, M. (2018). Evaluating barriers to adopting telemedicine worldwide: A systematic review. *Journal of Telemedicine and Telecare*, 24(1), 4–12. <https://doi.org/10.1177/1357633X16674087>
- Kruse, C. S., Krowski, N., Rodriguez, B., Tran, L., Vela, J., & Brooks, M. (2017). Telehealth and patient satisfaction: A systematic review and narrative analysis. *BMJ Open*, 7(8), 1–12. <https://doi.org/10.1136/bmjopen-2017-016242>
- Kumar, S., & Chong, I. (2018). Correlation analysis to identify the effective data in machine learning: Prediction of depressive disorder and emotion states. *International Journal of Environmental Research and Public Health*. <https://doi.org/10.3390/ijerph15122907>
- Kurtaliqui, F., Lancelot Miltgen, C., Viglia, G., & Pantin-Sohier, G. (2024). Using advanced mixed methods approaches: Combining PLS-SEM and qualitative studies. *Journal of*

- Business Research*, 172(November 2023).
<https://doi.org/10.1016/j.jbusres.2023.114464>
- Lagisetty, S. K., Devarajulu, P., & Moka, A. K. (2025). AI-Enhanced Telehealth Platforms: A Comprehensive Analysis of Automated Triage and Personalized Care Systems. 2025 5th Intelligent Cybersecurity Conference (ICSC), 370–377.
<https://doi.org/10.1109/ICSC65596.2025.11139916>
- Lakens, D. (2022). Sample Size Justification. In *Collabra: Psychology*.
<https://doi.org/10.1525/collabra.33267>
- Lee, C., Hoehn-Weiss, M. N., & Karim, S. (2021). Competing both ways: How combining Porter’s low-cost and focus strategies hurts firm performance. *Strategic Management Journal*, 42(12), 2218–2244. <https://doi.org/10.1002/smj.3279>
- Lee, E. C., Grigorescu, V., Enogieru, I., Smith, S. R., Wong Samson, L., Conmy, A. B., & De Lew, N. (2023). *Updated National Survey Trends in Telehealth Utilization and Modality (2021-2022): Issue Brief*.
<https://www.ncbi.nlm.nih.gov/books/NBK604393/>
- Lee, J. S., Bhatt, A., Pollack, L. M., Jackson, S. L., Chang, J. E., Tong, X., & Luo, F. (2024). Telehealth use during the early COVID-19 public health emergency and subsequent health care costs and utilization. *Health Affairs Scholar*, 2(1).
<https://doi.org/10.1093/haschl/qxae001>
- Lewin, K. (1947). Frontiers in Group Dynamics. *Human Relations*, 1(1), 5–41.
<https://doi.org/10.1177/001872674700100103>
- Lin, C.-C. C., Dievler, A., Robbins, C., Sripipatana, A., Quinn, M., & Nair, S. (2018). Telehealth In Health Centers: Key Adoption Factors, Barriers, And Opportunities. *Health Affairs*, 37(12), 1967–1974. <https://doi.org/10.1377/hlthaff.2018.05125>
- Liu, Q., Wald, N., Daskon, C., & Harland, T. (2024). Multiple-choice questions (MCQs)

- for higher-order cognition: Perspectives of university teachers. *Innovations in Education and Teaching International*.
<https://doi.org/10.1080/14703297.2023.2222715>
- Mansoor, M. A., & Ibrahim, A. F. (2025). Telemedicine beyond the Pandemic: Innovations, Challenges, and the Future of Digital Healthcare. In *Telemedicine - Models of Care [Working Title]*. IntechOpen.
<https://doi.org/10.5772/intechopen.1010583>
- Mathews, D., Abernethy, A., Butte, A. J., Ginsburg, P., Kocher, B., Novelli, C., Sandy, L., Smee, J., Fabi, R., Offodile II, A. C., Sherkow, J. S., Sullenger, R. D., Freiling, E., & Balatbat, C. (2023). Telehealth and Mobile Health: Case Study for Understanding and Anticipating Emerging Science and Technology. *NAM Perspectives*, 11(15).
<https://doi.org/10.31478/202311e>
- Maugeri, A., Barchitta, M., Basile, G., & Agodi, A. (2024). Public and Research Interest in Telemedicine From 2017 to 2022: Infodemiology Study of Google Trends Data and Bibliometric Analysis of Scientific Literature. *Journal of Medical Internet Research*, 26, e50088. <https://doi.org/10.2196/50088>
- Michael, E. P. (1985). Competitive Advantage: Creating and sustaining superior performance. In *The Free*. [https://library.uniq.edu.iq/storage/books/file/competitive Advantage/166677556024.pdf](https://library.uniq.edu.iq/storage/books/file/competitive%20Advantage/166677556024.pdf)
- Miller, R. L. (2015). Rogers' Innovation Diffusion Theory (1962, 1995). In *Information Seeking Behavior and Technology Adoption: Theories and Trends*.
<https://doi.org/10.4018/978-1-4666-8156-9.ch016>
- Mishra, P., Pandey, C. M., Singh, U., Gupta, A., Sahu, C., & Keshri, A. (2019). Descriptive statistics and normality tests for statistical data. *Annals of Cardiac Anaesthesia*.
https://doi.org/10.4103/aca.ACA_157_18

- Monaghesh, E., & Hajizadeh, A. (2020). The role of telehealth during COVID-19 outbreak: a systematic review based on current evidence. *BMC Public Health*, 20(1), 1193. <https://doi.org/10.1186/s12889-020-09301-4>
- Morelli, S., Daniele, C., D'Avenio, G., Grigioni, M., & Giansanti, D. (2024). Optimizing Telehealth: Leveraging Key Performance Indicators for Enhanced TeleHealth and Digital Healthcare Outcomes (Telemechron Study). *Healthcare*, 12(13), 1319. <https://doi.org/10.3390/healthcare12131319>
- Mudiyansele, S. B., Stevens, J., Watts, J. J., Toscano, J., Kotowicz, M. A., Steinfert, C. L., Bell, J., Byrnes, J., Bruce, S., Carter, S., Hunter, C., Barrand, C., & Hayles, R. (2019). Personalised telehealth intervention for chronic disease management: A pilot randomised controlled trial. *Journal of Telemedicine and Telecare*, 25(6), 343–352. <https://doi.org/10.1177/1357633X18775850>
- Mwende, W. G. (2024). The Role of Telehealth in Rural Healthcare Access. *Research Output Journal of Public Health and Medicine*, 3(2), 18–21. <https://doi.org/10.59298/ROJPHM/2024/321821>
- Nam, T. (2014). Determining the type of e-government use. *Government Information Quarterly*, 31(2), 211–220. <https://doi.org/10.1016/j.giq.2013.09.006>
- Nazar, N. I. M., Rahman, N. S. A., Alias, N. E., Zaini, S., Kamil, T. K. T. M., Jamani, N. A., & Elnaem, M. H. (2024). *Development and Feasibility Evaluation of a Telehealth System (Impak Sihat) to Empower Rural Population in Malaysia on the Quality Use of Medicines*. <https://doi.org/10.1101/2024.12.22.24319506>
- Ndwabe, H., Basu, A., & Mohammed, J. (2024). Post pandemic analysis on comprehensive utilization of telehealth and telemedicine. *Clinical EHealth*, 7, 5–14. <https://doi.org/10.1016/j.ceh.2023.12.002>
- Nelson, J. (2025). *The Impact of Telemedicine on Diabetes Management During the*

COVID-19 Pandemic: Insights Gained.

https://www.researchgate.net/publication/390582926_The_Impact_of_Telemedicine_on_Diabetes_Management_During_the_COVID-19_Pandemic_Insights_Gained

Nonda, P. K., Rafid Khan, M. S., Jiko, A.-I. A. J., Sultana, R., & Sajib, M. M. H. (2025).

The Evolution and Future of Telemedicine: Challenges, Innovations, and Global Adoption Trends. *Varendra International Journal for Interdisciplinary Research*, 01(01). <https://doi.org/10.64296/vijir.v1i1.04>

Norzaidi, M. D., Salwani, M. I., Chong, S. C., & Rafidah, K. (2008). A Study of Intranet

Usage and Resistance in Malaysia's Port Industry. *Journal of Computer Information Systems*, 49(1), 37–47. <https://doi.org/10.1080/08874417.2008.11645304>

Novielli, J., Kane, L., & Ashbaugh, A. R. (2023). Convenience Sampling Methods in

Psychology: A Comparison Between Crowdsourced and Student Samples. *Canadian Journal of Behavioural Science*. <https://doi.org/10.1037/cbs0000394>

Okesiji, A., Oyasiji, O., Akomolafe, O. O., & Kuponiyi, A. (2024). A Conceptual

Framework for Telemedicine as a Financially Sustainable Healthcare Model: Evaluating ROI, Market Expansion, and Business Growth Strategies. *International Journal of Multidisciplinary Futuristic Development*, 5(2), 28–43. <https://doi.org/10.54660/IJMFD.2024.5.2.28-43>

Olivencia, S. B., & Sasangohar, F. (2023). A sociotechnical framework for integration of

telehealth into clinical workflow. *IISE Transactions on Healthcare Systems Engineering*, 13(3), 248–259. <https://doi.org/10.1080/24725579.2023.2211083>

Olorunsogo, T. O., Balogun, O. D., Oluwatoyin Ayo-Farai, Ogundairo, O., Maduka, C. P.,

Okongwu, C. C., & Onwumere, C. (2024). Reviewing the evolution of U. S. telemedicine post-pandemic by analyzing its growth, acceptability, and challenges in remote healthcare delivery during Global Health Crises. *World Journal of Biology*

- Pharmacy and Health Sciences*, 17(1), 075–090.
<https://doi.org/10.30574/wjbphs.2024.17.1.0010>
- Omboni, S., Padwal, R. S., Alessa, T., Benczúr, B., Green, B. B., Hubbard, I., Kario, K., Khan, N. A., Konradi, A., Logan, A. G., Lu, Y., Mars, M., McManus, R. J., Melville, S., Neumann, C. L., Parati, G., Renna, N. F., Ryvlin, P., Saner, H., ... Wang, J. (2022). The worldwide impact of telemedicine during COVID-19: current evidence and recommendations for the future. *Connected Health*.
<https://doi.org/10.20517/ch.2021.03>
- Organization, W. H. (2022). *Consolidated Telemedicine Implementation Guide*.
<https://www.who.int/publications/i/item/9789240059184>
- Organization, W. H. (2023). *Classification of digital interventions, services and applications in health: a shared language to describe the uses of digital technology for health, 2nd ed* (2nd ed.). <https://www.who.int/publications/i/item/9789240081949>
- Osterwalder, A., & Pigneur, Y. (2010). *Business Model Generation: A Handbook for Visionaries, Game Changers, and Challengers*.
https://www.researchgate.net/publication/236650466_Business_Model_Generation_A_Handbook_for_Visionaries_Game_Changers_and_Challengers
- Patel, W. D., & Koyuncu, H. (2024). Advancements in Telehealth: Enhancing Breast Cancer Detection and Health Automation through Smart Integration of IoT and CNN Deep Learning in Residential and Healthcare Settings. *Journal of Advanced Research in Applied Sciences and Engineering Technology*, 45(2), 214–226.
<https://doi.org/10.37934/araset.45.2.214226>
- Pol, & Khedkar. (2024). Telemedicine and telehealth. *World Journal of Advanced Research and Reviews*, 24(1), 883–891.
<https://doi.org/10.30574/wjarr.2024.24.1.3106>

- Porteny, T., Brophy, S. A., & Burroughs, E. (2025). Experiences of Telehealth Reimbursement Policies in Federally Qualified Health Centers. *JAMA Network Open*, 8(2), e2459554. <https://doi.org/10.1001/jamanetworkopen.2024.59554>
- Ramirez, L. A. R. (2024). Current trends and future directions of mHealth in psychology: Challenges and promising perspectives. *Mexican Journal of Medical Research ICSA*, 12(24), 89–95. <https://doi.org/10.29057/mjmr.v12i24.12430>
- Research, P. (2025). *Telemedicine Market Growth Driven by Rising Demand for Remote Healthcare and Virtual Consultations*. <https://www.precedenceresearch.com/telemedicine-market>
- Rintyarna, B. S., Sasmiyanto, Insantuan, O. D., Widiawati, I., & Purwoko, R. Y. (2023). Telehealth in Remote Areas: A New Artificial Intelligence-Based Model. *International Journal of Science and Society*, 5(4), 243–254. <https://doi.org/10.54783/ijssoc.v5i4.782>
- Rogers, E. M. (2003). *Diffusion of Innovations* (Third Edit). <https://teddykw2.wordpress.com/wp-content/uploads/2012/07/everett-m-rogers-diffusion-of-innovations.pdf>
- Roy, J., Levy, D., & Senathirajah, Y. (2021). Defining Telehealth for Research, Implementation, and Equity (Preprint). *Journal of Medical Internet Research*, 24. <https://doi.org/10.2196/35037>
- Şahin, E., Veizi, B. G. Y., & Naharci, M. I. (2024). Telemedicine interventions for older adults: A systematic review. *Journal of Telemedicine and Telecare*, 30(2), 305–319. <https://doi.org/10.1177/1357633X211058340>
- Sahoo, S., Sahoo, J., Kumar, S., Lim, W. M., & Ameen, N. (2023). Distance is no longer a barrier to healthcare services: current state and future trends of telehealth research. *Internet Research*, 33(3), 890–944. <https://doi.org/10.1108/INTR-10-2021-0774>

- Sarrami-Foroushani, P., Travaglia, J., Debono, D., & Braithwaite, J. (2014). Implementing strategies in consumer and community engagement in health care: results of a large-scale, scoping meta-review. *BMC Health Services Research*, 14(1), 402. <https://doi.org/10.1186/1472-6963-14-402>
- Savira, F., Sanigorski, A., Namara, K. M., Hensher, M., Peeters, A., & Manias, E. (2024). Perspectives on telehealth implementation in Australia: An exploratory qualitative study with practice managers and general practitioners. *International Journal of Medical Informatics*, 188, 105473. <https://doi.org/10.1016/j.ijmedinf.2024.105473>
- Seifert, A., Batsis, J. A., & Smith, A. C. (2020). Telemedicine in Long-Term Care Facilities During and Beyond COVID-19: Challenges Caused by the Digital Divide. *Frontiers in Public Health*. <https://doi.org/10.3389/fpubh.2020.601595>
- Shachar, C., Engel, J., & Elwyn, G. (2020). Implications for Telehealth in a Postpandemic Future: Regulatory and Privacy Issues. In *JAMA - Journal of the American Medical Association*. <https://doi.org/10.1001/jama.2020.7943>
- Shafik, W. (2024). *Data-Driven Future Trends and Innovation in Telemedicine* (pp. 93–118). <https://doi.org/10.4018/979-8-3693-2141-6.ch005>
- Shaver, J. (2022). The State of Telehealth Before and After the COVID-19 Pandemic. *Primary Care: Clinics in Office Practice*, 49(4), 517–530. <https://doi.org/10.1016/j.pop.2022.04.002>
- Shirke, M. M., Shaikh, S. A., & Harky, A. (2020). Implications of telemedicine in oncology during the COVID-19 pandemic. In *Acta Biomedica*. <https://doi.org/10.23750/abm.v91i3.9849>
- Sittig, D. F., Wright, A., Coiera, E., Magrabi, F., Ratwani, R., Bates, D. W., & Singh, H. (2020). Current challenges in health information technology–related patient safety. *Health Informatics Journal*. <https://doi.org/10.1177/1460458218814893>

- Snoswell, C. L., North, J. B., & Caffery, L. J. (2020). Economic Advantages of Telehealth and Virtual Health Practitioners: Return on Investment Analysis. *JMIR Perioperative Medicine*. <https://doi.org/10.2196/15688>
- Song, J., Sawang, S., Drennan, J., & Andrews, L. (2015). Same but different? Mobile technology adoption in China. *Information Technology & People*, 28(1), 107–132. <https://doi.org/10.1108/ITP-10-2013-0187>
- Sonkamble, R. G., Shirke-Deshmukh, S., Katkar, V., Lunagaria, M., Tejani, G. G., & Mousavirad, S. J. (2025). A blockchain secured metaverse framework for scalable and immersive telemedicine. *Scientific Reports*, 15(1), 26655. <https://doi.org/10.1038/s41598-025-12068-6>
- Sounderajah, V., Patel, V., Varatharajan, L., Harling, L., Normahani, P., Symons, J., Barlow, J., Darzi, A., & Ashrafian, H. (2021). Are disruptive innovations recognised in the healthcare literature? A systematic review. *BMJ Innovations*, 7(1), 208–216. <https://doi.org/10.1136/bmjinnov-2020-000424>
- Statistics, L. (2015). *Mann-Whitney U test using SPSS Statistics*. Statistical Tutorials and Software Guides. <https://statistics.laerd.com/spss-tutorials/mann-whitney-u-test-using-spss-statistics.php>
- Su, Z., Li, C., Fu, H., Wang, L., Wu, M., & Feng, X. (2024). Development and prospect of telemedicine. *Intelligent Medicine*, 4(1), 1–9. <https://doi.org/10.1016/j.imed.2022.10.004>
- Summerfelt, W. T., Sulo, S., Robinson, A., Chess, D., & Catanzano, K. (2015). Scalable hospital at home with virtual physician visits: Pilot study. *American Journal of Managed Care*, 21(10), 675–684. <https://pubmed.ncbi.nlm.nih.gov/26633092/>
- Sunny, M. N. M., Sumaiya, U., Akter, M. H., Kabir, F., Munmun, Z. S., Nurani, B., Atayeva, J., & Amin, M. M. (2024). Telemedicine and Remote Healthcare: Bridging

- the Digital Divide. *South Eastern European Journal of Public Health*, 1500–1510. <https://doi.org/10.70135/seejph.vi.2920>
- Taherdoost, H., & Keshavarzsaleh, A. (2016). Critical Factors that Lead to Projects' Success/Failure in Global Marketplace. *Procedia Technology*, 22, 1066–1075. <https://doi.org/10.1016/j.protcy.2016.01.151>
- Teece, D. J. (2018). Business models and dynamic capabilities. *Long Range Planning*. <https://doi.org/10.1016/j.lrp.2017.06.007>
- Terry, O. (2020). Disruption Innovation and Theory. *Journal of Service Science and Management*, 13(03), 449–458. <https://doi.org/10.4236/jssm.2020.133030>
- Thompson, M., & Walker, O. (2025). *Global Integration of AI-Powered Telemedicine: Innovations, Challenges, and the Future of Healthcare Delivery*. <https://doi.org/10.20944/preprints202505.0323.v1>
- Topol, E. J., Steinhubl, S. R., & Torkamani, A. (2015). Digital Medical Tools and Sensors. *JAMA*, 313(4). <https://doi.org/10.1001/jama.2014.17125>
- Totten, A. M., Womack, D. M., Eden, K. B., McDonagh, M. S., Griffin, J. C., Grusing, S., & Hersh, W. R. (2016). *Telehealth: mapping the evidence for patient outcomes from systematic reviews*. <https://pubmed.ncbi.nlm.nih.gov/27536752/>
- Tuckson, R. V., Edmunds, M., & Hodgkins, M. L. (2017). Telehealth. *New England Journal of Medicine*, 377(16), 1585–1592. <https://doi.org/10.1056/NEJMSr1503323>
- Union, I. T. (2023). *Accessibility of Telehealth Services: Recommendation ITU-T F.780.2 (04/23) (v2 ed.)*. <https://www.itu.int/epublications/es/publication/itu-t-f-780-2-v2-2023-04-accessibility-of-telehealth-services/en>
- Velayati, F., Ayatollahi, H., Hemmat, M., & Dehghan, R. (2022). Telehealth Business Models and Their Components: Systematic Review. *Journal of Medical Internet Research*, 24(3), e33128. <https://doi.org/10.2196/33128>

- Venkatesh, Morris, Davis, & Davis. (2003). User Acceptance of Information Technology: Toward a Unified View. *MIS Quarterly*, 27(3), 425. <https://doi.org/10.2307/30036540>
- Venkatesh, V., & Bala, H. (2008). Technology Acceptance Model 3 and a Research Agenda on Interventions. *Decision Sciences*, 39(2), 273–315. <https://doi.org/10.1111/j.1540-5915.2008.00192.x>
- Venkatesh, V., Thong, J. Y. L., & Xu, X. (2016). Unified theory of acceptance and use of technology: A synthesis and the road ahead. *Journal of the Association for Information Systems*. <https://doi.org/10.17705/1jais.00428>
- Verhoef, P. C., Broekhuizen, T., Bart, Y., Bhattacharya, A., Dong, J. Q., Fabian, N., & Haenlein, M. (2021). Digital transformation: A multidisciplinary reflection and research agenda. *Journal of Business Research*, 122, 889–901. <https://doi.org/10.1016/j.jbusres.2019.09.022>
- Vystropova, O., & Bass, E. (2024). Communicative Component as Part of Professional Activity of Telehealth Physicians. *Virtual Communication and Social Networks*, 145–152. <https://doi.org/10.21603/2782-4799-2024-3-2-145-152>
- Wang, B., Yang, P., Zhang, L., Zhang, Y., & Li, X. (2024). Evaluating the Effectiveness of Remote Mobile Health Services for the Elderly: Standards and Best Practices. *International Journal of Interactive Mobile Technologies (IJIM)*, 18(23), 125–137. <https://doi.org/10.3991/ijim.v18i23.51311>
- Wells, M. J., Dukarm, P., Mills, A., & Bokinsky, R. L. (2025). Telehealth. In *The Oxford Handbook of Rehabilitation Psychology* (pp. 1227–1258). Oxford University Press. <https://doi.org/10.1093/oxfordhpb/9780197525456.013.0066>
- Werf, M. Vander, Bernard, J., Barta, D. T., Berg, J., Collins, T., Dowdy, M., Feiler, K., Moore, D. L., Sifri, C., Spargo, G., Taylor, C. W., Towle, C. B., & Wibberly, K. H. (2022). Pandemic Action Plan Policy and Regulatory Summary Telehealth Policy and

- Regulatory Considerations During a Pandemic. *Telemedicine and E-Health*, 28(4), 457–466. <https://doi.org/10.1089/tmj.2021.0216>
- Whitten, P., & Love, B. (2005). Patient and provider satisfaction with the use of telemedicine: overview and rationale for cautious enthusiasm. *Journal of Postgraduate Medicine*, 51(4), 294–300. <https://pubmed.ncbi.nlm.nih.gov/16388172/>
- Wickham, R. J. (2019). Secondary Analysis Research. *Journal of the Advanced Practitioner in Oncology*, 10(4). <https://doi.org/10.6004/jadpro.2019.10.4.7>
- Wootton, R. (2012). Twenty years of telemedicine in chronic disease management-an evidence synthesis. In *Journal of Telemedicine and Telecare*. <https://doi.org/10.1258/jtt.2012.120219>
- Wosik, J., Fudim, M., Cameron, B., Gellad, Z. F., Cho, A., Phinney, D., Curtis, S., Roman, M., Poon, E. G., Ferranti, J., Katz, J. N., & Tcheng, J. (2020). Telehealth transformation: COVID-19 and the rise of virtual care. In *Journal of the American Medical Informatics Association*. <https://doi.org/10.1093/jamia/ocaa067>
- Yang, R., Gao, S., & Jiang, Y. (2024). Digital divide as a determinant of health in the U.S. older adults: prevalence, trends, and risk factors. *BMC Geriatrics*, 24(1), 1027. <https://doi.org/10.1186/s12877-024-05612-y>
- Yang, Y. T. (2016). *Telehealth Parity Laws*. <https://doi.org/10.1377/hpb20160815.244795>
- Yawar, R. (2023). Microlearning-Supported Telehealth Approach for Addressing Developmental Disorders in Children: Challenges and Opportunities in Pakistan. *Global Journal of Intellectual & Developmental Disabilities*, 11(5). <https://doi.org/10.19080/GJIDD.2023.11.555821>
- Ye, J., He, L., & Beestrum, M. (2023). Implications for implementation and adoption of telehealth in developing countries: a systematic review of China's practices and experiences. *Npj Digital Medicine*, 6(1), 174. <https://doi.org/10.1038/s41746-023->

00908-6

- Yusifov, F. F., & Akhundova, N. E. (2022). Analysis of Demographic Characteristics Based on E-Demography Data. *Demography and Social Economy*. <https://doi.org/10.15407/dse2022.01.038>
- Zaineldeen, S., Hongbo, L., Koffi, A. L., & Hassan, B. M. A. (2020). Technology acceptance model' concepts, contribution, limitation, and adoption in education. *Universal Journal of Educational Research*, 8(11), 5061–5071. <https://doi.org/10.13189/ujer.2020.081106>
- Zhang, Y., Chandra, S., Peña, M. T., Lal, L., Summers, R. L., & Swint, J. M. (2023). Framework for Evaluating and Developing Sustainable Telehealth Programs. *Telemedicine and E-Health*, 29(9), 1421–1425. <https://doi.org/10.1089/tmj.2022.0407>
- Zhao, Y., Ni, Q., & Zhou, R. (2018). What factors influence the mobile health service adoption? A meta-analysis and the moderating role of age. *International Journal of Information Management*, 43, 342–350. <https://doi.org/10.1016/j.ijinfomgt.2017.08.006>
- Zhou, X., Snoswell, C. L., Harding, L. E., Bambling, M., Edirippulige, S., Bai, X., & Smith, A. C. (2020). The Role of Telehealth in Reducing the Mental Health Burden from COVID-19. *Telemedicine and E-Health*, 26(4), 377–379. <https://doi.org/10.1089/tmj.2020.0068>

APPENDIX A:

Q.1 If you have used telehealth services, how satisfied were you with the experience?

- Very Satisfied
- Somewhat Satisfied
- Neutral
- Somewhat Dissatisfied
- Very Dissatisfied

Q.2 What is the main reason you use telehealth services?

- Convenience
- Cost-effectiveness
- Better access to healthcare services
- Better communication with healthcare providers
- Other

Q.3 How do you feel about the quality of care received through telehealth services compared to in-person care?

- Telehealth services provide the same quality of care as in-person care
- Telehealth services provide better quality of care than in-person care
- Telehealth services provide worse quality of care than in-person care
- Not sure

Q.4 Have you ever experienced technical difficulties during a telehealth appointment?

- Yes
- No

Q.5 If you answered yes to above question, how satisfied were you with the support you received to resolve the technical difficulties?

- Very Satisfied

- Somewhat Satisfied
- Neutral
- Somewhat Dissatisfied
- Very Dissatisfied

Q.6 In your opinion, what are the advantages of telehealth services?

- Increased access to healthcare services
- Cost-effectiveness
- Convenience
- Reduced travel time and expenses
- Better communication with healthcare providers
- Other

Q.7 In your opinion, what are the disadvantages of telehealth services?

- Technical difficulties
- Limited physical examination options
- Reduced personal interaction with healthcare providers
- Difficulty with diagnosis and treatment of complex health issues
- Other

Q.8 Would you be willing to use telehealth services again in the future?

- Yes
- No

APPENDIX B: DATASET

	A	B	C	D	E	F	G	H	I
1	Have you	If you have u	What is the	How do you	Have you ever	If you answer	In your opinion, what	In your opinion, what are th	Would you be willi
2	1	4	1	2	1	4	Increased access to health	Technical difficulties	
3	1	5	1	1	1	5	Increased access to health	Technical difficulties	
4	1	4	3	2	2	5	Increased access to health	Technical difficulties; Limited p	
5	1	4	1	1	2	3	Increased access to health	Reduced personal interaction	
6	1	5	4	1	1	5	Increased access to health	Other	
7	1	4	2	3	1	3	Convenience; Reduced tra	Technical difficulties; Limited p	
8	1	4	2	2	2	4	Cost-effectiveness	Limited physical examination	
9	1	5	1	1	1	3	Increased access to health	Technical difficulties; Limited p	
10	1	5	3	2	2	5	Increased access to health	Limited physical examination	
11	1	4	1	1	2	5	Increased access to health	Limited physical examination	
12	2	3	1	3	2	3	Reduced travel time and	Difficulty with diagnosis and tre	
13	1	4	1	1	2	3	Reduced travel time and	Difficulty with diagnosis and tre	
14	2	3	5	4	2	3	Other	Other	
15	2	4	3	4	2	4	Increased access to health	Limited physical examination	
16	2	5	5	4	2	5	Other	Other	
17	1	5	3	2	2	5	Increased access to health	Reduced personal interaction	
18	1	4	1	2	2	3	Increased access to health	Difficulty with diagnosis and tre	
19	1	4	1	1	1	2	Increased access to health	Difficulty with diagnosis and tre	
20	1	3	5	2	1	3	Increased access to health	Limited physical examination	
21	2	5	3	1	2	5	Increased access to health	Limited physical examination	
22	1	5	2	1	2	3	Increased access to health	Limited physical examination	
23	1	5	1	1	2	5	Increased access to health	Difficulty with diagnosis and tre	
24	1	5	2	1	2	5	Increased access to health	Limited physical examination	
25	1	4	1	2	2	3	Convenience	Reduced personal interaction	
26	2	5	3	2	2	5	Increased access to health	Technical difficulties; Difficulty	
27	1	5	2	2	2	5	Increased access to health	Other	
28	1	5	4	2	2	3	Increased access to health	Other	
29	2	4	1	4	2	3	Increased access to health	Limited physical examination	
30	1	5	1	1	2	5	Convenience; Reduced tra	Technical difficulties; Limited p	
31	1	3	1	1	2	5	Increased access to health	Technical difficulties; Limited p	
32	2	4	4	1	2	4	Increased access to health	Limited physical examination	
33	2	3	5	4	1	2	Increased access to health	Technical difficulties; Limited p	
34	1	5	4	1	2	4	Better communication wi	Difficulty with diagnosis and tre	
35	1	5	1	1	2	5	Increased access to health	Limited physical examination	
36	1	5	1	3	1	4	Increased access to health	Technical difficulties; Limited p	
37	1	4	3	1	1	4	Reduced travel time and	Limited physical examination	
38	1	5	2	2	1	5	Increased access to health	Reduced personal interaction	
39	1	5	4	1	2	5	Increased access to health	Other	
40	1	5	4	1	2	5	Reduced travel time and	Difficulty with diagnosis and tre	
41	1	3	1	4	1	3	Increased access to health	Difficulty with diagnosis and tre	
42	1	4	3	2	1	4	Increased access to health	Technical difficulties; Limited p	
43	1	5	3	1	1	5	Increased access to health	Technical difficulties	
44	1	5	1	1	2	4	Increased access to health	Technical difficulties; Limited p	
45	1	5	1	1	2	5	Cost-effectiveness	Other	
46	2	4	1	2	1	4	Cost-effectiveness	Technical difficulties	
47	1	5	4	2	2	5	Increased access to health	Limited physical examination	
48	2	3	5	4	2	3	Increased access to health	Limited physical examination	
49	2	3	1	1	2	3	Convenience	Technical difficulties	
50	2	3	4	2	2	3	Convenience	Other	
51	1	3	1	1	1	2	Cost-effectiveness; Conve	Technical difficulties; Limited p	
52	1	4	1	2	1	3	Increased access to health	Limited physical examination	